

New synonymy and new data on the distribution of the mayflies from Korea and the Russian Far East (Ephemeroptera)

Y.J. Bae, N.Ju. Kluge & D.J. Chun

Bae, Y.J., Kluge, N.Ju. & Chun, D.J. 1998. New synonymy and new data on the distribution of the mayflies from Korea and the Russian Far East (Ephemeroptera). *Zoosystematica Rossica*, 7(1): 89-94.

Based on comprehensive examination of type specimens of Northeast Asian mayflies, the following new synonymy is established: *Ecdyonurus bajkova* Kluge, 1986 = *E. sub-spinosus* (Braasch & Soldan, 1988), syn. n.; *Epeorus curvatulus* Matsumura, 1931 = *E. anatolii* Sinitshenkova, 1981, syn. n.; *E. (Iron) aesculus* Imanishi, 1934 = *Iron koreanicus* Braasch & Soldan, 1988, syn. n.; *Ephemerella (Cincticostella) levnidovae* Tshernova, 1952 = *E. (C.) castanea* Allen, 1971, syn. n.; *E. (Ephemerella) dentata* Bajkova, 1967 = *E. (E.) keijoensis* Allen, 1971, syn. n.; *E. (E.) kozhovi* Bajkova, 1967 = *E. (E.) notofascia* Yoon & Bae, 1988, syn. n. Taxonomic discussions on some problematic species and new distributional data are provided.

Y.J. Bae, Department of Biology, Seoul Women's University, 126 Kongnung-dong, Nowon-gu, Seoul 139-774, Korea.

N.Ju. Kluge, Department of Entomology, Biological Faculty, St.Petersburg State University, Universitetskaya nab. 7/9, St.Petersburg 199034, Russia.

D.J. Chun, Zoological Institute, Russian Academy of Sciences, Universitetskaya nab. 1, St.Petersburg 199034, Russia.

Introduction

About one hundred and fifty species of mayflies occur in the Far East of Russia which is close to two thirds of all known Northeast Asian mayflies (Tshernova et al., 1986; Bae, 1997). Tshernova, Bajkova, Sinitshenkova, Kluge, and Tiunova mainly contributed to the knowledge of mayfly fauna of the Far East of Russia. On the other hand, sixty-six species of mayflies have been reported from Korea since Imanishi (1940) (Bae et al., 1994; Bae, 1997). The North Korean mayfly fauna has been thoroughly investigated recently (Braasch & Soldan, 1988; Bae & Soldan, 1997; Bae & Andrikovics, 1997).

Although geographically adjacent, mayflies from Korea and the Far East of Russia have been scarcely treated together until 1980s because of the communication problems between those countries. For this study, we intensively examined type and

non-type material from Korea and the Far East of Russia. We deal herein with new synonymy, taxonomic discussions on some problematic species, and new distributional data.

Material

Mayflies from the Far East of Russia (FE Russia), North Korea (N Korea), and South Korea (S Korea) were examined. For detailed information on type material deposited at Zoological Institute in St.Petersburg (ZIN), see Kluge (1995). For numerous non-type specimens (many of them reared) deposited at St.Petersburg State University (SPbU), see various Kluge's previous papers. For type and non-type material deposited at Seoul Women's University (SWU), see Bae et al. (1994) and Bae's various previous papers.

Taxonomic account

Family BAETIDAE

Baetis (Baetis) fuscatus (Linnaeus, 1761)

Baetis nla: Imanishi, 1940: 221 (larva; N Korea, Japan, Manchuria); Yoon & Bae, 1988a: 111 (S Korea).

Baetis fuscatus (L.): Müller-Liebenau, 1969: 128 (larva, imago; Europe); Kluge, 1980: 562 (Siberia); Tshernova et al., 1986: 133 (imago; FE Russia); Bae & Soldan, 1997 (N Korea).

Baetis (Baetis) fuscatus (L.): Novikova & Kluge, 1987: 8; Kluge, 1997: 194 (larva; Palaearctic).

Material examined. Numerous larvae, imagines, and imagines reared from larvae (Europe, Siberia, FE Russia, N Korea, S Korea; SPbU, SWU).

Imanishi (1940) described larvae of this species as "*Baetis nla*" from North Korea, Japan, and Manchuria. Since then, there have been many faunistic studies in Korea listing this species under the arbitrary name "*Baetis nla*" which refers to any *fuscatus*-type baetid (see Yoon & Bae, 1988a). This is a transpalaearctic species known from Europe to East Asia. The form distributed in Siberia and the Far East differs from that of Europe by orange colour of turban eyes of male imago (Kluge, 1980). It is a very common species in the Far East of Russia and in Korea occurring from upper to down stream areas, sometimes in polluted streams.

Baetis (Nigrobaetis) muticus (Linnaeus, 1758)

Baetis muticus (L.): Müller-Liebenau, 1969: 180 (larva, imago; Europe).

Baetis KUa: Yoon & Bae, 1988a: 111 (S Korea).

Baetis (Nigrobaetis) muticus (L.): Novikova & Kluge, 1987: 10; Novikova & Kluge, 1994: 635 (larva, imago; Europe, East Kazakhstan); Kluge, 1997: 190 (larva); Bae & Soldan, 1997 (N Korea).

Alainites muticus (L.): Waltz et al., 1994: 34 (Europe).

Material examined. Numerous larvae, imagines, and imagines reared from larvae from Europe, the Urals and Caucasus, one larva from Eastern Kazakhstan (SPbU). *Baetis* KUa: Yoon & Bae, 1988 (N Korea, S Korea; SWU).

This species has been known from Korea as "*Baetis* KUa" since Yoon & Bae (1988a). Based on examination of a good series of larval material from Europe and Asia, we determine this species as *Baetis (Nigrobaetis) muticus*. Formerly the easternmost point of recorded distribution of *B. (N.) muticus* was Eastern Kazakhstan (Novikova & Kluge, 1994).

Waltz et al. (1994) suggested a new classification for selected baetid groups, in which the genus *Alainites* Waltz & McCafferty, 1994 was established with the type species *Ephemerella mutica* Linnaeus, 1758 (wrongly spelled as "*Baetis muticus* L., 1758"). According to our investigation (Novikova & Kluge, 1994), the type species of *Alainites* is undoubtedly congeneric with the type species of *Takobia* Novikova & Kluge, 1987, and both can be placed in the subgenus (or genus) *Nigrobaetis* Kazlauskas in Novikova & Kluge, 1987 (see Kluge, 1997: 189).

Family HEPTAGENIIDAE

Ecdyonurus (Afghanurus) bajkova Kluge, 1986

Paracinygmula zhiltzovae Bajkova, 1975: 56 (nom. praecoc.) (larva; FE Russia).

Ecdyonurus zhiltzovae (Bajkova): Kluge, 1983: 31 (imago; FE Russia).

Ecdyonurus KUa: Yoon & Bae, 1984: 14 (larva; S Korea) (associated with *E. bajkova* Kluge by Bae et al., 1994).

Ecdyonurus bajkova Kluge in Tshernova et al., 1986: 117 (nomen novum pro *Paracinygmula zhiltzovae* Bajkova) (imago; FE Russia); Kluge, 1988: 300; Kluge, 1995: 19 (type deposition).

Nixe subspinosa Braasch & Soldan, 1988: 25 (imago; N Korea), *syn. n.*

Ecdyonurus subspinosa (Braasch & Soldan): Bae et al., 1994: 40 (S Korea).

Ecdyonurus (Afghanurus) bajkova Kluge: Kluge, 1997: 200 (larva; FE Russia).

Material examined. Holotype and paratypes (larvae) of *Paracinygmula zhiltzovae* Bajkova, 1975 = *Ecdyonurus bajkova* Kluge, 1986 (FE Russia; ZIN). Larvae, imagines and imagines reared from larvae (FE Russia, N Korea, S Korea; SPbU, SWU).

This species was originally described from larvae from the Far East of Russia (Bajkova, 1975) and Korea (Yoon & Bae, 1984). The adults were reared from larvae from the Far East of Russia and described by Kluge (1983). Soldan, on the other hand, collected the adult of the same species from North Korea in 1986 and described it under the name *Nixe subspinosa* Braasch & Soldan, 1988. Soldan also collected larvae and reared adults of *N. subspinosa* from North Korea (see Bae & Soldan, 1997).

This species was placed in the genus *Paracinygmula*, *Ecdyonurus*, or *Nixe*. The type species of *Paracinygmula* Bajkova, 1975 and the type species of *Nixe* Flowers, 1980 are undoubtedly congeneric with the type species of *Afghanurus* Demoulin, 1964

(Kluge, 1988), and all of them can be placed in the genus *Ecdyonurus* Eaton, 1868 s. l.

***Ecdyonurus (Afghanurus) joernensis* Bengtsson, 1909**

- Ecdyonurus joernensis* Bengtsson, 1909: 19 (female imago; Europe).
- Heptagenia mongolica* Bajkova & Varychanova, 1978: 111 (larva; Mongolia).
- Heptagenia dentata* Braasch, 1979: 69 (imago; Mongolia).
- Ecdyonurus mongolicus* (Bajkova & Varychanova): Kluge, 1980: 573 (larva, imago; Siberia); Bajkova, 1984: 116 (Mongolia).
- Ecdyonurus* KUb: Yoon & Bae, 1984: 14 (larva; S Korea).
- Ecdyonurus joernensis mongolicus* (Bajkova & Varychanova): Tshernova et al., 1986: 177 (imago; FE Russia); Kluge, 1988: 300; Bae et al., 1994: 39 (N Korea).
- Nixe mongolica* (Bajkova & Varychanova): Braasch & Soldan, 1988: 25 (N Korea).
- Ecdyonurus (Afghanurus) joernensis* Bengtsson: Kluge, 1997: 200 (larva; Scandinavia, Russia, Mongolia).

Material examined. Numerous larvae, imagines and imagines reared from larvae (Europe, the Urals, Siberia, Mongolia, FE Russia, S Korea; SPbU, SWU).

This is a transpalearctic species occurring from Europe to East Asia. On its systematic position see discussion above, under *Ecdyonurus (Afghanurus) bajkovae*; the usage of the generic name *Heptagenia* for this species was undoubtedly wrong.

***Epeorus curvatulus* Matsumura, 1931**

- Epeorus curvatulus* Matsumura, 1931: 1477 (Japan); Imanishi, 1934: 392 (imago, larva; Japan); Imanishi, 1940: 250 (larva; N Korea, S Korea, Manchuria); Braasch & Soldan, 1988: 27 (N Korea); Kluge, 1997: 205 (larva).
- Epeorus (Belovius) curvatulus* Matsumura: Tshernova, 1981: 326 (imago; Japan).
- Epeorus* (s. str.) *anatolii* Sinitshenkova, 1981: 814 (larva; FE Russia), *syn. n.*; Kluge, 1995: 19 (type deposition).
- Epeorus rautiani* Sinitshenkova, 1982: 52 (larva, imago; Siberia) (synonymized with *E. anatolii* by Tiunova, 1987: 7); Tshernova et al., 1986: 117 (imago; Siberia); Kluge, 1995: 27 (type deposition);

Material examined. Larvae and imagines (S Korea, FE Russia; SWU, SPbU). Holotypus (larva) of *E. anatolii* Sinitshenkova, 1981 (FE Russia; ZIN). Paratypes (larvae, imagines and imagines reared from larvae) of *E. rautiani* Sinitshenkova, 1982 (Siberia; ZIN);

In the original description of the subgenus *Belovius* Tshernova, 1981, this taxon was

characterized by imaginal features only; a number of unrelated species of which important larval characters were unknown were placed there, *E. curvatulus* among them. Sinitshenkova (1981) gave a diagnosis of the subgenus *Belovius* based on larval characters: each of tergalliae of the pairs II-VII has a wide proximal lobe separated from remainder of tergallia by posterior costa. With this diagnosis, *Belovius* became a natural taxon. According to this diagnosis, *E. curvatulus* must be placed not in *Belovius*, as each tergallia of *E. curvatulus* has the posterior costa on its posterior margin. Based on this character, Sinitshenkova described *E. anatolii* as belonging not to *Belovius*, but to the subgenus *Epeorus* s. str.

***Epeorus (Iron) aesculus* Imanishi, 1934**

- Epeorus aesculus* Imanishi, 1934: 384 (imago, larva partim; Japan); Imanishi, 1940: 250 (larva; N Korea).
- Iron aesculus* (Imanishi): Sinitshenkova, 1978: 50 (larva, imago; FE Russia); Tshernova et al., 1986: 120 (imago; FE Russia).
- Epeorus (Iron) aesculus* Imanishi: Kluge & Tiunova, 1989: 8 (larva, imago; FE Russia); Kluge, 1997: 206 (larva; FE Russia).
- Iron koreanicus* Braasch & Soldan, 1988: 25 (larva; N Korea), *syn. n.*

Material examined. Larvae, imagines and imagines reared from larvae (FE Russia, N Korea, S Korea; SPbU, SWU).

Originally *E. aesculus* was described from imagines and two forms of larvae tentatively attributed to this species (Imanishi, 1934). Sinitshenkova (1978) redescribed imagines and true larvae of *E. aesculus*, and distinguished it from *Iron maculatus* Tshernova, 1949. Kluge & Tiunova (1989) gave an additional description of this species. Soldan (pers. comm.) recently agreed that *Iron koreanicus* is conspecific with Kluge & Tiunova's (1989) concept of *E. (I.) aesculus*.

Family EPHEMERELLIDAE

***Ephemerella (Cincticostella) levanidovae* Tshernova, 1952**

- Ephemerella levanidovae* Tshernova, 1952: 274 (larva, FE Russia).
- Ephemerella orientalis* Tshernova, 1952: 279 (imago; FE Russia) (synonymy established by Tshernova et al., 1986: 138).
- Ephemerella (Cincticostella) levanidovae* Tshernova: Allen, 1971: 516 (larva, FE Russia); Tshernova et

al., 1986: 138 (imago; FE Russia); Kluge, 1995: 41 (type deposition).
Ephemerella (Cincticostella) castanea Allen, 1971: 514 (larva; S Korea), *syn. n.*
Cincticostella levanidovae (Tshernova): Tiunova, 1987: 9 (FE Russia).
Cincticostella castanea (Allen): Yoon & Bae, 1988b: 29 (larva, imago; S Korea).

Material examined. Lectotype and paralectotypes (larvae) of *Ephemerella levanidovae* Tshernova, 1952 (FE Russia; ZIN). Larvae, imagines, and imagines reared from larvae (FE Russia, S Korea; SPbU, SWU).

We recognize this synonymy based on a good series of reared material of *E. (C.) levanidovae* and *E. (C.) castanea* from the type localities in the Far East of Russia and in Korea respectively. This species commonly occurs in clean mountain streams.

Ephemerella (Drunella) solida Bajkova, 1980

Ephemerella trispina na: Imanishi, 1940: 193 (larva; N Korea).
Ephemerella solida Bajkova, 1980: 796 (larva; FE Russia).
Ephemerella (Drunella) solida (Bajkova): Tshernova et al., 1986: 140 (imago; FE Russia); Kluge, 1997: 210 (larva; FE Russia).
Drunella solida (Bajkova): Tiunova, 1978: 7 (FE Russia); 1988: 5 (imago; FE Russia).

Material examined. Larvae, imagines, and imagines reared from larvae (FE Russia, N Korea, S Korea; SPbU, SWU).

Previously the species described by Imanishi (1940) under arbitrary name "*Ephemerella trispina* na" was incorrectly associated with *Ephemerella triacantha* by Tshernova (1952). *E. (D.) solida* is distinguished from *E. (D.) triacantha* by the presence of stout setae on fore femora, lack of a longitudinal ridge on fore femora, and distinct colour pattern on thorax and abdomen. Previously, this species has been frequently misidentified as *E. (D.) triacantha* in faunistic studies in Korea.

Ephemerella (Drunella) triacantha Tshernova, 1949

Ephemerella trispina naa: Imanishi, 1940: 194 (larva; N Korea).
Ephemerella triacantha Tshernova, 1949: 151 (larva; Altai).
Ephemerella tenax Tshernova, 1952: 273 (larva; FE Russia) (synonymy established by Kluge, 1995: 43).
Ephemerella (Drunella) triacantha (Tshernova): Edmunds, 1959: 546; Tshernova et al., 1986: 139

(imago; FE Russia); Kluge, 1997: 210 (larva; FE Russia).

Ephemerella (Drunella) trispina Ueno: Yoon & Kim, 1981: 37 (S Korea).
Drunella triacantha (Tshernova): Yoon & Bae, 1988a: 166 (larva, imago; S Korea).

Material examined. Larvae, imagines, and imagines reared from larvae (FE Russia, N Korea, S Korea; PSU, SWU). Lectotype and paralectotype (larvae) of *E. tenax* Tshernova, 1952 (FE Russia; ZIN).

See discussion under *E. (D.) solida*, above.

Ephemerella (Ephemerella) dentata Bajkova, 1967

Ephemerella dentata Bajkova, 1967: 331 (larva; FE Russia); Kluge, 1995: 40 (type deposition).
Ephemerella (Ephemerella) keijoensis Allen, 1971: 526 (larva; S Korea), *syn. n.*
Ephemerella (Ephemerella) dentata Bajkova: Kluge, 1997: 212 (larva; FE Russia).

Material examined. Holotype (larva) of *E. dentata* Bajkova, 1967 (FE Russia; ZIN). Larvae (FE Russia; SPbU) and imagines reared from larvae (S Korea; SWU).

Based on the examination of type material of *E. dentata* and a good series of reared material of *E. (E.) keijoensis* from the type locality (Seoul, Korea), we established the above synonymy.

Ephemerella (Ephemerella) kozhovi Bajkova, 1967

Ephemerella nba: Imanishi, 1940: 202 (larva; N Korea) (associated with *E. notofascia* by Yoon & Bae, 1988b).
Ephemerella kozhovi Bajkova, 1967: 327 (larva, imago; FE Russia); Kluge, 1995: 40.
Ephemerella (Ephemerella) kozhovi Bajkova: Tshernova et al., 1986: 138 (imago; FE Russia); Kluge, 1997: 212 (larva; FE Russia).
Ephemerella notofascia Yoon & Bae, 1988b: 34 (larva; S Korea), *syn. n.*

Material examined. Holotype and paratypes (imagines and larvae) of *E. kozhovi* Bajkova, 1967 (FE Russia; ZIN). Holotype and paratypes (larvae) of *E. notofascia* Yoon & Bae, 1988 (S Korea; SWU). Larvae, imagines and imagines reared from larvae (FE Russia, S Korea; SPbU, SWU).

We establish the above synonymy by comparison of the type specimens of both species.

Acknowledgements

This work has been done during the senior author's research visit to the St. Petersburg State Uni-

versity in 1996 which was supported by Korean Science and Engineering Foundation # 961-0508-063-2. The work was fulfilled using scientific collections of the Zoological Institute, Russian Academy of Sciences, which obtain financial support from the Science and Technology Ministry of the Russian Federation (Reg. No. 98-03-16).

References

- Allen, R.K. 1971. New Asian *Ephemerella* with notes (Ephemeroptera: Ephemerellidae). *Can. Entomol.*, 103: 512-528.
- Bengtsson, S. 1909. Beiträge zur Kenntnis der paläarktischen Ephemeren. *Lunds Univ. Årsskrift, N. F., Afd. 2*, 2(5): 1-19.
- Bae, Y.J. 1997. A historical review of Ephemeroptera systematics in Northeast Asia. In: *Ephemeroptera & Plecoptera. Biology - Ecology - Systematics (Proc. VIII Intern. Conf. on Ephemeroptera and XII Intern. Symp. on Plecoptera, Aug. 1995, Losanne)*: 405-417. Fribourg/Switzerland.
- Bae, Y.J. & Andrikovics, S. 1997. Mayfly (Ephemeroptera) fauna of North Korea (2). *Insecta Koreana*, 14: 153-160.
- Bae, Y.J. & Soldan, T. 1997. Mayfly (Ephemeroptera) fauna of North Korea (1). *Insecta Koreana*, 14: 137-152.
- Bae, Y.J., Yoon, I.B. & Chun, D.J. 1994. A catalogue of the Ephemeroptera of Korea. *Entomol. Res. Bull., Korean entomol. Inst., Korea Univ. Seoul*, 20: 31-50.
- Bajkova, O.Ya. 1967. New species of *Ephemerella* Walsh (Ephemeroptera, Ephemerellidae) from the Far East and East Siberia. *Entomol. Obozr.*, 46(2): 327-335. (In Russian).
- Bajkova, O.Ya. 1975. A new genus of Ephemeroptera from the Primorye (Ephemeroptera: Heptageniidae). *Izv. Sibir. Otd. Akad. Nauk SSSR, Ser. biol. Nauk*, 1(5): 54-57. (In Russian).
- Bajkova, O.Ya. 1980. A new species of mayfly (Ephemeroptera, Ephemerellidae) from Primorye region. *Entomol. Obozr.*, 59(4): 796-799. (In Russian).
- Bajkova, O.Ya. 1984. Contributions to the knowledge of mayflies (Ephemeroptera) of North Mongolia. In: V. Landa et al. (eds.). *Proc. IV Intern. Conf. on Ephemeroptera*: 115-116. ČSAV. (In Russian).
- Bajkova, O.Ya. & Varychanova, K.V. 1978. Mayflies of Mongolia. In: *Prirodnye usloviya i resursy Prisubkhugulya. Trudy Sov.-mong. kompleks. Khubsugul. eksped.*, 6: 111-121. Irkutsk - Ulan Bator. (In Russian).
- Braasch, D. 1979. Die Eintagsfliegen (Ephemeroptera, Heptageniidae) der Mongolisch-Deutschen Biologischen Expeditionen 1964 und 1977. *Entomol. Nachr.*, 23(5): 65-76.
- Braasch, D. & Soldan, T. 1988. Heptageniidae aus Nordkorea (KVDR), nebst einigen Bemerkungen zu ihrem generischen Status (Insecta, Ephemeroptera). *Faun. Abh. staatl. Mus. Tierkd. Dresden*, 16(2): 23-28.
- Edmunds, G.F., jr. 1959. Subgeneric groups within the mayfly genus *Ephemerella* (Ephemeroptera: Ephemerellidae). *Ann. entomol. Soc. Amer.*, 52(5): 543-547.
- Imanishi, K. 1934. Mayflies from Japanese torrents. IV. Notes on the genus *Epeorus*. *Annot. zool. Japan.*, 14: 381-395.
- Imanishi, K. 1940. Ephemeroptera of Manchoukuo, Inner Mongolia, and Chosen. *Rep. limnol. Surv. Kwant. and Manch.*: 169-263. Kyoto. (In Japanese).
- Kluge, N.Ju. 1980. To the knowledge of mayflies (Ephemeroptera) of Taimyr national district. *Entomol. Obozr.*, 59(3): 561-579. (In Russian).
- Kluge, N.Ju. 1983. New and little-known mayflies of the USSR Far East. Genus *Ecdyonurus* (Ephemeroptera, Heptageniidae). In: *Ekologiya i sistematika presnovodnykh organizmov Dal'nego Vostoka* [Ecology and systematics of the freshwater organisms in the Far East]: 27-36. Vladivostok. (In Russian).
- Kluge, N.Ju. 1988. Revision of genera of the family Heptageniidae (Ephemeroptera). I. Diagnoses of tribes, genera and subgenera of the subfamily Heptageniinae. *Entomol. Obozr.*, 67(2): 291-313. (In Russian).
- Kluge, N.Ju. 1995. *A catalogue of the type-specimens in the collection of the Zoological Institute, Russian Academy of Sciences. Insecta, Ephemeroptera*. 52 pp. St.Petersburg. (In Russian).
- Kluge, N.Ju. 1997. Order mayflies. Ephemeroptera. In: S.Ya. Tsalolikhin (ed.). *Opredelitel' presnovodnykh bespozvonochnykh Rossii i sopredel'nykh stran* [Key to freshwater invertebrates of Russia and adjacent lands], 3: 176-220. St.Petersburg. (In Russian).
- Kluge, N.Ju. & Tjunova, T.M. 1989. Palearctic mayflies of the group *longimanus* of the subgenus *Iron* (Ephemeroptera, Heptageniidae). *Vestnik Zool.*, 1989(4): 7-14. (In Russian).
- Matsumura, S. 1931. *6000 Illustrated Insects of the Japanese Empire*. 191+1497 p. Tokyo. (In Japanese).
- Müller-Liebenau, I. 1969. Revision der europäischen Arten der Gattung *Baetis* Leach, 1815 (Insecta, Ephemeroptera). *Gewässer und Abwässer*, 48/49: 1-214.
- Novikova, E.A. & Kluge, N.Ju. 1987. Systematics of the genus *Baetis* (Ephemeroptera, Baetidae) with description of a new species from Middle Asia. *Vestnik Zool.*, 1987(4): 8-19. (In Russian).
- Novikova, E.A. & Kluge, N.Ju. 1994. Mayflies of the subgenus *Nigrobaetis* (Ephemeroptera, Baetidae). *Baetis*. *Entomol. Obozr.* 73(3): 623-644. (In Russian).
- Sinitshenkova, N.D. 1978. A redescription of larval and imaginal stages of *Iron aesculus* (Imanishi), 1934 and the evolution of the gill apparatus in the larvae of the g. *Iron* Eaton, 1881 (Ephemeroptera, Heptageniidae). *Bull. Mosk. Obshch. Ispyt. Prir.*, 83(5): 49-56. (In Russian).
- Sinitshenkova, N.D. 1981. To the knowledge of nymphs of the genus *Epeorus* Eaton (Ephemeroptera, Heptageniidae). *Entomol. Obozr.*, 60(4): 813-823. (In Russian).

- Sinitshenkova, N.D. 1983. A new species of mayflies *Epeorus rautiani* sp. n. from Transbaikal (Ephemeroptera, Heptageniidae). *Bull. Mosk. Obshch. Ispyt. Prir.*, 87(1): 52-58. (In Russian).
- Tiunova, T.M. 1987. *Mayflies from river Kedrovaya and their ecological and physiological characters*. Diss. Abstr., Zool. Inst. Acad. Sci. USSR, Leningrad. 25 p. (In Russian).
- Tiunova, T.M. 1988. New data on mayflies of the family Ephemerellidae (Ephemeroptera) of the Far East fauna. In: *Fauna, sistematika i biologiya presnovodnykh bespozvonochnykh* [Fauna, systematics and biology of freshwater invertebrates]: 3-6. Vladivostok. (In Russian).
- Tshernova, O.A. 1949. Mayfly nymphs from Lake Teletskoye and River Bija. *Trudy zool. Inst. Akad. Nauk SSSR*, 7: 139-158. (In Russian).
- Tshernova, O.A. 1952. Mayflies (Ephemeroptera) of the Amur river basin and adjacent waters and their role in the nutrition of Amur fishes. *Trudy Amur. ikhtiol. Eksped.*, 3: 229-360. (In Russian).
- Tshernova, O.A. 1981. On the systematics of adult mayflies of the genus *Epeorus* Eaton, 1881 (Ephemeroptera, Heptageniidae). *Entomol. Obozr.*, 60(2): 323-336. (In Russian).
- Tshernova, O.A., Kluge, N.Ju., Sinitshenkova, N.D. & Belov, V.V. 1986. 5. Order Ephemeroptera. In: Lehr, P.A. (ed.). *Opredelitel' nasekomykh Dal'nego Vostoka SSSR* [Keys to the insects of the Far East of the USSR], 1: 99-142. Leningrad. (In Russian).
- Waltz, R.D., McCafferty, W.P. & Thomas, A. 1994. Systematics of *Alainites* n. gen., *Dipheter*, *Indobaetis*, *Nigrobaetis* n. stat., and *Takovia* n. stat. (Ephemeroptera, Baetidae). *Bull. Soc. Hist. nat. Toulouse*, 130: 33-36.
- Yoon, I.B. & Bae, Y.J. 1984. The classification of the Heptageniidae (Ephemeroptera) in Korea. *Entomol. Res. Bull., Korean entomol. Inst., Korea Univ., Seoul*, 10: 1-34.
- Yoon, I.B. & Bae, Y.J. 1988a. I. Order Ephemeroptera. In: *Illustrated encyclopaedia of animals and plants of Korea*, vol. 30 [Animals (Aquatic Insects)]: 95-184. Ministry of Education, Korea. (In Korean).
- Yoon, I.B. & Bae, Y.J. 1988b. The classification of the Ephemerellidae (Ephemeroptera) in Korea. *Entomol. Res. Bull., Korean entomol. Inst., Korea Univ., Seoul*, 14: 23-44.
- Yoon, I.B. & Kim, M.L. 1981. A taxonomic study on the larvae of Ephemerellidae (Ephemeroptera) in Korea. *Entomol. Res. Bull., Korean entomol. Inst., Korea Univ., Seoul*, 8: 33-59. (In Korean).

Received 15 October 1997