## PRIVATE LIBRARY OF WILLIAM L. PETERS

## SYSTEMATIC CHANGES IN CERTAIN EPHEMEROPTERA STUDIED BY R. K. ALLEN

GEORGE F. EDMUNDS, JR.<sup>1</sup> AND CHAD M. MURVOSH<sup>2</sup> <sup>1</sup>Department of Biology, University of Utah, Salt Lake City, Utah 84112 <sup>2</sup>Department of Biological Sciences, University of Nevada, Las Vegas, Nevada 89154

Abstract.—Heptagenia bella (Allen & Cohen) is placed herein as Nixe bella, NEW COMBI-NATION. Serratella thailandensis Allen NEW SYNONYM is a junior objective synonym of Cincticostella gosei (Allen). The species Ephemerella (Dentatella) bartoni (Allen) is properly referred to as Eurylophella (Dentatella) bartoni Allen. Caurinella idahoensis Allen is confirmed to be a valid genus and species of uncertain affinities. The genus Vietnamella Tshernova is transferred from Ephemerellinae to Teloganodinae.

Key Words.-Insecta, Ephemeroptera, mayfly, taxonomy

Edmunds & Murvosh (1995) presented an obituary for the late Richard K. Allen. This paper discusses and corrects some systematic situations resulting from the taxonomic work of Allen, and detailed in an unpublished bibliography and list of taxa (GFE & CMM, unpublished), which is available from either of us. In most cases no comment is made on published name changes made by Allen or others.

Labels on Type Material. — It is essential for those subsequently studying types or paratypes of species described by R. K. Allen to note that in some cases the slides, at least, bear names that apparently were changed before publication, and that slides of holotypes or supplementary types are not always labeled as such. Usually there is a red margin on a slide label where the specimen is part of a holotype and a blue edge slide label designates a part from a paratype. This too is inconsistent in the collection. Localities and dates will allow these specimens to be associated with their correct names; in a few cases obvious types or supplementary types in vials or parts on slides are not labeled as such. There are also specimens or slides labeled as types or supplementary types that are manuscript names.

Systematics. — We have examined the types of Heptagenia bella Allen & Cohen (1977) of Mexico at the California Academy of Sciences and note that the characters place this species as Nixe bella, a genus described after the name bella was proposed. Allen (personal communication) was aware of the need for the generic reassignment. Other specimens of Nixe from farther south appear to belong to one or more undescribed but similar species.

Allen (1975:20) named Ephemerella (Cincticostella) gosei Allen based on Gose's figure 38 labeled Ephemerella TEB (Gose, 1969) based on specimens from Chanta Buri, Thailand, 20 Jun 1961. Allen (1980: 76) named Serratella thailandensis Allen, based on the same Ephemerella TEB (Gose, 1969) and raised Cincticostella to generic rank. Serratella thailandensis NEW SYNONYM is a junior objective synonym of Cincticostella gosei.

Allen & Murvosh (1983) and Allen (1990) place *Epeorus* (*Iron*) margarita Edmunds and Allen (1964) from the U. S. and Mexico in *Iron*, thus restoring

Vol. 71(3)

*Iron* as a full genus. Recognition of *Iron* as a genus is now common but the generic placement of some Asian species of the complex are still to be resolved and will certainly influence the final resolution of the taxonomy of the species of *Iron* and related taxa.

The unnamed species *Thraulodes* sp. F of Allen & Brusca (1978) (Central America) is a member of the genus *Farrodes*. The correct generic assignment was recognized by R. K. Allen and H. M. Savage (personal communications). *Farrodes* occurs from Argentina to Texas (see Davis 1991). *Thraulodes* sp. G from Mexico was recognizable by several mayfly specialists as belonging to the genus *Terpides*, but this has been indicated earlier by Savage (1987) on a distribution map of *Terpides*. Savage (personal communication) confirmed that this map was based in part on Allen's *Thraulodes* sp. G.

Allen (1977) described Ephemerella bartoni Allen and placed it in the subgenus Dannella. Ephemerella bartoni did not fit into the definitions and keys of any of the existing subgenera of the Ephemerellidae. McCafferty (1978) transferred bartoni from the subgenus Dannella to Eurylophella. Allen (1980) raised the subgenera of Ephemerella to generic rank, retained bartoni in Dannella and placed it in a new subgenus Dentatella. Gill 4 of bartoni does not cover the remaining gills as fully as in other species of Eurylophella and the elongation of abdominal segments 8 and 9 is relatively less than in other Eurylophella. Although Allen felt strongly that he was correct about the generic placement of bartoni, we feel certain, as did McCafferty, that bartoni is the most plesiomorphic species of Eurylophella. We concede that the examination of either males or mature eggs from females or mature larvae would be needed to provide overwhelming proof of the generic assignment of *bartoni*, but the evidence is adequate that *bartoni* is cladistically a member of Eurylophella and is informative because it is the least specialized larva of the genus (the adult is unknown). McCafferty and Wang (1994) placed Dentatella as a subgenus of Eurylophella.

When Allen (1984) described Caurinella idahoensis Allen as a new genus and species of Ephemerellidae, from a single specimen, he provided no figures. Interestingly, the type specimen was first sent to R. W. Baumann at Brigham Young University who noted the unique nature of the specimen and forwarded it to Allen. When Allen described the species, he had forgotten who had sent the specimen (Allen, personal communication). Some entomologists privately have expressed doubt concerning the validity of the genus. The caudal filaments are similar to those of Serratella, but this character is shared also by the Asian Uracanthella and Cincticostella and some populations of the European Ephemerella ignita (Poda). We have examined the type of Caurinella idahoensis and believe it to be a distinct valid genus. The characters of the larva suggest that *Caurinella* is not a specialized derivative of *Serratella*. When the adults are discovered, its systematic position should be clarified. We include here a lateral view of the diagnostic apical abdominal segments of the larvae (Fig. 1). Vincent F. Lee loaned us an additional specimen of C. idahoensis (IDAHO. VALLEY Co.: Eggers Cr., Trib. Silver Creek, 24 May 1978, R. C. Biggam) from the California Academy of Sciences, San Francisco. The type specimen and the Eggers Creek specimen are female larvae, roughly the same size and both have well developed wing pads that are not yet as dark as they would be if the specimens were ready to emerge.

In the bibliography of Allen's mayfly papers we noted that Henry (1995) was

158



Figure 1. Caurinella idahoensis. Lateral view of abdominal segment 7 to 10 of larva.

raising the subgenus *Neochoroterpes* (Allen, 1974) of *Choroterpes* to generic status; McCafferty et al. (1993) and Henry (1993) have since elevated *Neochoroterpes* to generic rank.

Allen (1980) placed the genus Vietnamella Tshernova (1972) (V. thani Tshernova) as a subgenus of *Cincticostella* Allen (1971), which it superficially resembles. Allen (1984) restored Vietnamella as a valid genus in a new subtribe Vietnamellae of the tribe Ephemerellini of Ephemerellinae. This created a new problem because the presence of gills on abdominal segments 2-7 in V. thani is a fundamental character of the subfamily Teloganodinae. Gills on abdominal segments 3-7 are basic traits of the Ephemerellinae, Ephemerellini, subtribe Ephemerellae, and gills on 4-7 the hallmark of Ephemerellini, subtribe Timpanogae. Thus, Allen gave no characters for distinguishing *Vietnamella* from the subfamily Teloganodinae. You & Su (1987) have described larvae and adult males and females of a second species of Vietnamella (V. dabishanensis) from China. The adult characters are those of Teloganodinae. The genus Vietnamella is herein transferred to Teloganodinae. The cladistic relationships with typical Teloganodes and a derived group of *Teloganodes* are not known but the very flat larvae with prominent spines on the head, pronotum and forefemora differ markedly from other Teloganodinae. The name Vietnamellini should be retained as a tribe, but in Teloganodinae. Thus Teloganodinae has two tribes, Vietnamellini and Teloganodini.

## ACKNOWLEDGMENT

We thank Janice G. and W. L. Peters for calling our attention to the synonymy of *Serratella thailandensis* Allen and for reviewing the manuscript; Vincent F. Lee (Calif. Acad. Sci. San Francisco) provided a specimen of *Caurinella idahoensis*; and W. P. McCafferty also kindly reviewed an earlier version of the manuscript.

## LITERATURE CITED

- Allen, R. K. 1971. New Asian *Ephemerella* with notes (Ephemeroptera: Ephemerellidae). Canad. Entomol., 103: 512-528.
- Allen, R. K. 1974. Neochoroterpes, a new subgenus of Choroterpes Eaton from North America. Canad. Entomol., 106: 161-168.
- Allen, R. K. 1975. *Ephemerella* (*Cincticostella*): A revision of the nymphal stages (Ephemeroptera: Ephemerellidae). Pan-Pacif. Entomol., 51: 16-22.

1995

- Allen, R. K. 1977. A review of *Ephemerella (Dannella)* and the description of a new species (Ephemeroptera: Ephemerellidae). Pan-Pacif. Entomol., 53: 215-217.
- Allen, R. K. 1980. Geographic distribution and reclassification of the subfamily Ephemerellinae (Ephemeroptera: Ephemerellidae). In Flannagan, J. F. & K. E. Marshall (eds.). Advances in ephemeroptera biology. Plenum, New York.
- Allen, R. K. 1984. A new classification of the family Ephemerellidae and the description of a new genus. Pan-Pacif. Entomol. 60: 245-247.
- Allen, R. K. 1990. The distribution of southwest North American mayfly genera (Ephemeroptera) in the Mexican transition zone. In Campbell, I. C. (ed.). Mayflies and stoneflies: life histories and biology. Kluwer, Dordrecht.
- Allen, R. K. & R. C. Brusca. 1978. Generic revisions of mayfly nymphs II. Thraulodes in North and Central America (Leptophlebiidae). Canad. Entomol., 110: 413-433.
- Allen, R. K. & S. D. Cohen. 1977. Mayflies (Ephemeroptera) of Mexico and Central America: new species, descriptions and records. Canad. Entomol., 109: 399-414.
- Allen, R. K. & C. M. Murvosh. 1983. Taxonomy and zoogeography of the mayflies (Ephemeroptera: Insecta) of Baja California. Ann. Entomol. Soc. Am., 76: 425–433.
- Davis, J. R. 1991. A new species of *Farrodes* (Ephemeroptera: Leptophlebiidae: Atalophlebiinae) from southern Texas. Proc. Entomol. Soc. Wash., 89: 407–416.
- Edmunds, G. F., Jr. & R. K. Allen. 1964. Rocky Mountain species of *Epeorus (Iron)*. Eaton. (Ephemeroptera: Heptageniidae). J. Kansas Entomol. Soc., 37: 275-288.
- Edmunds, G. F. Jr. & C. M. Murvosh. 1995. Obituary: Richard K. Allen 1925–1992. Pan-Pac. Entomol., 71: 1-3.
- Gose, K. 1969. Mayflies (Ephemeroptera) from Thailand. Nat. S W Asia 6: 125-136.
- Henry, B. C. 1993. A revision of *Neochoroterpes* (Ephemeroptera: Leptophlebiidae) new status. Trans. Am. Entomol. Soc., 119: 317-333.
- Henry, B. C. (in press). Phylogeny of Neochoroterpes (Leptophlebiidae: Atalophlebiinae). In Ciborowski, J. J. H. and L. Corkum (eds.). Current directions in ephemeroptera research. Canadian Scholars' Publishing, Inc., Toronto.
- McCafferty, W. P. 1978. A natural subgeneric classification of *Ephemerella bartoni* and related species (Ephemeroptera: Ephemerellidae). Great Lakes Entomol., 11: 209-216.
- McCafferty, W. P., R. S. Durfee & B. C. Kondratieff. 1993. Colorado mayflies (Ephemeroptera): an annotated inventory. Southwest. Natur., 38: 252–274.
- McCafferty, W. P. & T-Q. Wang. Phylogenetics and the classification of the *Timpanoga* complex (Ephemeroptera: Ephemerellidae). J. N. Am. Benthol. Soc., 13: 569-579.
- Savage, H. M. 1987. Biogeographical classification of the Neotropical Leptophlebiidae (Ephemeroptera) based upon geological centers of ancestral origin and ecology. Stud. Neotrop. Fauna and Environ., 22: 199-222.
- Tshernova, O. A. 1972. Some new species of mayflies from Asia (Ephemeroptera: Heptageniidae, Ephemerellidae). Entomol. Rev. (Engl. Transl. Entomol. Obozr.), 51: 604-617.
- You, Da-shou & Su, Cui-rong. 1987. A new Vietnamella from China. Acta Zootaxon. Sinica, 12: 176–180.