

SINS.

ENTOMOLOGISKE MEDDELELSE

UDGIVNE AF

ENTOMOLOGISK FORENING

VED

ALB. KLÖCKER.

2nd. Ser.

ANDEN RÆKKE.

TREDIE BIND. FEMTE HEFTET.

3 vol. 5th. No.

FEBRUAR 1910.

Indhold:

Mordellistena Engelharti. Eine neue Mordellide aus Dänemark. Beschrieben von J. Schilsky in Berlin.	pag. 255
Snyltere i Edderkoppeæg. Af J. P. Kryger.	— 257
Résumé. Notes and observations on some parasites on spiders' eggs. By J. P. Kryger.	— 280
Om Forekomst, Indsamling og Præparation af Staphyliner. Af Joh. P. Johansen. — 286	
Nye Bidrag til Fortegnelserne over Danmarks Neuropterer og Trichopterer. Af Esben Petersen, Silkeborg.	— 305
Description of a new species of Ephemerida from Denmark. By Esben Pe- tersen, Silkeborg.	— 313
Mindre Meddelelser.	— 315
Litteratur.	— 318

KJØBENHAVN.

ENTOMOLOGISK FORENINGS FORLAG.

HOVEDKOMMISSIONÆR: H. HAGERUPS BOGHANDEL.


1910.

Description of a new species of Ephemerida from Denmark.

By
Esben Petersen,
Silkeborg.

Rhithrogena Ussingi n. sp. ♂. The eyes above greyish brown, divided by a bluish band, black below. Notum light brown anteriorly, pitchy brown posteriorly. Abdomen above light brown with the joinings of the intermediate segments lighter. Segments 8—10 darker, and the last segment with two longitudinal black streaks. Venter of abdomen lighter than the dorsum and of a uniform colour. Wings hyaline. Costa and subcosta greyish yellow-brown; the other longitudinal nervures and the cross nervures brown. The marginal and submarginal areas slightly tinged with greyish yellow becoming darker in the pterostigmate region. Fore legs brown; knees darker, and tibiæ darker at the tip. Femoræ

with a distinct dark band. Femoræ and tarsi of intermediate and hind legs brownish yellow; tibiæ yellow. Femoræ with a longitudinal dark streak on the outside as well as on the inside. Limbs of forceps brown. Penis yellow. Setæ light greyish brown at the base, lighter towards the tip. At the base with hardly visible brownish joinings. L. 10 mm. V. 10 mm; Setæ 22 mm.



Forceps of Rh. Ussingi
(From below).

The description is made from examples preserved in alcohol.

Two males are found by the author at a small rivulet (Højen Bæk), near Vejle, 4. June 1906. I dedicate the species to my friend, Mr. H. Ussing, Randers, to whom I am much indebted for valuable additions to my collection of Neuropterous insects.
