



Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <https://www.tandfonline.com/loi/tmdn20>

The complete mitochondrial genome of *Epeorus herklotsi* (Ephemeroptera: Heptageniidae) and its phylogeny

Xin-Yan Gao, Shu-Sheng Zhang, Le-Ping Zhang, Dan-Na Yu, Jia-Yong Zhang & Hong-Yi Cheng

To cite this article: Xin-Yan Gao, Shu-Sheng Zhang, Le-Ping Zhang, Dan-Na Yu, Jia-Yong Zhang & Hong-Yi Cheng (2018) The complete mitochondrial genome of *Epeorus herklotsi* (Ephemeroptera: Heptageniidae) and its phylogeny, *Mitochondrial DNA Part B*, 3:1, 303-304, DOI: 10.1080/23802359.2018.1445482

To link to this article: <https://doi.org/10.1080/23802359.2018.1445482>



© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 28 Feb 2018.



Submit your article to this journal [↗](#)



Article views: 327



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 9 View citing articles [↗](#)

The complete mitochondrial genome of *Epeorus herklotsi* (Ephemeroptera: Heptageniidae) and its phylogeny

Xin-Yan Gao^a, Shu-Sheng Zhang^b, Le-Ping Zhang^a, Dan-Na Yu^{a,c}, Jia-Yong Zhang^{a,c} and Hong-Yi Cheng^{a,c}

^aCollege of Chemistry and Life Science, Zhejiang Normal University, Jinhua, Zhejiang Province, China; ^bZhejiang Wuyanling National Nature Reserve, Taishun, Zhejiang Province, China; ^cKey Lab of Wildlife Biotechnology, Conservation and Utilization of Zhejiang Province, Zhejiang Normal University, Jinhua, Zhejiang Province, China

ABSTRACT

The mitochondrial genome of *Epeorus herklotsi* (Ephemeroptera: Heptageniidae) is a circular molecule of 15,801 bp in length with a base composition of 32.7% A, 32.9% T, 21.5% C, 13.0% G, including extra *tRNA^{Met}* gene. The *IMQM* tRNA cluster is found in *E. herklotsi* as well as *Paraformura youi* and two species of *Epeorus* (KM244708, KJ493406), while the typical *IQM* tRNA cluster is found in *Paegniodes cupulatus*. In BI and ML phylogenetic trees, the monophyly of the families Heptageniidae, Baetidae, and Ephemerellidae are highly supported. *E. herklotsi* is a sister clade to *Epeorus* sp2. (KJ493406).

ARTICLE HISTORY

Received 7 February 2018
Accepted 13 February 2018

KEYWORDS

Ephemeroptera;
mitochondrial genome;
Epeorus herklotsi; phylogeny

The phylogenetic relationships of Ephemeroptera using morphological or molecular methods still exists disputes (Kristensen 1981; Ogden and Whiting 2003; Zhang et al. 2008; Simon and Hadrys 2013; Li et al. 2014; Misof et al. 2014). Thirteen complete mitochondrial genomes and six partial mitochondrial genomes of Ephemeroptera are available (Zhang et al. 2008; Li et al. 2014; Tang et al. 2014; Zhou et al. 2016). In the 19 known mayfly mitochondrial genomes, gene rearrangements are found in five species (e.g. *Paraformura youi*, *Epeorus* sp1., *Epeorus* sp2., *Siphuriscus chinensis*, *Alainites yixiani*), especially in Heptageniidae two types of *IQM* and *IMQM* tRNA cluster were found. Hence, we sequenced the mitochondrial genome of *Epeorus herklotsi* (Ephemeroptera: Heptageniidae) to analyse the characteristics of mitochondrial gene arrangement and to discuss the phylogenetic relationships within Ephemeroptera.

The samples of *E. herklotsi* identified by Dr. JY Zhang were collected in Lishui, Zhejiang province, China. All samples of *E. herklotsi* were stored in Lab of HY Cheng, College of Chemistry and Life Science, Zhejiang Normal University. And the total genomic of DNA was isolated from one leg of *E. herklotsi* using Ezup Column Animal Genomic DNA Purification Kit (Sangon Biotech Company, Shanghai, China). The conserved primers and modified primers for PCR amplification were designed according to Zhang et al. (2008). PCR products were sent to Sangon Biotech Company for sequencing with both strands.

The mt genome of *E. herklotsi* is a circular molecule of 15,801 bp in length, with the overall base composition of

32.7% A, 32.9% T, 21.5% C, 13.0% G, including extra *tRNA^{Met}* gene. The gene rearrangement with *IMQM* tRNA cluster was found in *E. herklotsi*, which is similar to *P. youi* (Zhang et al. 2008) and two species of *Epeorus* (KM244708, KJ493406) (Tang et al. 2014), while is different to *Paegniodes cupulatus* with the typical *IQM* tRNA cluster (Zhou et al. 2016).

Bayesian inference (BI) and maximum likelihood (ML) trees were constructed using the 13 PCGs from 20 species using *S. chinensis* (Li et al. 2014) as outgroup (Figure 1). Each alignment was performed by Gblock 0.91b (Castresana 2000) using default settings in order to select conserved regions of the nucleotide. BI and ML analysis were performed by MrBayes3.1.2 (Huelsenbeck and Ronquist 2001) and RAxML 8.2.0 (Stamatakis 2014), respectively. In BI and ML phylogenetic trees, the monophyly of the families Heptageniidae, Baetidae, and Ephemerellidae is highly supported (Figure 1), but the monophyly of Siphonuridae is failed. *Siphonurus* sp. (Siphonuridae) is a clade sister to *Ameletus* sp. (Ameletidae), however, *Siphonurus immanis* (Siphonuridae) is a sister clade to *Ephemerella orientalis* (Ephemeridae). The monophyly of *Epeorus* is highly supported and *E. herklotsi* is a sister clade to *Epeorus* sp2. (KJ493406). In addition, Baetidae (*Baetis* sp. + *Alainites yixiani*) is a sister clade to Teloganodidae which is not consist with view of Ogden and Whiting (2005). We found the long branch attraction existed in Baetidae, which may affect the phylogenetic relationship of Baetidae.




Figure 1. Phylogenetic tree of the relationships among 20 species of Ephemeroptera, including *E. herklotsi* based on the nucleotide dataset of the 13 mitochondrial protein-coding genes. The Bayesian posterior probability values and the maximum-likelihood bootstrap values are indicated above nodes. The GenBank numbers of all species are shown in the figure.

Acknowledgements

The authors would like to thank Ya-Jie Gao for aid in taxon sampling.

Nucleotide sequence accession number: The complete mitochondrial genome of *E. herklotsi* has been assigned the GenBank accession number MG870104.

Disclosure statement

The authors declare no conflict of interest.

Funding

This work was supported by Zhejiang Provincial Natural Science Foundation (Grant Y18C040006) and by the National Natural Science Foundation of China (Grant 31370042).

References

- Castresana J. 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol.* 17:540–552.
- Huelsenbeck JP, Ronquist F. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics.* 17:754–755.
- Kristensen NP. 1981. Phylogeny of insect orders. *Annu Rev Entomol.* 26:135–157.
- Li D, Qin JC, Zhou CF. 2014. The phylogeny of Ephemeroptera in Pterygota revealed by the mitochondrial genome of *Siphuriscus chinensis* (Hexapoda: Insecta). *Gene.* 545:132–140.
- Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, Frandsen PB, Ware J, Flouri T, Beutel RG. 2014. Phylogenomics resolves the timing and pattern of insect evolution. *Science.* 346:763–767.
- Ogden TH, Whiting MF. 2003. The problem with “the Paleoptera problem” sense and sensitivity. *Cladistics.* 19:432–442.
- Ogden TH, Whiting MF. 2005. Phylogeny of Ephemeroptera (mayflies) based on molecular evidence. *Mol Phylogenet Evol.* 37:625–643.
- Simon S, Hadrys H. 2013. A comparative analysis of complete mitochondrial genomes among Hexapoda. *Mol Phylogenet Evol.* 69:393–403.
- Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics.* 30:1312–1313.
- Tang M, Tan M, Meng G, Yang S, Su X. 2014. Multiplex sequencing of pooled mitochondrial genomes a crucial step toward biodiversity analysis using mito-metagenomics. *Nucleic Acids Res.* 42:e166.
- Zhang J, Zhou C, Gai Y, Song D, Zhou K. 2008. The complete mitochondrial genome of *Parafronurus youi* (Insecta: Ephemeroptera) and phylogenetic position of the Ephemeroptera. *Gene.* 424:18–24.
- Zhou D, Wang YY, Sun JZ, Han YK. 2016. The complete mitochondrial genome of *Paegniodes cupulatus* (Ephemeroptera: Heptageniidae). *Mitochondrial DNA A DNA Mapp Seq Anal.* 27:925–926.