

Y. J. Bae (Ed.). 2001. The 21st Century and Aquatic Entomology in East Asia. Proc. 1st Symp. AESEA.
Korean Soc. Aquatic Entomol., Korea. pp. 55-84.

A Checklist of Japanese Ephemeroptera

ISHIWATA Shin-ichi

Kanagawa Environmental Research Center
1-3-39 Shinomiya, Hiratsuka, Kanagawa 254-0014, Japan

Abstract. Thirteen families, 36 genera, and 140 species of Ephemeroptera were confirmed for the Japanese Islands (Hokkaido, Honshu, Shikoku, Kyushu, and Okinawa). Out of them, one genus and eight species were reported for the first time. The checklist incorporated 21 new synonyms (including one generic synonym), eight new combinations, and one new status. Five species were indicated as *nomina dubia*.

Key words. Ephemeroptera, checklist, Japan

Introduction

Considerable contributions to mayfly taxonomy in Japan were given by Gose's monograph (Gose, 1979a, 1979b, 1979c, 1979d, 1979e, 1980a, 1980b, 1980c, 1980d, 1980e, 1980f, 1980e), in which a total of 95 species and one subspecies were listed as Japanese mayflies. In the monograph, however, 25 out of 35 species and three out of five genera named by Matsumura (1904, 1911, 1931a, 1931b, 1933) were overlooked. On the other hand, a checklist of Japanese insects compiled by Y. Hirashima (Entomological Laboratory of Kyushu University, 1990), which listed 103 species of Ephemeroptera for the Japanese Islands (Hokkaido, Honshu, Shikoku, Kyushu and Okinawa), included 20 of Matsumura's species but lacked 23 species named by Gose (1985). Besides Makibayashi (1998) declared that the Gose's species are to be *nomina dubia* because there were neither an indication nor designation of types for the new taxa.

Most of the species established by Matsumura in his earlier publications (Matsumura, 1904, 1931a, 1933) are available since they were published with descriptions. However, some generic names are not available because of the lack of type designation (Hubberd, 1988, 1989). For establishing new species, Gose gave a diagnostic key to distinguish them from the other species (Gose, 1979c, 1980a, 1980b, 1980c, 1980d, 1980e, 1980f, 1985). Clearly, they are available enough since they satisfy Acts 9.1, 10.1, 11.1-5, and 13.1.1 of the International Code of Zoological Nomenclature (4th Ed.).

This paper lists up to 140 valid species of Japanese mayflies based on comprehensive literatures revived. The families are phylogenetically arranged according to McCafferty (1996) with an additional family rank for Dipteromimidae (Kluge *et al.* 1995). Within a family, genera and species are arranged alphabetically. Symbols used in the text are as follows: *: newly recorded species, [()]: geographical distribution, “ ”: arbitrary name in the original text, [!]: obvious error of spelling, and ?: ?-mark in the original literature.

The types in the present study are deposited in the following institutions: CERK (Center for Ecological Research, Kyoto University, Japan); HU (The collection in the Laboratory of Systematic Entomology, Hokkaido University, Japan); NU (Nara Women's University, Japan); INHS (Illinois Natural History Survey, USA).

Only following 5 species were listed as *nomina dubia* because there were no chance for them to be applied by the original names: i.e., *Ephoron nakamurae* Matsumura, 1941, *Ephemerella akiana* Sato, 1969, *Ephemerella gose* Okazaki 1984, *Ameletus towadensis* Matsumura 1931, and *Isonychia mitsukonis* Morishita, 1960. *Ephoron nakamurae* which had

already been known to be *nomen dubium* by Ishiwata (1996). The names *Ephemera akiana* and *Isonychia mitsukonis* by Sato (1969) and Morishita (1961) respectively, lacked any descriptions or types, thus I cannot verify those species. *Ephemerella gose* by Okazaki (1984) based on the egg stage, also cannot be verified. *Ameletus towadensis* cannot be verified from the original description and none of Matsumura's specimens of the species exist in his collection.

Checklist

Leptophlebiidae (Tobiiro-kagerou-ka)

Genus *Chiusanophlebia* Uéno, 1969 (Ryukyu-tobiiro-kagerou-zoku)

Chiusanophlebia asahinai Uéno, 1969 (Ryukyu-tobiiro-kagerou) [Japan (Kyushu: Amamioshima Is., Okinawa: Okinawa-honto, Ishigaki-jima Is.)]
asahinai Uéno, 1969 (*Chiusanophlebia*).

Genus *Choroterpes* Eaton, 1881 (Hime-Tobiiro-kagerou-zoku)

Subgenus *Euthraulus* Barnard, 1932 (Toyo-hime-tobiiro-kagerou-azoku)

Choroterpes altioculus Kluge 1984 (Hime-tobiiro-kagerou) [Japan (Honshu, Shikoku, Kyushu), China, Korea, Russia]
altioculus Kluge 1984 (*Choroterpes (Euthraulus)*).
trifurcata auct., nec Uéno, 1928 (*Choroterpes*).

Genus *Choroterpides* Ulmer, 1939 (Kuro-tobiiro-kagerou-zoku)

**Choroterpides nigella* Kang & Yang, 1994 (Kuro-tobiiro-kagerou) [Japan (Okinawa: Iriomote-jima Is., Ishigaki-jima Is.), Taiwan]
nigella Kang & Yang, 1994 b (*Choroterpides*).

Genus *Paraleptophlebia* Lestage, 1917 (Tobiiro-kagerou-zoku)

Paraleptophlebia japonica (Matsumura, 1931) COMB. N. (Nami-tobiiro-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, Russia]
japonicus Matsumura, 1931a (*Baetis*).
chocolata Imanishi, 1937b (*Paraleptophlebia*) SYN. N.; Kluge, 1997 (*Leptophlebia (Neoleptophlebia)*).
chocorata [!] Imanishi, 1940 (*Paraleptophlebia*).

Note: After examination of the type species of *Baetis japonicus* Matsumura, 1931 in the Matsumura Collection, HU, I found that the species could not belong to Baetidae but to Leptophlebiidae. Although I have not been successful in locating the holotype of *Paraleptophlebia chocolata* Imanishi, 1937, I found the male specimens identified to *P. chocolata* by Imanishi in the Imanishi Collection, CERK, and specimens which I collected in Hiragawa River, Shiga-ken, the type locality of *P. chocolata* are identical to *P. japonica* in the respect of size and morphology of genitalia. Therefore, I regard *P. chocolata* as a synonym of *P. japonica*.

Paraleptophlebia spinosa Uéno, 1931 (Toge-tobiiro-kagerou) [Japan (Honshu, Shikoku)]
spinosa Uéno, 1931b (*Paraleptophlebia*).

Paraleptophlebia westoni Imanishi, 1937 (Uesuton-tobiirou-kagerou) [Japan (Hokkaido, Honshu, Kyushu)]

(?) *cincta*: Uéno, 1928 (*Paraleptophlebia*); Imanishi, 1940 (*Paraleptophlebia*); Bae et al., 2000.

westoni Imanishi, 1937b (*Paraleptophlebia*).

"sp. n!" Hashimoto, 1955 (*Paraleptophlebia*).

cincta: Gose, 1962 (*Paraleptophlebia*) Misidentification.

"sp. PA" Gose, 1962 (*Paraleptophlebia*).

Genus *Thraulus* Eaton, 1881 (Togeera-kagerou-zoku)

**Thraulus fatuus* Kang & Yang, 1994 (Usuguro-togeera-kagerou) [Japan (Okinawa: Okinawa-honto), Taiwan]

fatuus Kang & Yang, 1994b (*Thraulus*).

Thraulus grandis Gose, 1980 (Oo-togeera-kagerou) [Japan (Honshu, Kyushu, Okinawa: Okinawa-honto)]

"sp." Horasawa, 1931 (*Thraulus*).

grandis Gose, 1980c (*Thraulus*).

"sp. 2" Ueno, 1995 (*Thraulus*).

**Thraulus macilentus* Kang & Yang, 1994 (Hime-togeera-kagerou) [Japan (Okinawa: Okinawa-honto), Taiwan]

macilentus Kang & Yang, 1994b (*Thraulus*).

Potamanthidae (Kawa-kagerou-ka)

Genus *Potamanthus* Pictet, 1843 (Kawa-kagerou-zoku)

Subgenus *Potamanthodes* Ulmer, 1920 (Kiilo-kawa-kagerou-azoku)

Potamanthus formosus Eaton, 1892 (Kiilo-kawa-kagerou) [Japan (Honshu, Shikoku, Kyushu), Myanmar, China, Korea, Malaysia, Thailand, Vietnam]

formosus Eaton, 1892 (*Potamanthus*); Uéno, 1969 (*Potamanthodes*).

luteus: Uéno, 1928 (*Potamanthus*) Misidentification.

iyonis Matsumura, 1931a (*Potamanthus*).

kamonis Imanishi, 1940 (*Potamanthus* (*Potamanthodes*)).

Note: In Japan this species was described at first under the name of *Potamanthus luteus* (Linnaeus, 1767) by Uéno (1928) from central Japan (Setagawa River and Tenryugawa River). Imanishi (1940) regarded that *P. luteus* reported by Uéno (1928) did not belong to the subgenus *Potamanthus*, but to *Potamanthodes* and described a new species under the name of *Potamanthus* (*Potamanthodes*) *kamonis* Imanishi, 1940. Although I have not been successful in examining Uéno's *P. luteus*, it is highly probable that nymphs of Uéno's *P. luteus* will eventually prove to be nymphs of *P. formosus*. Uéno (1969) reported adults of this species from Taiwan and synonymized *P. kamonis* with *P. formosus*, nevertheless the name *P. kamonis* has been used as a valid name incorrectly (Uéno, 1973; Gose, 1970, 1985) until the world revision of Potamanthidae given by Bae & McCafferty (1991). Gose (1985) stated without precise data that *P. formosus* was distributed from Hokkaido to Kyushu. My observations suggest that distribution of this species in Japan extends to northern part of Honshu (Aomori-ken) and not to

Hokkaido.

Subgenus *Potamanthus* Pictet, 1843 (Kawa-kagerou-azoku)

- **Potamanthus huoshanensis* Wu, 1987 (Oo-kawa-kagerou) [Japan (Honshu), China]
huoshanensis Wu, 1987 (*Potamanthus*).
luteus oriens Bae & McCafferty, 1991 (*Potamanthus (Potamanthus)*) Misidentification.
"sp." Kobayashi & Nishino, 1992 (*Potamanthus*).

Note: It has been believed for many years that a single species, *P. kamonis*, was widely distributed in Japan (Gose, 1985). However, Kobayashi & Nishino (1990) reported the adults of *Potamanthus* sp. collected on the shores of Lake Biwa by light trap. The adults were also collected from Yokkaichi-shi (Yokkaichi City Government, 1990) by me. Bae & McCafferty (1991) assigned *Potamanthus* sp. to *P. luteus* from the descriptive and collecting data. I have examined *Potamanthus* sp. and found that it is conspecific with *Potamanthus huoshanensis* Wu, 1987.

Ephemeridae (Mon-kagerou-ka)

Genus *Ephemera* Linnaeus, 1758 (Mon-kagerou-zoku)

- Ephemera formosana* Ulmer, 1919 (Taiwan-mon-kagerou) [Japan (Okinawa: Okinawa-honto, Iriomote-jima Is., Ishigaki-jima Is.), Taiwan]
formosana Ulmer, 1919 (*Ephemera*); Gose, 1981 (*Ephemera*).
cf. *formosana* Uéno (*Ephemera*).
"sp." Futai, 1963 (*Ephemera*).
japonica: Takara & Azuma, 1972 (*Ephemera*) Misidentification.
koshuensis [!]: Takara & Azuma, 1972 (*Ephemera*) Misidentification.

Ephemera japonica McLachlan, 1875 (Futasuji-mon-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), Russia]
japonica McLachlan, 1875 (*Ephemera*); Uéno, 1928 (*Ephemera*?).

Ephemera orientalis McLachlan, 1875 (Toyo-mon-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), China, Korea, Mongolia, Russia]
orientalis McLachlan, 1875 (*Ephemera*).
kuccharonis Matsumura, 1931a (*Ephemera*) SYN. N.
lineata: Uéno & Okamoto, 1932 (*Ephemera*) Misidentification.

Note: Based on type material examination of *E. kuccharonis* in the Matsumura Collection, HU, I recognized *E. kuccharonis* is a synonym of *E. orientalis*.

Ephemera strigata Eaton, 1892 (Mon-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), China, Korea, Mongolia, Russia]
strigata Eaton, 1892 (*Ephemera*).
kuwamayai Navás, 1919 (*Ephemera*) SYN. N.
iwatensis Matsumura, 1931a (*Ephemera*) SYN. N.
jezonica Matsumura, 1931a (*Ephemera*) SYN. N.

Note: Judging from the original description of *Ephemera kuwamayai* Navás (1919), the abdominal terga having triangle dark markings are identical to those of *Ephemera strigata* Eaton, 1892, as is wing coloration and collecting data. Based on type material examination of both *E. iwatensis* and *E. jezonica* in the Matsumura Collection, HU, I could find no significant

morphological differences between the two. These two species are identical to *E. strigata*. Therefore, I recognize these three species, *E. kuwamayai*, *E. iwatensis*, and *E. jezonica* as synonyms of *E. strigata*.

Ephemera akiana Sato, 1969 *nomen dubium*.
akiana Sato, 1969 (*Ephemera*) *nomen dubium*.

Polymitarcyidae (Shiroiro-kagerou-ka)

Genus *Ephoron* Williamson, 1802 (Shiroiro-kagerou-zoku)

Ephoron eophilum Ishiwata, 1996 (Akatsuki-shiro-kagerou) [Japan (Honshu)]
eophilum Ishiwata, 1996 (*Ephoron*).

Ephoron limnobium Ishiwata, 1996 (Biwako-shiro-kagerou) [Japan (Honshu)]
limnobium Ishiwata, 1996 (*Ephoron*).

Ephoron shigae (Takahashi, 1924) (Oo-shiro-kagerou) [Japan (Honshu, Shikoku, Kyushu),
Korea, Russia (Far East)]
shigae Takahashi, 1924 (*Polymitarsis* [!]); Uéno, 1931c (*Polymitarcis* [!]); Lestage, 1938
(*Polymitarcys*); Uéno, 1973 (*Ephoron*).
"sp." Uéno, 1930 (*Polymitarcys*); Uéno, 1931c (*Polymitarcis* [!]).
"sp. NI" Hashimoto, 1955 (*Polymitarcis* [!]).

Ephoron nakamurae Matsumura, 1941 *nomen dubium*.
nakamurae Matsumura, 1941 (*Ephoron*) *nomen dubium*.

Caenidae (Himeshoro-kagerou-ka)

Genus *Brachycercus* Curtis, 1834 (Mitsutoge-hime-kagerou-zoku)

Brachycercus japonicus Gose, 1980 (Mitsutoge-hime-kagerou) [Japan (Honshu)]
japonica [!] Gose, 1980f (*Brachycercus*).
japonicus: Bae, 1997(*Brachycercus*).

Genus *Caenis* Stephens, 1835 (Himeshoro-kagerou-zoku)

Caenis horaria (Linnaeus) 1758 (Himeshoro-kagerou) [Japan (Honshu)]
horaria Linnaeus, 1758 (*Ephemera*); Uéno, 1950 (*Caenis*).
dimidiata Stephens, 1835 (*Caenis*).
lactea sensu Pictet, 1843 (*Caenis*).
lactella Eaton, 1884 (*Caenis*).

Caenis nishinoae Malzacher, 1996 (Biwako-himeshoro-kagerou) [Japan (Honshu), China,
Korea].
nishinoae Malzacher, 1996 (*Caenis*).

Ephemerellidae (Madara-kagerou-ka)

Genus *Cincticostella* Allen, 1971 (Toyo-madara-kagerou-zoku)

Cincticostella levanidovae (Tscheranova, 1952) (Kasutanea-madara-kagerou) [Japan (Kyushu: Tsushima Is.), China, Korea, Russia]
"sp. nay" Imanishi, 1940 (*Ephemerella*).
levanidovae Tscheranova, 1952 (*Ephemerella*); Edmunds, 1959 (*Ephemerella (Serratella)*); Allen, 1971 (*Ephemerella (Cincticostella)*); Ishiwata, 1993 (*Cincticostella*).
orientalis Tscheranova, 1952 (*Ephemerella*); Bajkova, 1979 (*Ephemerella (Cincticostella)*); Edmunds, 1959 (*Ephemerella (Serratella)*).
castanea Allen, 1971 (*Ephemerella (Cincticostella)*); Ishiwata, 1987 (*Cincticostella*).
delicata Allen, 1971 (*Ephemerella (Cincticostella)*).
costanea [!]: Gose, 1980d (*Ephemerella (Cincticostella)*).

Cincticostella nigra (Uéno, 1928) (Kuro-madara-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu)]
nigra Uéno, 1928 (*Ephemerella*); Uéno, 1931c (*Chitonophora(?)*); Allen, 1980 (*Cincticostella (Cincticostella)*); Gose, 1980d (*Ephemerella (Cincticostella)*); Ishiwata, 1987 (*Cincticostella*).

Cincticostella okumai (Gose, 1980) (Okuma-madara-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), Russia (Kuril Is.)]
nigra: Horasawa, 1931 (*Ephemerella*); Allen 1971 (*Ephemerella (Cincticostella)*) Misidentification.
"sp. EC" Gose, 1962 (*Ephemerella*).
okumai Gose, 1980d (*Ephemerella (Cincticostella)*); Ishiwata, 1987 (*Cincticostella*).

Cincticostella tschernovae (Bajkova, 1962) (Cherunoba-madara-kagerou) [Japan (Hokkaido, Honshu), Korea, Russia]
"sp. nax" Imanishi, 1940 (*Ephemerella*).
tschernovae Bajkova, 1962 (*Ephemerella*); Bajkova, 1979 (*Ephemerella (Cincticostella)*); Ishiwata, 1987 (*Cincticostella*).
imanishii Allen, 1971 (*Ephemerella (Cincticostella)*).

Genus *Drunella* Needham, 1909 (Toge-madara-kagerou-zoku)

Drunella basalis (Imanishi, 1937) (Oo-madara-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, Russia (Kuril Is.)]
("?) *tuberculata*: Uéno, 1928 (*Ephemerella*).
basalis Imanishi, 1937a (*Ephemerella*); Edmunds, 1959 (*Ephemerella (Drunella)*); Ishiwata, 1987 (*Drunella*).

Drunella cryptomeria (Imanishi, 1937) (Futakobu-madara-kagerou) [Japan (Honshu)]
cryptomeria Imanishi, 1937a (*Ephemerella*); Ishiwata, 1987 (*Drunella*).
"sp. (trispina group)" Tanaka, 1966 (*Ephemerella*).
bicornis Gose, 1980d (*Ephemerella (Drunella)*) SYN. N.; Ishiwata, 1987 (*Drunella*).

Note: Imanishi (1937) originally described *Ephemerella cryptomeria* based on male imagoes, and the nymphs have never been described. Gose (1980d) originally described *Ephemerella (Drunella) bicornis* by nymphal stages. The nymphs determined by Gose as *E. bicornis* are identical to nymphs collected from the type locality of *E. cryptomeria*, Kufune, Kyoto-fu, from which I reared adult males. Therefore, I regard *E. bicornis* as a synonym of *E. cryptomeria*.

Drunella ishiyamana Matsumura, 1931 (Yoshino-madara-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, Russia (Primor'ye, Sakhalin, Kuril Is.)]
ishiayama Matsumura, 1931a (*Drunella*).

cryptomeria auct., nec Imanishi, 1937a.

"sp. nG" Imanishi, 1940 (*Ephemerella*).

ishiayama [!]: Edmunds, 1959 (*Ephemerella* (*Drunella*)).

yoshinoensis Gose, 1963 (*Ephemerella*) SYN. N.

Note: *Ephemerella yoshinoensis* was first reported as "*Ephemerella nG*" based on the nymph by Imanishi (1940). Gose (1974) regarded the nymph "*Ephemerella nG*" as *E. yoshinoensis* and have erroneously made *E. yoshinoensis* a synonym of *Ephemerella cryptomeria* Imanishi without explanation (Gose, 1980d). Hereafter, *E. yoshinoensis* should be known by the name *E. cryptomeria* auct., nec Imanishi (1937a) (see, note under *Drunella cryptomeria*, above).

Although I have not been successful in locating the holotype of *E. yoshinoensis* (types missing, Gose personal communication), I have collected specimens from Gojo-shi, Nara-ken, the type locality of *E. yoshinoensis*. All of them are identical to *D. ishiyamana* in the respects of size, coloration, and ratio of femora to tibiae as *D. ishiyamana*. Therefore *E. yoshinoensis* is valid and should be synonymized with *D. ishiyamana*.

Drunella kohnoi (Allen, 1971) (Kono-madara-kagerou) [Japan (Honshu, Shikoku, Kyushu)]

kohnoi Allen, 1971 (*Ephemerella* (*Drunella*)).

kohnaoe [!]: Allen, 1980 (*Drunella* (*Tribrochella*)); Uchida, 1986 (*Drunella*).

kohnoi [!]: Gose, 1980d (*Ephemerella* (*Drunella*)).

kohnoae [-]: Gose, 1988 (*Ephemerella* (*Drunella*)).

Drunella sachalinensis (Matsumura, 1931) (Futamata-madara-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), Russia (Kuril Is.)]

sachalinensis Matsumura, 1931a (*Ephemerella*).

trispina: Uéno, 1931c (*Ephemerella*) Misidentification.

bifurcata Allen, 1971 (*Ephemerella* (*Drunella*)) SYN. N.; Ishiwata, 1987 (*Drunella*).

Note: Based on type material examination (*Ephemerella sachalinensis* Matsumura, 1931, in Matsumura Collection, HU; *Ephemerella* (*Drunella*) *bifurcata* Allen, 1971, in INHS), I regard *E. bifurcata* as a synonym of *E. sachalinensis*.

Drunella triacantha (Tshernova, 1949) (Ezo-mitsutoge-madara-kagerou) [Japan (Hokkaido), Korea, Russia]

trispina "subsp. naa" Imanishi, 1940 (*Ephemerella*).

triacantha Tshernova, 1949 (*Ephemerella*); Edmunds, 1959 (*Ephemerella* (*Drunella*)); Ishiwata et al., 2000 (*Drunella*).

trispina ezoensis Gose, 1980d (*Ephemerella* (*Drunella*)); Ishiwata, 1987 (*Drunella*).

Drunella trispina (Uéno, 1928) (Mitsutoge-madara-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), Russia (Kuril Is.)]

trispina Uéno, 1928 (*Ephemerella*); Allen, 1980 (*Drunella* (*Tribrochella*)); Gose, 1980d (*Ephemerella* (*Drunella*)); Ishiwata, 1987 (*Drunella*).

Genus *Ephacerella* Paclt, 1994 (Shirinaga-madara-kagerou-zoku)

Acerella Allen, 1971, nom. preocc.

Ephacerella Paclt, 1994, replacement name for *Acerella* Allen nec Berlese, 1909.

Ephacerella longicaudata (Uéno, 1928) (Shirinaga-madara-kagerou) [Japan (Honshu, Shikoku,

Kyushu), Korea, Russia)]

longicaudata Uéno, 1928 (*Ephemerella*); Edmunds, 1959 (*Ephemerella (Serratella)*); Allen, 1971 (*Ephemerella (Acerella)*); Allen, 1980 (*Acerella*); Kluge, 1997 (*Ephemerella (Ephacerella)*).

Genus *Ephemerella* Walsh, 1862 (Madara-kagerou-zoku)

Ephemerella atagosana Imanishi, 1937 (Hosoba-madara-kagerou) [Japan (Honshu, Shikoku, Kyushu), China, Korea, Russia]

atagosana Imanishi, 1937a (*Ephemerella*); Gose, 1980e (*Ephemerella (Cincticostella)*); Tshernova et al., 1986 (*Ephemerella (Ephemerella)*); Ishiwata, 1987 (*Cincticostella*); Ishiwata, 1993 (*Ephemerella*).

"sp. nb" Imanishi, 1940 (*Ephemerella*).

"sp. nbx" Imanishi, 1940 (*Ephemerella*); Bae et al., 2000.

dentata Bajkova, 1967 (*Ephemerella*); Tshernova et al., 1986 (*Ephemerella (Ephemerella)*).

denticula Allen, 1971 (*Ephemerella (Ephemerella)*) SYN.N.; Ishiwata, 1987 (*Ephemerella*).

keijoensis Allen, 1971 (*Ephemerella (Ephemerella)*).

Note: I have examined the type of *Ephemerella (Ephemerella) denticula* Allen, 1971, in IMNS. Although type series of *Ephemerella atagosana* Imanishi, 1937 are not preserved in the Imanishi Collection, CERK (Ishiwata, personal observation), based on examination of adult specimens reared from nymphs collected from the type locality, Mt. Atago, Kyoto-fu, I regard *E. dentica* as a synonym of *E. atagosana*.

Ephemerella aurivillii (Bengtsson, 1908) (Kita-madara-kagerou) [Japan (Hokkaido, Honshu), Korea, North Europe, Central and eastern Asia, North America]

aurivillii Bengtsson, 1908 (*Chitonophora*); Allen & Edmunds, 1965 (*Ephemerella (Ephemerella)*); Bajkova, 1972 (*Ephemerella*); Gose, 1988 (*Ephemerella (Ephemerella)*).

aroni Eaton, 1908 (*Ephemerella*).

norda McDunnough, 1924 (*Ephemerella*).

aurivilliusi Lestage, 1930 (*Chitonophora*).

concinna Traver, 1934 (*Ephemerella*).

"sp. nN" Imanishi, 1940 (*Ephemerella*).

taeniata Tshernova, 1952 (*Ephemerella*); Gose, 1980e (*Ephemerella (Ephemerella)*).

maxima Allen, 1971 (*Ephemerella (Ephemerella)*) SYN. N.

ezoensis Gose, 1980d (*Ephemerella (Cincticostella)*) SYN. N.; Ishiwata, 1993 (*Cincticostella*).

Note: Based on type material examination of *Ephemerella (Ephemerella) maxima* in IMNS, I regarded it as conspecific with *Ephemerella aurivillii* (Bengtsson, 1908). The type material of *Ephemerella (Cincticostella) ezoensis* are lost (Gose, personal communication), so the nymphs of *E. (C.) ezoensis* determined by Gose were compared with *E. aurivillii* and they were found to be indistinguishable morphologically. Therefore, I regard these two species, *E. (E.) maxima* and *E. (C.) ezoensis*, as synonyms of *E. aurivillii*.

Ephemerella cornuta Gose, 1980 Emendation (Tsuno-madara-kagerou) [Japan (Honshu, Shikoku, Kyushu)]

"sp. EA" Gose, 1962 (*Ephemerella*).

cornutus [!] Gose, 1980e (*Ephemerella (Ephemerella)*); Ishiwata, 1987 (*Ephemerella*).

Note: Ishiwata & Inada (1996) suggested that this species should be a junior homonym for *Ephemerella cornuta* Morgan, 1911. However, when Gose (1980, October 15) described his *E.*

cornutus, Morgan's *E. cornuta* had been already transferred to *Drunella* (Allen 1980, March). Therefore, Gose's *Ephemerella cornuta* is valid.

Ephemerella imanishi Gose, 1980 (Imanishi-madara-kagerou) [Japan (Honshu, Shikoku, Kyushu), Korea]

"sp. nF" Imanishi, 1940 (*Ephemerella*).

imanishi Gose, 1980e (*Ephemerella (Ephemerella)*); Okazaki, 1984 (*Ephemerella*).

Ephemerella ishiwatai Gose, 1985 (Ishiwata-madara-kagerou) [Japan (Honshu, Kyushu)]
ishiwatai Gose, 1985 (*Ephemerella (Ephemerella)*); Ishiwata, 1987 (*Ephemerella*).

**Ephemerella notata* Eaton, 1887 (Ki-madara-kagerou) [Japan (Honshu, Kyushu), Europe, Russia]

notata Eaton, 1887 (*Ephemerella*); Tshernova et al., 1986 (*Ephemerella (Ephemerella)*).

ishiwatai: Kobayashi, 1986 (*Ephemerella*) Misidentification.

"sp. 2" Kobayashi & Nishino, 1992 (*Ephemerella*).

Ephemerella setigera Bajkova, 1965 (Kushige-madara-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, Russia]

"sp. EB" Gose, 1962 (*Ephemerella*).

setigera Bajkova, 1965 (*Ephemerella*); Gose, 1980e (*Ephemerella (Serratella)*); Yamasaki, 1987 (*Serratella*); Ishiwata, 1987 (*Ephemerella*); Kluge, 1997 (*Ephemerella (Torleya)*).

Genus *Torleya* Lestage, 1917 (Erabuta-madara-kagerou-zoku)

**Torleya glareosa* Kang & Yang, 1995 (Yaeyama-madara-kagerou) [Japan (Okinawa: Iriomote-jima Is., Ishigaki-jima Is.), Taiwan]

"sp." Tanida, 1974 (*Ephemerella*).

glareosa Kang & Yang, 1995 (*Torleya*).

Torleya japonica (Gose, 1980) (Erabuta-madara-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu)]

"sp. ED" Gose, 1962 (*Ephemerella*).

japonica Gose, 1980d (*Ephemerella (Torleya)*); Ishiwata, 1987 (*Torleya*).

Genus *Uracanthella* Belov, 1979 (Aka-madara-kagerou-zoku)

Uracanthella chinoi (Gose, 1980) (Chino-madara-kagerou) [Japan (Honshu)]

"sp. EE" Chino, 1975 (*Ephemerella*).

Chinoi [!] Gose, 1980e (*Ephemerella (Serratella)*).

chinoi Gose, 1985 (*Ephemerella (Serratella)*); Ishiwata, 1987 (*Serratella*); Takemon, 1990 (*Uracanthella*).

Uracanthella punctisetae (Matsumura, 1931) COMB. N. (Aka-madara-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, Russia]

punctisetae Matsumura, 1931a (*Drunella*).

rufa Imanishi, 1937a (*Ephemerella*) SYN. N.; Gose, 1980e (*Ephemerella (Serratella)*); Uchida, 1986 (*Serratella*); Tshernova et al., 1986 (*Ephemerella (Uracanthella)*); Hatta & Ishiwata, 1990 (*Uracanthella*).

Note: I examined the holotype of *Drunella punctisetae* Matsumura, 1931 in the Matsumura

Collection, HU. Although I have not been successful in locating the holotype of *Ephemerella rufa* Imanishi, 1937, I found the male specimen identified with *E. rufa* by Imanishi in the Imanishi Collection, CERK, the specimens I have collected from Ichihara, near Kamogawa River, Kyoto-fu, the type locality of *E. rufa* are identical with *U. punctisetae* in regard to size, coloration, and detail morphology of the genitalia. Therefore, I regard *E. rufa* as a synonym of *D. punctisetae*.

Ephemerella gose Okazaki, 1984 *nomen dubium*.
gose Okazaki, 1984 (*Ephemerella*) *nomen dubium*.

Genus *incertae sedis*

Ephemerella elongatula McLachlan, 1875 (Hoso-madara-kagerou) [Japan (Honshu), China (?)]
elongatula McLachlan, 1875 (*Leptophlebia*); Eaton, 1885 (*Ephemerella*).

Ameletidae (Hime-futao-kagerou-ka)

Genus *Ameletus* Eaton, 1887 (Hime-futao-kagerou-zoku)
Chimura Navás, 1915.

Ameletus aethereus (Navás, 1915) Emendation (Nabasu-hime-futao-kagerou) [Japan (Honshu)]
aetherea Navás, 1915 (Chimura).
aetherea [!] Edmunds, 1960 (*Ameletus*).

Ameletus costalis (Matsumura, 1931) (Maeguro-hime-futao-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea, Russia]
costalis Matsumura, 1931a (*Ameletus*).

Ameletus croceus Imanishi, 1932 (Yokosuji-hime-futao-kagerou) [Japan (Honshu)]
crocus Imanishi, 1932 (*Ameletus*).

Ameletus kyotensis Imanishi, 1932 (Kyoto-hime-futao-kagerou) [Japan (Honshu)]
kyotensis Imanishi, 1932 (*Ameletus*).
kyotoensis [!]: Gose, 1979b (*Ameletus*).

Ameletus montanus Imanishi, 1930 (Hime-futao-kagerou) [Japan (Hokkaido, Honshu, Kyushu), Korea]
montanus Imanishi, 1930 (*Ameletus*).
metanus [!]: Gose, 1979b (*Ameletus*).

Ameletus subalpinus Imanishi, 1932 (Kurobe-hime-futao-kagerou) [Japan (Honshu)]
subalpinus Imanishi, 1932 (*Ameletus*).

Ameletus towadensis Matsumura, 1931 *nomen dubium*
towadensis Matsumura, 1931a (*Ameletus*) *nomen dubium*.

Baetidae (Ko-kagerou-ka)

Genus *Acentrella* Bengtsson, 1912 (Mijikao-futaba-ko-kagerou-zoku)

***Acentrella gnom* (Kluge, 1983)** (Mitsuo-mijikao-futaba-ko-kagerou) [Japan (Honshu, Kyushu), Russia, Korea, Taiwan]
gnom Kluge, 1983 (*Pseudocloeon*); Tshernova et al., 1986 (*Baetis (Acentrella)*); Waltz & McCafferty, 1987 (*Acentrella*); Kobayashi, 1992 (*Acentrella*).
ultimum Müller-Liebenau, 1985 (*Pseudocloeon*).

***Acentrella lata* (Müller-Liebenau, 1985)** (Minami-mijikao-futaba-ko-kagorou) [Japan (Okinawa: Okinawa-honto), Taiwan]
latum Müller-Liebenau, 1985 (*Pseudocloeon*); Nakasone et al., 1998 (*Pseudocloeon*).
lata: Waltz & McCafferty, 1987 (*Acentrella*).

***Acentrella sibirica* (Kazlauskas)** (Mijikao-futaba-ko-kagerou) [Japan (Honshu), Korea, Russia]
sibiricum Kazlauskas, 1963 (*Pseudocloeon*).
nosegawaensis: Gose, 1980c **Misidentification**, nec Gose, 1965 (*Pseudocloeon*).
sibiricus: Tshernova et al., 1986 (*Baetis (Acentrella)*).
sibirica: Waltz & McCafferty, 1987 (*Acentrella*); Ishiwata, 2000 (*Acentrella*).
sp. "B" Kobayashi, 1992 (*Acentrella*).

Note: The nymph described as *Pseudocloeon nosegawaensis* by Gose (1980c) is referred to a species of *Acentrella* because adult specimens reared from the nymphs agreed well with characteristics of *Acentrella* having sharply projected anteronotal protuberance (ANp) on mesothorax (Kluge, 1994; Edmunds & Waltz, 1996).

***Acentrella suzukiella* Matsumura, 1931** (Suzuki-mijikao-futaba-ko-kagerou) [Japan (Honshu)]
suzukiella Matsumura, 1931a (*Acentrella*).

Genus *Baetiella* Uéno, 1931 (Futaba-ko-kagerou-zoku)

Pseudocloeon Matsumura, 1931a. nom. preocc.

Neobaetiella Müller-Liebenau, 1985.

Matsumuracloeon Hubbard, 1989 **SYN.** N. replacement name for *Pseudocloeon* Matsumura, 1931a nec Klapálek, 1905.

Note: Based on examination of the type material, I recognized *Pseudocloeon aino* Matsumura, the type species of *Pseudocloeon* Matsumura, 1931, is conspecific with *Baetiella japonica*, the type species of *Baetiella* Uéno 1931. Although the former generic name is senior to the latter, the name *Baetiella* is valid because the name *Pseudocloeon* Matsumura, 1931, is preoccupied by *Pseudocloeon* Klapálek, 1905 (Hubbard, 1988). *Matsumuracloeon* Hubbard, 1989, a replacement name, for *Pseudocloeon* Matsumura (Hubbard, 1989), is therefore a junior synonym of *Baetiella*.

***Baetiella bispinosa* (Gose, 1980)** (Togetoge-futaba-ko-kagerou) [Japan (Okinawa: Iriomote-jima Is., Ishigaki-jima Is.), Taiwan]
bispinosus Gose, 1980c (*Pseudocloeon*).
macani Müller-Liebenau, 1985 (*Neobaetiella*).
bispinosa: Waltz & McCafferty, 1987 (*Baetiella*).

***Baetiella japonica* (Imanishi, 1930)** (Futaba-ko-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu)]
"n. sp.?" Uéno, 1928 (*?Acentrella*).

japonica, Imanishi, 1930 (*Acentrella*).

aino Matsumura, 1931a (*Pseudocloeon*) SYN. N.; Waltz & McCafferty, 1987 (*Baetiella*);

Hubbard, 1990 (*Matsumuracloeon*).

japonica: Uéno, 1931a (*Baetiella*).

nosegawaensis Gose, 1965 (*Baetiella*) SYN. N.

japonica [!]: Gose, 1980c (*Pseudocloeon*).

japonicum: Müller-Liebenau, 1985 (*Pseudocloeon*).

Note: *Acentrella japonica* Imanishi, 1930 is a senior synonym of *Pseudocloeon aino* Matsumura (see the note on the genus *Baetiella* above) and *Baetiella nosegawaensis* Gose. Kluge (1983) synonymized *Baetiella nosegawaensis* Gose, 1965 with *Baetiella tuberculata* (Kazlauskas, 1963). I have examined Korean specimens of *B. tuberculata*, and found that the species is much different from *B. nosegawaensis*, in having the median tubercles on posterior margin of each tergum. According to Gose (1980c), *B. nosegawaensis* is distinguished from *B. japonica* by the presence of a projection at basal segment of male genital forceps. However, my examination of reliable materials of *B. japonica* detected the presence of the projection. Although the type material of both of *B. japonica* and *B. nosegawaensis* were missing (personal observation and Gose, personal communication, respectively), I recognize that *Baetiella nosegawaensis* should be considered as a synonym of *Baetiella japonica*.

Genus *Baetis* Leach, 1815 (Ko-kagerou-zoku).

Note: The genus *Baetis* tends to be split into many subgroups by many authors. Waltz *et al.* (1994) and McCafferty & Waltz (1995) created a new genus and gave subgeneric rank based on morphological differences in nymphs. Kobayashi (1987) revised the Japanese species of *Baetis* and presented a nymphal key to seven nominal species together with 13 informal species (20 species in total). He also assigned them to some species groups. Although twenty-two species are listed here, generic or subgeneric assignment is not given. Further taxonomical studies, including generic or subgeneric assignment are needed.

Baetis acuminatus Gose, 1980 (Togari-ko-kagerou) [Japan]

acuminatus Gose, 1980b (*Baetis*).

Baetis atagonis Imanishi, 1937 (Atago-ko-kagerou) [Japan (Honshu)]

atagonis Imanishi, 1937b (*Baetis*).

Baetis bicaudatus Dodds, 1923 (Futao-ko-kagerou) [Japan (Hokkaido, Honshu), Mongolia, Russia, North America]

bicaudatus Dodds, 1923 (*Baetis*); Kluge, 1997 (*Baetis (Baetis)*); Ishiwata *et al.*, 2000 (*Baetis*).

"sp. L" Kobayashi, 1987 (*Baetis*).

Baetis ceculus Imanishi, 1937 (Kerukusu-ko-kagerou) [Japan (Honshu)]

ceculus Imanishi, 1937b (*Baetis*).

ceculus [!]: Gose, 1980b (*Baetis*).

Baetis chocoratus Gose, 1980 (Tobiiro-ko-kagerou) [Japan (Honshu, Shikoku, Kyushu)]

chocoratus Gose, 1980a (*Baetis*).

Baetis florens Imanishi, 1937 (Furorensu-ko-kagerou) [Japan (Honshu)]

florens Imanishi, 1937b (*Baetis*).

Baetis frexifemora Gose, 1980 (Udemagari-ko-kagerou) [Japan (Honshu)]
frexifemora Gose, 1980b (*Baetis*).

Baetis fuscatus (Linnaeus, 1761) (Usuguro-ko-kagerou) [Japan, Europe, Russia, Korea]
fuscatus Linnaeus, 1761 (*Baetis*); Novikova & Kluge, 1987 (*Baetis (Baetis)*).
bioculatus: Uéno, 1928 (*Baetis*).
"sp. nla" Imanishi, 1940 (*Baetis*).

Baetis hyugensis Gose, 1980 (Hyuga-ko-kagerou) [Japan (Honshu)]
hyugensis Gose, 1980a (*Baetis*).

Baetis iriomotensis Gose, 1980 (Iriomote-ko-kagerou) [Japan (Okinawa: Iriomote-jima Is.)]
"sp. (No.1)" Uéno, 1969 (*Baetis*).
iriomotensis Gose, 1980b (*Baetis*).

Baetis muticus (Linnaeus, 1758) (Kiso-ko-kagerou) [Japan (Honshu), Korea, Russia, England, Europe]
muticus Linnaeus, 1758 (*Ephemera*); Novikova & Kluge, 1987 (*Baetis (Nigrobaetis)*); Waltz et al., 1994 (*Alainites*).
pumilus Burmeister, 1839; Horasawa, 1931 (*Baetis*).
furcatus Navás 1934 (*Baetis*).

Baetis pseudofrequentus Müller-Liebenau, 1985 (Minami-ko-kagerou) [Taiwan, Japan (Okinawa: Iriomotejima Is.)]
pseudofrequentus Müller-Liebenau, 1985 (*Baetis*); Kobayashi, 1989 (*Baetis*); Kang et al., 1994 (*Baetis (Tenuibaetis)*).

Baetis sacishimensis Uéno, 1969 (Sakishima-ko-kagerou) [Japan (Okinawa: Ishigaki-jima Is., Iriomote-jima Is.)]
sacishimensis Uéno, 1969 (*Baetis*); Waltz et al., 1994 (*Alainites*).

Baetis sahoensis Gose, 1980 (Saho-ko-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu)].
sahoensis Gose, 1980a (*Baetis*).

Baetis shinanonis Uéno, 1931 (Shinano-ko-kagerou) [Japan (Honshu)]
shinanonis Uéno, 1931a (*Baetis*).

Baetis takamiensis Gose, 1980 (Takami-ko-kagerou) [Japan (Honshu)]
takamiensis Gose, 1980a (*Baetis*).
Baetis thermicus Uéno, 1931 (Shirohara-ko-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu, Okinawa), Korea, Russia (Kuril Is.), Taiwan]
thermicus Uéno, 1931a (*Baetis*).

Baetis totukawensis Gose, 1980 (Totukawa-ko-kagerou) [Japan (Honshu)]
totukawensis Gose, 1980a (*Baetis*).

Baetis tsushimaensis Gose, 1980 (Tsushima-ko-kagerou) [Japan (Kyushu)]
tsushimaensis Gose, 1980a (*Baetis*).

Baetis uenoi Gose, 1980 (Ueno-ko-kagerou) [Japan (Okinawa: Iriomote-jima Is.)]
"sp. No.2, new species" Uéno, 1969 (*Baetis*).
uenoi Gose, 1980b (*Baetis*).

Baetis yamatoensis Gose, 1980 (Yamato-ko-kagerou) [Japan (Honshu)]
yamatoensis Gose, 1980a (*Baetis*).

Baetis yoshinensis Gose, 1980 (Yoshino-ko-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu, Okinawa: Okinawa-honto), Taiwan]
yoshinensis Gose, 1980a (*Baetis*).

Genus *Centropilum* Eaton, 1869 (Usuba-ko-kagerou-zoku)

Centropilum rotundum Takahashi, 1929 (Usuba-ko-kagerou) [Japan (Honshu, Shikoku)]
rotundum Takahashi, 1929 (*Centropilum*).

Genus *Cloeon* Leach, 1815 (Futaba-kagerou-zoku)

Cloeon dipterum (Linnaeus, 1761) (Futaba-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu)]
diptera Linnaeus, 1761 (*Ephemerata*).
dipterum: Leach, 1815 (*Cloeon*); McLachlan, 1875 (*Cloeon*); Matsumura, 1904 (*Cloeon*); Kluge, 1997 (*Cloeon* (*Cloeon*)).

Cloeon kyotonis Matsumura, 1931 (Kyoto-futaba-kagerou) [Japan (Honshu)]
kyotonis Matsumura, 1931a (*Cloeon*).

Cloeon maikonis (Takahashi, 1924) (Amefuri-futaba-kagerou) [Japan (Honshu)]
maikonis Takahashi, 1924 (*Cloeon*).

Cloeon marginale Hagen, 1858 (Akamon-futaba-kagerou) [Japan (Okinawa: Okinawa-honto), Taiwan, China (South), India, Indonesia, Philipines, Sri Lanka]
marginalis Hagen, 1858 (*Cloe.*).
marginale: Eaton, 1871 (*Cloeon*); Uéno, 1969 (*Cloeon*).

Cloeon okamotoi Takahashi, 1924 (Aoba-futaba-kagerou) [Japan (Honshu, Shikoku)]
okamotoi Takahashi, 1924 (*Cloeon*).

Cloeon ryogokuense Gose, 1980 Emendation (Tamari-futaba-kagerou) [Japan (Honshu, Kyushu)]
ryogokuensis [!] Gose, 1980a (*Cloeon*).

Cloeon tamagawanum (Matsumura, 1931) COMB. N. (Tamagawa-futaba-kagerou) [Japan (Honshu)]
tamagawanum Matsumura, 1931a (*Procloeon*); Hubbard, 1988 (*Promatsumura*).

Genus *Procloeon* Bengtsson, 1915 (Himeusuba-ko-kagerou-zoku)

Procloeon bimaculatum (Eaton, 1885) (Himeusuba-ko-kagerou) [Japan (Okinawa: Ishigaki-jima Is.), Taiwan, Java, Sumatra, Thailand, Sri Lanka]

bimaculatum Eaton, 1885 (*Cloeon*); Kimmins, 1947 (*Procloeon*); Uéno, 1969 (*Cloeon*).

Genus *Promatsumura* Hubbard, 1988 (Matsumura-ko-kagerou-zoku)

Procloeon Matsumura, 1931a, nom. preocc. Not available, no type designated.

Promatsumura Hubbard, 1988, replacement name for *Procloeon* Matsumura, 1931a nec *Procloeon* Bengtsson, 1915.

Promatsumura nipponicum (Matsumura, 1931) (Matsumura-ko-kagerou) [Japan (Honshu)]

nipponicum Matsumura, 1931a (*Procloeon*); Hubbard, 1988 (*Promatsumura*).

Dipteromimidae (Gaganbo-kagerou-ka)

Genus *Dipteromimus* McLachlan, 1875 (Gaganbo-kagerou-zoku)

Dipteromimodes Matsumura, 1931b

Dipteromimus tipuliformis McLachlan, 1875 (Gaganbo-kagerou) [Japan (Honshu, Shikoku, Kyushu)]

tipuliformis McLachlan, 1875 (*Dipteromimus*).

suzukii Matsumura, 1931a (*Dipteromimodes*).

Siphlonuridae (Futao-kagerou-ka)

Genus *Siphlonurus* Eaton, 1868 (Futao-kagerou-zoku)

Siphlonurus binotatus (Eaton, 1892) (Oo-futao-kagerou) [Japan (Hokkaido, Honshu, Shikoku), China]

binotatus Eaton, 1892 (*Siphlurus*).

alternatus: Uéno, 1928 (*Siphlurus* (?)).

maculosus Takahashi, 1930 (*Siphlonurus*).

grandiosa Matsumura, 1931a (*Siphlonisca*).

Siphlonurus sanukensis Takahashi, 1929 (Nami-futao-kagerou) [Japan (Honshu, Shikoku), Korea]

sanukensis Takahashi, 1929 (*Siphlurus*); Imanishi, 1933 (*Siphlonurus*).

Siphlonurus yoshinoensis Gose, 1979 (Yoshino-futao-kagerou) [Japan (Honshu)]

yoshinoensis Gose, 1979c (*Siphlonurus*).

Siphlonurus zhelochovtsevi Tshernova, 1952 (Ezo-futao-kagerou) [Japan (Hokkaido), Russia]

zhelochovtsevi Tshernova, 1952 (*Siphlonurus*); Itou, 1989 (*Siphlonurus*).

Isonychiidae (Chira-kagerou-ka)

Genus *Isonychia* Eaton, 1871 (Chira-kagerou-zoku)

Subgenus *Isonychia* Eaton (Chira-kagerou-azoku)

Isonychia japonica (Ulmer, 1919) (Chira-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), China, Korea, Russia]

japonicus Ulmer, 1919 (*Chirotenetes*); Uéno, 1928 (*Chirotenetes* (?)).

violacea Matsumura, 1931a (*Sparrea*).
japonica: Uéno & Okamoto, 1932 (*Isonychia*).

Subgenus *Prionoides* Kondoratief & Voshell (Shima-chira-kagerou-azoku)

Isonychia shima (Matsumura, 1931) SP. REV. & COMB. N. (Shima-chira-kagerou) [Japan (Hokkaido, Honshu)]
shima Matsumura, 1931a (*Rhoenanthus*).
jozana Matsumura, 1931a (*Siphlonisca*) SYN. N.
sukashii Matsumura, 1931a (*Siphlonisca*) SYN. N.

Note: *Rhoenanthus shima* Matsumura, 1931 was declared to be a *nomen dubium* by Bae & MacCafferty (1991). However, it is a valid species since the female syntype has been preserved in the Matsumura Collection, HU. Based on page priority, the last two names in the list above are placed as junior synonyms of *Isonychia shima* (Matsumura, 1931).

Isonychia mitsukonis Morishita, 1961 *nomen dubium*
mitsukonis Morishita, 1961 (*Isonychia*) *nomen dubium*.

Subgenus *incertae sedis*

Isonychia valida Navás, 1919 (Navasu-chira-kagerou) [Japan (Honshu)]
validus Navás, 1919 (*Chirotonetes*).
valida; Imanishi, 1933 (*Isonychia*).

Oligoneuriidae (Hitoriga-kagerou-ka)

Genus *Oligoneuriella* Ulmer, 1924 (Hitoriga-kagerou-zoku)

Oligoneuriella rhenana Imhoff, 1852 (Hitoriga-kagerou) [Europe, Japan (Honshu)]
rhenana Imhoff, 1852 (*Oligoneuriella*); Uéno 1950 (*Oligoneuriella*).

Heptageniidae (Hirata-kagerou-ka)

Genus *Bleptus* Eaton, 1885 (Obi-kagerou-zoku)

Bleptus fasciatus Eaton, 1885 (Obi-kagerou) [Japan (Honshu, Shikoku, Kyushu), Korea]
fasciatus Eaton, 1885 (*Bleptus*).

Genus *Cinygma* Eaton, 1885 (Kita-tanigawa-kagerou-zoku)

Cinygma lyriformis (McDunnough, 1924) (Oo-ezo-kagerou) [Japan (Hokkido)], Korea, Russia, North America
lyriformis McDunnough, 1924 (*Ecdyonurus*); Kuhara et al., 1993 (*Cinygma*).
peterseni Lestage, 1930 (*Ecdyonurus*).

Genus *Cinygmoidea* Matsumura, 1931 (Ezo-tanigawa-kagerou-zoku)

Cinygmoidea hekachii Matsumura, 1931 (Hekachi-ezo-tanigawa-kagerou) [Japan (Hokkaido)]
hekachii Matsumura, 1931a (*Cinygmoidea*); Matsumura, 1933 (*Ecdyonurus* (*Cinygmoidea*)).

Genus *Cinygmula* McDunnoug, 1933 (Miyama-tanigawa-kagerou-zoku)

Note: The first comprehensive study of the genus *Cinygmula* in Russian Far East and Central Asia was given by Tshernova & Belov (1982), who demonstrated the similarity of the two genera, *Cinygmula* and *Rhithrogena*. Kluge (1988, 1997) regarded *Cinygmula* as a subgenus of the genus *Rhithrogena*, because there are few morphological differences in adults. In Japan and North America, however, no comprehensive works on *Cinygmula* has been carried out.

Cinygmula adusta (Imanishi, 1935) (Chairo-miyama-tanigawa-kagerou) [Japan (Honshu, Shikoku)]

adusta Imanishi, 1935 (*Cinygma*); Tshernova & Belov, 1982 (*Cinygmula*).

Cinygmula cava (Ulmer, 1927) (Ezo-miyama-tanigawa-kagerou) [Japan (Hokkaido), Mongolia, Russia]

cavum Ulmer, 1927 (*Cinygma*).

cavum ?: Uéno & Okamoto, 1932 (*Cinygma*).

cava: Tshernova & Belov, 1982 (*Cinygmula*); Kluge, 1988 (*Rhithrogena (Cinygmula)*); Ishiwata, 1999 (*Cinygmula*).

Cinygmula dorsalis (Imanishi, 1935) (Sesuji-miyama-tanigawa-kagerou) [Japan (Honshu)]

dorsalis Imanishi, 1935 (*Cinygma*); Tshernova & Belov, 1982 (*Cinygmula*).

Cinygmula hirasana (Imanishi, 1935) (Miyama-tanigawa-kagerou) [Japan (Honshu), Korea sp. nov. ? Uéno, 1928 (*Ecdyurus*).

hirasana Imanishi, 1935 (*Cinygma*); Tshernova & Belov, 1982 (*Cinygmula*); Kluge, 1997 (*Rhithrogena (Cinygmula)*).

kurobensis Imanishi, 1935 (*Cinygma*); Tshernova & Belov, 1982 (*Cinygmula*).

Cinygmula putoranica Kluge, 1980 (Kuro-miyama-tanigawa-kagerou) [Japan (Hokkaido), Russia]

putoranica Kluge, 1980 (*Cinygmula*); Kluge, 1988 (*Rhithrogena (Cinygmula)*); Kuhara et al., 1993 (*Cinygmula*).

Cinygmula sapporensis (Matsumura, 1904) SP. REV. & COMB. N. (Kuro-miyama-tanigawa-kagerou) [Japan (Hokkaido, Honshu), Korea, Russia]

sapporensis Matsumura, 1904 (*Siphlurus*); Matsumura, 1931b (*Cinygma*); Matsumura, 1933 (*Ecdyonurus (Cinygma)*).

sapporense Matsumura, 1931a (*Cinygma*).

sapporensis: Imanishi, 1932 (*Ameletus*) Misidentification.

grandifolia Tshernova, 1952 (*Cinygmula*) SYN. N.; Kluge, 1988 (*Rhithrogena (Cinygmula)*).

Note: After examination of type specimens of *Siphlurus sapporensis* Matsumura, 1904, I found that this is a valid species and a senior synonym of *Cinygmula grandifolia* Tshernova, 1952. This species was previously synonymized with *Ameletus costalis* (Matsumura) by Imanishi (1932).

Cinygmula vernalis (Imanishi, 1935) (Haruno-miyama-tanigawa-kagerou) [Japan (Honshu)]

vernalis Imanishi, 1935 (*Cinygma*); Tshernova & Belov, 1982 (*Cinygmula*).

Genus *Ecdyonurus* Eaton, 1868 (Tanigawa-kagerou-zoku)

Note: The concept of the genus *Ecdyonurus* by Kluge (1988) is adopted here until more

reasonable classification is proposed in future. The egg morphology of Japanese species of *Ecdyonurus* indicates several groups in the genus (Okazaki, 1984). In addition, the Japanese species of the genus are distinguishable from European species by the shape of male genitalia and hind margin of nymphal pronotum. In European species, the penis-lobes spread outwards distally to form a broad T-shape apex and the lateral projections of nymphal pronotum are elongated backwards distinctly on either side of mesonotum. Lacking in such features, the Japanese species currently assigned to *Ecdyonurus* should be examined in detail.

Ecdyonurus bajkovaе Kluge, 1986 (Onihime-tanigawa-kagereou) [Japan (Honshu, Shikoku, Kyushu), Korea, Russia]

*bajkovaе Kluge, 1986 (Ecdyonurus); Kluge, 1997 (Ecdyonurus (*Afghanurus*)), replacement name for *Paracinygma zhiltzovae* Bajkova, 1975; Ishiwata, 2000 (Ecdyonurus).*

"sp. A" Yamasaki, 1987 (Ecdyonurus).

subspinosa Braasch & Soldan, 1988 (Nixe).

Ecdyonurus flavus Takahashi, 1929 (Kiiro-tanigawa-kagerou) [Japan (Honshu)]

flavus Takahashi, 1929 (Ecdyurus); Imanishi, 1936 (Ecdyonurus).

**Ecdyonurus fractus* (Kang & Yang, 1994) COMB. N. (Minami-madara-tanigawa-kagerou) [Japan (Okinawa: Iriomote-jima Is.), Taiwan]

fracta Kang & Yang, 1994a (Electrogena).

"sp." Takara & Azuma, 1972 (Bleptus) Misidentification.

"sp." Takara & Azuma, 1972 (Siphlonurus) Misidentification.

Ecdyonurus hyalinus (Ulmer, 1912) COMB. N. (Minami-tanigawa-kagerou) [Japan (Okinawa: Iriomote-jima Is., Ishigaki-jima Is.), Taiwan]

hyalinus Ulmer, 1912 (Ecdyurus); Uéno, 1931c (Ecdyonurus); Kang & Yang, 1994a (*Afronurus*).

"sp. No.2" Uéno, 1928 (Ecdyurus).

Ecdyonurus kibunensis Imanishi, 1936 (Kibune-tanigawa-kagerou) [Japan (Honshu, Shikoku, Kyushu), Korea, Russia]

kibunensis Imanishi, 1936 (Ecdyonurus).

Ecdyonurus naraensis Gose, 1968 SP. REV. (Ashiguro-tanigawa-kagerou) [Japan (Honshu)]

naraensis Gose, 1968 (Ecdyonurus).

Note: After examination of specimens collected from the type locality (Gojo-shi, Nara-ken) (type species lost, Gose personal communication), I have found that this is a valid name for a species previously synonymized with *Rhiithrogena satsuki* Imanishi by Gose (1979e). See the note on *Rhiithrogena tetrapunctiger* below.

Ecdyonurus scalaris Kluge, 1983 (Hime-tanigawa-kagerou) [Japan (Honshu, Kyushu), Korea, Russia]

scalaris Kluge, 1983 (Ecdyonurus); Ishiwata, 2000 (Ecdyonurus).

Ecdyonurus tigris Imanishi, 1936 (Tora-tanigawa-kagerou) [Japan (Honshu, Shikoku, Kyushu)]

tigris Imanishi, 1936 (Ecdyonurus).

Ecdyonurus tobilironis Takahashi, 1929 (Kuro-tanigawa-kagerou) [Japan (Honshu, Shikoku,

Kyushu)]

tobiironis Takahashi, 1929 (*Ecdyonurus*).

suzukianum Matsumura, 1931a (*Cinygma*) SYN. N.

Note: After examination of type specimen of *Cinygma suzukianum* Matsumura, 1931, I found that this is a junior synonym of *Ecdyonurus tobiironis* Takahashi, 1929. Imanishi (1936) examined this specimen in the Matsumura Collection and labeled "*Ecdyonurus tobiironis*". There are, however, no synonymies in his listings.

Ecdyonurus viridis (Matsumura, 1931) COMB. N. (Midori-tanigawa-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Okinawa: Okinawa-honto)]

viridis Matsumura, 1931a (*Heptagenia*).

yoshidae: Uéno, 1938 (*Ecdyonurus*) Misidentification.

hyalinus: Uéno, 1969 (*Ecdyonurus*) Misidentification.

Ecdyonurus yoshidae Takahashi, 1924 (Shiro-tanigawa-kagerou) [Japan (Honshu, Shikoku, Kyushu)]

yoshidae Takahashi, 1924 (*Ecdyonurus*).

japonicus Uéno, 1928 (*Ecdyurus*).

levis: Ishiwata, 1997 (*Ecdyonurus*) Misidentification.

Genus *Epeorus* Eaton, 1881 (Hirata-kagerou-zoku)

Epeorus aesculus Imanishi, 1934 (Kiuro-hirata-kagerou) [Japan (Hokkaido, Honshu, Kyushu), Korea, Russia]

aesculus Imanishi, 1934 (*Epeorus*); Sinitchenkova, 1978 (*Iron*); Kluge, 1988 (*Epeorus* (*Iron*)).

Epeorus cumulus Imanishi, 1939 STAT. N. (Kumurusu-hirata-kagerou) [Japan (Honshu)]

curvatulus f. *cumulus* Imanishi, 1939 (*Epeorus*).

curvatulus *cumulus*: Takemon, 1990 (*Epeorus*).

Epeorus curvatulus Matsumura, 1931 (Ueno-hirata-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu)]

"sp. ?" Uéno, 1928 (*Iron*).

curvatulus Matsumura, 1931a (*Epeorus*).

uenoi auct., nec Matsumura, 1933 (*Epeorus*).

Note: This species was firstly reported as nymphal stage under the name "*Iron* sp. ?" by Uéno (1928). Imanishi (1934) erroneously assigned this species to the nymph of *Epeorus uenoi* (Matsumura) and described the adults.

**Epeorus erratus* Braasch, 1981 (Minami-hirata-kagerou) [Japan (Okinawa: Ishigaki-jima Is., Iriomote-jima Is.), Taiwan]

psi Eaton, 1885 (*Epeorus*); Ulmer, 1912 (*Epeorus*); Uéno, 1931 (*Epeorus*); Imanishi, 1934 (*Epeorus*) Misidentification.

erratus Braasch 1981 (*Epeorus*).

"sp.", Uéno, 1928 (*Epeorus*).

Epeorus hiemalis Imanishi, 1934 (Onaga-hirata-kagerou) [Japan (Honshu)]

hiemalis Imanishi, 1934 (*Epeorus*).

Epeorus ikanonis Takahashi, 1924 (Nami-hirata-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu)]

ikanonis Takahashi, 1924 (*Epeorus*).

hanazonensis Matsumura, 1931a (*Epeorus*).

Epeorus latifolium Uéno, 1928 (Erumon-hirata-kagerou) [Japan (Hokkaido, Honshu, Shikoku), Korea]

latifolium Uéno, 1928 (*Epeorus*); Kluge, 1988 (*Epeorus* (*Belovius*)).

Epeorus l-nigrus Matsumura, 1931 Emendation (Matsumura-hirata-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu)]

latifolium: Horasawa, 1929 (*Epeorus*) Misidentification.

L-nigrum [!] Matsumura, 1931a (*Epeorus*).

Note: Based on type material examination of *Epeorus l-nigrus* Matsumura, 1931 in the Matsumura Collection, HU, I recognize it is a valid name for a species confused with *Epeorus latifolium* Uéno, 1928 by previous authors.

Epeorus napaeus Imanishi, 1934 (Tani-hirata-kagerou) [Japan (Honshu, Shikoku, Kyushu)]
napaeus Imanishi, 1934 (*Epeorus*).

Epeorus nipponicus (Uéno, 1931) COMB. N. (Yumimon-hirata-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea]

nipponicus Uéno, 1931a (*Iron*).

curvatulus f. *curvatulus* Imanishi, 1939 (*Epeorus*) Misidentification.

curvatulus *curvatulus*: Takemon, 1990 (*Epeorus*) Misidentification.

curvatulus auct., nec Matsumura 1931a (*Epeorus*).

Note: After original description of *Iron nipponicus* by Uéno (1931b), this species has been listed by some authors (Uéno, 1932; Imanishi, 1934), but after that it has been overlooked by previous authors. Although I have not been successful in locating the holotype of *I. nipponicus*, I have collected specimens from Nakabusa River, Shinanozaka, Nagano-ken, the type locality of *I. nipponicus*. All of them are identical in regard to size, coloration, and detail of the male genitalia as *I. nipponicus*.

Epeorus uenoi (Matsumura, 1933) [Kita-hirata-kagerou] [Japan (Hokkaido)]

uenoi Matsumura, 1933 nec Imanishi, 1934 (*Iron*).

Note: Matsumura (1933) described a new species under the name of *Iron uenoi* and cited the name of author as "Iron uenoi Imanishi" erroneously. Judging from the examination of the specimens identified with *Epeorus uenoi* (Matsumura, 1933) by Imanishi in the Imanishi Collection, CERK, *uenoi* sensu Imanishi (1934) is not referred to the true *uenoi*. See the note on *Epeorus curvatulus* above.

Genus *Heptagenia* Walsh, 1863 (Kihada-hirata-kagerou-zoku)

Kageronia Matsumura, 1931b.

Heptagenia flava Rostock, 1878 (Sato-kihada-hirata-kagerou) [Japan (Hokkaido, Honshu), China, Europe, Russia]

flava Rostock, 1877 (*Heptagenia*); Kluge, 1988 (*Heptagenia* (*Heptagenia*)).

"sp." Horasawa, 1931 (*Heptagenia*).

"sp. na" Imanishi, 1940 (*Heptagenia*).

"sp. nII" Hashimoto, 1955 (*Heptagenia*).

"sp. HB" Gose, 1962 (*Heptagenia*).

kamonis: Kushiro Insect Lover's Society, 1995 (*Potamanthus*) Misidentification.

***Heptagenia kihada* Matsumura, 1931** (Kihada-hirata-kagerou) [Japan (Honsyu, Shikoku, Kyushu), Korea]

kihada Matsumura, 1931a (*Heptagenia*); Kluge, 1988 (*Heptagenia (Kageronia)*).

suzukiella Matsumura, 1931a (*Kageronia*).

***Heptagenia kyotoensis* Gose, 1963** (Kyoto-kihada-hirata-kagerou) [Japan (Honshu, Shikoku, Kyushu), Korea]

"sp. nb" Imanishi, 1940 (*Heptagenia*).

"sp. nIV" Hashimoto, 1955 (*Heptagenia*).

kyotoensis Gose, 1963 (*Heptagenia*).

***Heptagenia pectoralis* Matsumura, 1931** (Munaguro-kihada-hirata-kagerou) [Japan (Hokkaido, Honshu, Kyushu)]

pectoralis Matsumura, 1931a (*Heptagenia*).

Genus *Rhithrogena* Eaton, 1881 (Hime-hirata-kagerou-zoku)

***Rhithrogena japonica* Uéno, 1928** (Hime-hirata-kagerou) [Japan (Hokkaido, Honshu, Shikoku, Kyushu), Korea]

japonica Uéno, 1928 (*Rhithrogena*).

***Rhithrogena minazuki* Imanishi, 1936** (Minazuki-hime-hirata-kagerou) [Japan (Hokkaido, Honshu, Okinawa: Okinawa-honto), Russia (Sakhalin)]

minazuki Imanishi, 1936 (*Rhithrogena*).

***Rhithrogena parva* (Ulmer, 1912)** (Taiwan-hime-hirata-kagerou) [Japan (Okinawa: Iriomote-jima Is, Ishigaki-jima Is.), Taiwan]

parvus Ulmer, 1912 (*Ecdyurus*).

formosicola Matsumura, 1931a (*Rhithrogena*).

vitrea Walker var. *parva* Ulmer: Uéno, 1931c (*Rhithrogena*).

parva: Uéno, 1969 (*Rhithrogena*).

***Rhithrogena tateyamana* Imanishi, 1936** (Tateyama-hime-hirata-kagerou) [Japan (Honshu)]

tateyamana Imanishi, 1936 (*Rhithrogena*).

magawana Imanishi, 1936 (*Rhithrogena*).

***Rhithrogena tetrapunctigera* Matsumura, 1931** Emendation (Satsuki-hime-hirata-kagerou) [Japan (Honshu), Korea].

4-punctigera [!] Matsumura, 1931a (*Rhithrogena*).

satsuki Imanishi, 1936 (*Rhithrogena*) SYN. N.

"sp. na" Imanishi, 1940 (*Rhithrogena*).

"sp. nl" Hashimoto, 1955 (*Rhithrogena*).

gojoensis Gose, 1968 (*Ameletus*) SYN. N.

Note: Holotype specimen of *Rhithrogena tetrapunctigera* Matsumura, 1931a is a female imago, I collected male imagoes of this species near the type locality, Matsumoto-shi, Nagano-ken. The first figure of male genitalia of *Rhithrogena satsukii* Imanishi 1936 was illustrated by Gose (1979e). But the male genitalia of Gose's illustration were that of *Ecdyonurus naraensis* Gose,

1968. Because Gose (1979e) erroneously synonymized *E. naraensis* with *R. satsuki*, and provided drawing of the male genitalia of *E. naraensis* under the name *R. satsukii*. I could not examine the type specimen of *R. satsuki* because no type specimens have been preserved in the Imanishi Collection, CERK. However, my examination of reliable materials of *R. satsuki* showed the presence of four spots on the ventral thorax, absence of dark longitudinal streak on the femora, and widely separated penis-lobes, which were the diagnostic characters of *R. tetrapunctigera*.

Gose (1968) originally described *gojoensis* under the genus *Ameletus* that had four spots on the ventral thorax. This should be transferred to the genus *Rhithrogena* because of the following characters figured by Gose; the cubital intercalaries of forewing not consist of a series of veinlets (consist of two pairs) (Gose, 1968: fig. 1A), basal segment of fore tarsi less than one-third length of segment 2 (*ibid.* fig. 1D) and penis-lobes widely separated to base (*ibid.* fig. 1F). These are unique characteristics among genus *Rhithrogena*. I have not been successful in locating the holotype of *A. gojoensis*, but I have collected specimens from Gojo-shi, Nara-ken, the type locality of *A. gojoensis*. All of them are identical to *R. tetrapunctigera* in the respects of body size, coloration, and detail morphology of male genitalia. Therefore, I regard that both *R. satsuki* and *A. gojoensis* as synonyms of *R. tetrapunctigera*.

Genus *incertae sedis*

Ecdyonurus bifasciatus Navás, 1933 (Futaobi-tanigawa-kagerou) [Japan (Honshu)]
bifasciatus Navás, 1933 (*Ecdyonurus*?).

Acknowledgements. I am very grateful to Mr. N. Kobayashi, Yokohama, Japan, for helpful information, and Drs. J. Aoki (Yokohama National University, Japan), Y. Takemon (Osaka Prefectural University, Japan) and J. Zloty (University of Calgary, Canada) for their critical comments on the manuscript.

References

- Allen, R.K. 1971. New Asian *Ephemerella* with notes (Ephemerellidae). *Canad. Entomol.* 103: 512-528.
- Allen, R.K. 1980. Geographic distribution and reclassification of the subfamily Ephemerellinae (Ephemeroptera: Ephemerellidae). pp.71-91 In: Flannagan, J. F. & K. E. Marshall (eds). *Advances in Ephemeroptera Biology*. Plenum Publ. Co., New York.
- Allen, R.K., and G.F. Edmunds, Jr. 1965. A revision of the *Ephemerella* VII. The subgenus *Ephemerella* in North America (Ephemeroptera: Ephemerellidae). *Musc. Pub. Ent. Soc. Amer.* 3: 243-282.
- Bae, Y.J. 1997. A historical review of Ephemeroptera systematics in Northeast Asia. pp. 405-417 In: Landolt, P. & M. Sartori (eds). *Ephemeroptera & Plecoptera: Biology-Ecology-Systematics*. MTL. Fribourg, Switzerland.
- Bae, Y.J., N.Ju. Kluge, and D.J. Chun. 1998. New synonymy and new distributional data of mayflies from Korea and Russian Far East (Ephemeroptera). *Zoosyst. Rossica* 7: 89-94.
- Bae, Y.J., J.E. Lee, and I.B. Yoon. 2000. Northeast Asian Ephemeroptera in Imanishi's 1940 report. *Entomol. Sci.* 3(2): 391-397.
- Bae, Y.J. and W.P. McCafferty. 1991. Phylogenetic systematics of the Potamanthidae (Ephemeroptera). *Trans. Amer. Entomol. Soc.* 117(3/4): 1-143.
- Bae, Y.J. and I.B. Yoon. 1997. A revised catalogue of the Ephemeroptera of Korea. *Entomol. Res. Bull.*, Seoul 23: 29-39.
- Bae, Y.J., I.B. Yoon, and D.J. Chun. 1994. A catalogue of the Ephemeroptera of Korea. *Entomol.*

- Res. Bull.*, Seoul 20: 31-50.
- Bajkova, O.Ya. 1962. New species in the genus *Ephemerella* Walsh (Ephemeroptera) from mountain tributaries of the Amur basin. *Izv. Trkhookean. Nauchn. Issled. Inst. Rybu. Khoz. Okeanog.* 48: 202-206. [in Russian]
- Bajkova, O.Ya. 1965. The mayfly fauna of the Far East. *Voprosy geografii Dal'nego Vostoka. Dan'nevostochny filial im. V.L. Komarova SO AN SSSR* 7: 301-330. [in Russian]
- Bajkova, O.Ya. 1967. New species of *Ephemerella* Walsh (Ephemeroptera, Ephemerellidae) from Far East and East Siberia. *Entomol. Obozr.* 46: 327-335. [in Russian]
- Bajkova, O.Ya. 1972. On the mayflies of the Amur basin: I. Imagines (Ephemeroptera: Ephemerellidae). *Izv. Tikhookean. Nauchn. Issled. Inst. Rybn. Khoz. Okeanog.* 77: 178-206. [in Russian]
- Bajkova, O.Ya. 1975. New genus of Ephemeroptera from the Primorye (Ephemeroptera: Heptageniidae). *Izv. Sib. Otd. Akad. Nauk. SSSR Biol. Nauk.* 1: 54-57. [in Russian]
- Bajkova, O.Ya. 1979. A comparative review of the mayfly fauna and of Soviet Far East. pp. 15-20 In: Sowa, R. (ed.). *Proc. 2nd Int. Conf. Ephemeroptera*. Polska Akademia Nauk. Warszawa-Keakow. [in Russian]
- Barnard, K.H. 1932. South African may-flies (Ephemeroptera). *Trans. Royal Soc. South Africa* 20: 201-259.
- Belov, V.V. 1979. A new species of mayflies (Ephemeroptera, Ephemerellidae) for the fauna of the USSR. *Doklady Akademii Nauk Ukrainskoi SSR, ser. B*, 1979: 577-580. [in Russian]
- Bengtsson, S. 1908. Berattelse over en resa i entomologisk syte till mellaesta Sverige Sommaren 1907. *Kungliga Svenska Vetenskapsakademien Arsbok* 6: 237-246. Stockholm.
- Bengtsson, S. 1912. Neue Ephemeriden aus Schweden. *Entomol. Tids.* 33: 107-117.
- Bengtsson, S. 1914. Bemerkungen über die nordischen Arten der Gattung *Clœon* Leach. *Entomol. Tids.* 35: 210-220.
- Bengtsson, S. 1915. Eine Namensänderung. *Entomol. Tids.* 36: 34.
- Braasch, D. 1981. Zum Artstatus des *Epeorus psi* Eaton, 1885 von Taiwan. *Disch. Ent. Z., N. F.* 28, Heft 1-III: 113-115.
- Braasch, D. and T. Soldán. 1988. Heptageniidae aus Nordkorea (KVDR), nebst einigen Bemerkungen zu ihrem generischen Status (Insecta, Ephemeroptera). *Faun. Abh. Mus. Tierkd. Dresden* 12: 28.
- Burmeister, H.C.C. 1839. Handbuch der Entomologie. Vol. 2, Part 2 *Neuroptera*. Reimer, Berlin, xii: 757-1050.
- Chino, Y. 1975. Madarakagerouka isshunituite. [On the notes of Japanese species of Ephemerellidae]. *Tansueibutsu* 13: 10-11. [in Japanese]
- Curtis, J. 1834. Descriptions of some nondescript British species of mayflies of anglers. *London and Edinburgh Philo. Mag. J. Sci.* 3(4): 120-125, 212-218.
- Dodds, G.S. 1923. Mayflies from Colorado, description of certain species and notes on others. *Trans. Am. Entomol. Soc.* 69: 93-116.
- Eaton, A.E. 1868. An outline of a re-arrangement of the genera of the Ephemeridae. *Entomol. Month. Mag.* 5: 82-91.
- Eaton, A.E. 1869. On *Centroptilum*, a new genus of the Ephemeridae. *Entomol. Month. Mag.* 6: 131-132.
- Eaton, A.E. 1870. On some new British species of Ephemeridae. *Trans. Ent. Soc. Lond.* 1870: 1-8.
- Eaton, A.E. 1871. A monograph on the Ephemeridae. *Trans. Ent. Soc. Lond.* 1871: 1-164, pl. 1-6.
- Eaton, A.E. 1881a. An announcement of new genera of the Ephemeridae. *Entomol. Month. Mag.* 17: 191-197.
- Eaton, A.E. 1881b. An announcement of new genera of the Ephemeridae. *Entomol. Month. Mag.*

- 18: 21-27.
- Eaton, A.E. 1885 (1883-1888). A revisional monograph of recent Ephemeroidae or mayflies. *Trans. Linn. Soc., 2nd Ser. Zool.* 3: 1-346.
- Eaton, A.E. 1892. On two new and some other Japanese species of Ephemeroidae. *Entomol. Month. Mag.* 38: 302-303.
- Eaton, A.E. 1907-1908. Description of a new species of Ephemeroidae. pp. 149-151 In: *Bidrag til en Fortegnelse over arkisk norges Neuropter fauna*. by Esben Petersen. Tromsø Museums Aarshefter 25(1902): 119-153.
- Edmunds, G.F., Jr. 1959. Subgeneric groups within the mayfly genus *Ephemerella* (Ephemeroptera: Ephemerellidae). *Ann. Entomol. Soc. Amer.* 52: 543-547.
- Edmunds, G.F., Jr. 1960. Two generic synonymy in the Siphlonuridae (Ephemeroptera). *Bull. Brooklyn Entomol. Soc.* 55: 24.
- Edmunds, G.F., Jr. and R.D. Waltz. 1996. Ephemeroptera. pp. 126-163 In: Merritt, R.W. and K.W. Cummins (eds). *An introduction to the aquatic insects of North America*. 3rd ed. Kendall/Hunt Publishing Company, USA.
- Entomological Laboratory, Faculty of Agriculture, Kyushu University. 1990. Ephemeroptera. pp. 20-23 In: Entomological Laboratory, Faculty of Agriculture, Kyushu University and Japan Wildlife Research Center (eds). *A Checklist of Japanese Insects*. Addenda et corrigenda. Isseido Printing Co., Fukuoka.
- Futai, K. 1963. Ishigaki-jima no Suiseikontyu [Aquatic insects from Ishigaki-jima ls.]. *Kansai Shizenkagakushi* 20: 39-41. [in Japanese]
- Gose, K. 1962. Ephemeroptera. pp. 6-24 In: Tsuda, M. (ed.). *Aquatic Entomology*. Hokuryukan, Tokyo. [in Japanese]
- Gose, K. 1963. Two new mayflies from Japan. *Kontyū* 31: 142-145.
- Gose, K. 1965. Description of two new species of Baetidae from Japan (Ephemeroptera). *Kontyū* 33: 218-220.
- Gose, K. 1968. Two new mayflies from Japan. *Kontyū* 36: 147-150.
- Gose, K. 1979a. The mayflies of Japanese. General parts. *Aquabiology* 1: 38-44. [in Japanese]
- Gose, K. 1979b. The mayflies of Japanese. (1) Key to families, genera and species. *Aquabiology* 2: 40-45. [in Japanese]
- Gose, K. 1979c. The mayflies of Japanese. (2) Key to families, genera and species. *Aquabiology* 3: 58-60. [in Japanese]
- Gose, K. 1979d. The mayflies of Japanese. (3) Key to families, genera and species. *Aquabiology* 4: 43-47. [in Japanese]
- Gose, K. 1979e. The mayflies of Japanese. (4) Key to families, genera and species. *Aquabiology* 5: 51-53. [in Japanese]
- Gose, K. 1980a. The mayflies of Japanese. (5) Key to families, genera and species. *Aquabiology* 6: 76-79. [in Japanese]
- Gose, K. 1980b. The mayflies of Japanese. (6) Key to families, genera and species. *Aquabiology* 7: 122-123. [in Japanese]
- Gose, K. 1980c. The mayflies of Japanese. (7) Key to families, genera and species. *Aquabiology* 8: 211-215. [in Japanese]
- Gose, K. 1980d. The mayflies of Japanese. (8) Key to families, genera and species. *Aquabiology* 9: 286-288. [in Japanese]
- Gose, K. 1980e. The mayflies of Japanese. (9) Key to families, genera and species. *Aquabiology* 10: 366-368. [in Japanese]
- Gose, K. 1980f. The mayflies of Japanese. (10) Key to families, genera and species. *Aquabiology* 11: 454-457. [in Japanese]
- Gose, K. 1980g. The mayflies of Japanese. (11) Life histories. *Aquabiology* 12: 59-62. [in Japanese]

- Gose, K. 1981. A revision of the genus *Ephemera* (Ephemeroptera) in Japan. *Biol. Inl. Wat.* 2: 11-14. [in Japanese]
- Gose, K. 1985. Ephemeroptera. pp. 7-32 In: Kawai, T. (ed.). *An Illustrated Book of Aquatic Insects of Japan*. Univ. Tokai Press. Tokyo. [in Japanese]
- Gose, K. 1988. Ephemeroptera. pp. 7-32 In: Kawai, T. (ed.). *An Illustrated Book of Aquatic Insects of Japan*. 3rd ed., Univ. Tokai Press. Tokyo. [in Japanese]
- Hagen, H.A. 1858. Synopsis der Neuroptera Ceylons. *Verh. Zool. Bot. Ges. Wien.* 8: 471-488.
- Hashimoto, I. 1955. Studies on some mayflies in Saitama district (1). *Bull. Nat. Mus. Chichibu.* 67-79. [in Japanese]
- Hatta, K. and S. Ishiwata. 1990. Fauna of aquatic insects in the major rivers of the Tokai District, Central Japan (III). Ephemerellidae (Ephemeroptera) in the major rivers of the Tokai District. *J. Nagoya Women's Univ.* 36: 167-178. [in Japanese]
- Horasawa, I. 1929. Description of *Epeorus latiforium* Uéno. *Kontyû* 3: 253-258. [in Japanese]
- Horasawa, I. 1931. Short notes on Japanese Ephemeridae (2). *Kontyû* 5(3): 121-124. [in Japanese]
- Hubbard, M.D. 1988. *Promatsumura*, replacement name for *Procloeon* Matsumura, 1931 (Ephemeroptera: Baetidae) with designation of type-species. *Insecta Mundi* 2: 240.
- Hubbard, M.D. 1989. *Matsumuracloeon*, a replacement name for *Pseudocloeon* Matsumura, 1931 (Ephemeroptera: Baetidae). *Flor. Entomol.* 72: 388.
- Hubbard, M.D. 1990. *Mayflies of the World. A Catalog of the Family and Genus Group Taxa (Insect: Ephemeroptera)*. Flora & Fauna Handbook 8. Sandhill Crane Press, INC. Gainesville, Florida.
- Hwang, J.M. and Y.J. Bae. 1999. Systematics of the Caenidae (Ephemeroptera) in Korea. *Korean J. Entomol.* 29(3): 239-245.
- Imanishi, K. 1930. Mayflies from Japanese Torrents. New mayflies of the genera *Acentrella* and *Ameletus*. *Trans. Nat. Hist. Soc. Formosana* 20: 265-266.
- Imanishi, K. 1932. Mayflies from Japanese Torrents II. Further notes on the genus *Ameletus*. *Annal. Zool. Japon.* 13: 525-533.
- Imanishi, K. 1933. Mayflies from Japanese Torrents III. Third notes on the genus *Ameletus* with a list of the Japanese Siphlonuridae. *Insecta Matsumurana* 8(2): 64-69.
- Imanishi, K. 1934. Mayflies from Japanese Torrents IV. Notes on the genera *Epeorus*. *Annal. Zool. Japon.* 14: 380-394.
- Imanishi, K. 1935. Mayflies from Japanese Torrents V. Notes on the genera *Cinygma* and *Heptagenia*. *Annal. Zool. Japon.* 15: 213-223.
- Imanishi, K. 1936. Mayflies from Japanese Torrents VI. Notes on the genera *Ecdyonurus* and *Rhihogena*. *Annal. Zool. Japon.* 15: 538-549.
- Imanishi, K. 1937a. Mayflies from Japanese Torrents VII. Notes on the genus *Ephemerella*. *Annal. Zool. Japon.* 16: 321-339.
- Imanishi, K. 1937b. Mayflies from Japanese Torrents VIII. Notes on the genera *Paraleptophlebia* and *Baetis*. *Annal. Zool. Japon.* 16: 330-337.
- Imanishi, K. 1939. Mayflies from Japanese Torrents X. Life forms and life zones of mayfly nymphs. II. Ecological structure illustrated by life zone arrangement. *Mem. Coll. Sci., Kyoto Imp. Univ. ser B* 14: 1-35.
- Imanishi, K. 1940. Ephemeroptera of Manchuria, Inner Mongolia, and Korea. *Rep. Limnol. Surv. Kwant. Manch.* 1940: 169-263. [in Japanese]
- Imhoff, L. 1852. Bericht über Verhandl. d. naturf. Gesellsch. Basel X: 177-180.
- International Commission on Zoological Nomenclature. 1999. *International Code of Zoological Nomenclature*. 4th ed.
- Ishiwata, S. 1987. Structure and keys of the family Ephemerellidae (1). Structure and keys to genera from the family Ephemerellidae. *Aqu. Org. Kanagawa Pref.* 9: 27-34. [in Japanese]

- Ishiwata, S. 1993. A checklist of the family Ephemeralidae (Ephemeroptera) of Japan, Korea, and Far East of Russia. *Report of the Studies on the Structure and Function of River Ecosystems of the Far East* 2: 5-19.
- Ishiwata, S. 1996. A study of the genus *Ephoron* from Japan (Ephemeroptera, Polymitarcyidae). *Cand. Entomol.* 128: 551-572.
- Ishiwata, S. 1997. Mayflies from the Tanzawa mountains. pp. 290-296 In: *Fauna and Flora of the Tanzawa mountains*. Comprehensive Research Report on Natural Environment of Tanzawa Mountains. Kanagawa Prefectural Government. [in Japanese]
- Ishiwata, S. 1999. Mayflies of Rishiri-to Island. *Rishiri Studies* 18: 21-27. [in Japanese]
- Ishiwata, S. 2000. Notes on mayflies in Kanagawa Prefecture. *Nat. Hist. Rep. Kanagawa* 21: 73-82 [in Japanese]
- Ishiwata, S. and K. Inada. 1996. Mayflies common in Japan. *Biol. Inl. Wat.* 11: 36-44. [in Japanese]
- Ishiwata, S., T.M. Tiunova, and R.B. Kuranishi. 2000. The mayflies (Insecta: Ephemeroptera) collected from the Kamchatka Peninsula and the North Kuril Islands in 1996-1997. In: Komai, T. (ed.), *Results of Recent Research on Northeast Asian Biota*. Natural History Research, Special Issue 7: 67-75. Tokyo.
- Itou, M. 1989. Aquatic insects of Kushiro Marsh I, Ephemeroptera, Trichoptera and Planipennia. *Sylvicola* 7: 53-57.
- Kang, S., H. Chang, and C. Yang. 1994. A revision of the genus *Baetis* in Taiwan (Ephemeroptera, Baetidae). *J. Taiwan Mus.* 47(2): 9-44.
- Kang, S. and C. Yang. 1994a. Heptageniidae of Taiwan (Ephemeroptera). *J. Taiwan Mus.* 47(1): 5-36.
- Kang, S. and C. Yang. 1994b. Leptophlebiidae of Taiwan (Ephemeroptera). *J. Taiwan Mus.* 47(1): 57-82.
- Kang, S. and C. Yang. 1995. Ephemeralidae of Taiwan (Insecta, Ephemeroptera). *Bull. Nat. Mus. Nat. Sci.* 5: 95-116.
- Kazlauskas, R. 1963. New and little known mayflies (Ephemeroptera) from the USSR. *Entomol. Obozr.* 42: 582-589. [in Russian]
- Kimmins, D.E. 1960. The Ephemeroptera types of species described by A.E. Eaton, R. McLachlan and F. Walker, with particular reference to those in the British Museum (Natural History). *Bull. British Mus. (Nat. Hist.) Entomol.* 9(5): 269-318.
- Klapálek, F. 1905. Plecopteran und Ephemeren aus Java, gesammelt von Pro. K. Klaepelin 1904. *Mit. Natur. Mus. Hamburg* 22: 103-107.
- Kluge, N.Ju. 1980. To the knowledge of mayflies (Ephemeroptera) of Taimyr national district. *Entomol. Obozr.* 59(3): 561-579. [in Russian]
- Kluge, N.Ju. 1983. New and little known mayflies of the family Baetidae (Ephemeroptera) from the Premor'ye Territory. *Entomol. Obozr.* 61: 65-79. [in Russian]
- Kluge, N.Ju. 1984. Mayflies of the subgenus *Euthraulus* Barnard (Ephemeroptera, Leptophlebiidae, Genus *Choroterpes*) of the fauna of the USSR. *Entomol. Obozr.* 63: 722-728. [in Russian]
- Kluge, N.Ju. 1988. Revision of genera of the family Heptageniidae (Ephemeroptera). 1. Diagnoses of tribes, genera and subgenera of the subfamily Heptageniinae. *Entomol. Obozr.* 67: 291-313.
- Kluge, N.Ju. 1994. Pterothorax structure of mayflies (Ephemeroptera) and its use in systematics. *Bull. Soc. Entomol. Franc.* 99: 41-61.
- Kluge, N.Ju. 1997. Ephemeroptera. pp. 177-220, 304-328 In: Tsololikhin (ed.). *Key to Freshwater Invertebrates of Russia and Adjacent Lands*. St. Petersburg. [in Russian]
- Kluge, N.Ju. 1998. Phylogeny and higher classification of Ephemeroptera. *Zoosyst. Rossica*. 7(2): 255-269.

- Kluge, N.Ju., D. Studemann, P. Landolt, and T. Gonsern. 1995. A reclassification of Siphlonuroidea (Ephemeroptera). *Mitt. Schweiz. Ent. Ges.* 68: 103-132.
- Kobayashi, N. 1986. Biology of Japanese mayflies. *Insectarium* 23(7): 18-23. [in Japanese]
- Kobayashi, N. 1987. Kankyoushihyou tositeno Kokagerou [Baetis, indicator of water environment]. pp. 41-60 In: Yasuno, M. & T. Iwakuma (eds). *Symposium, Suiki ni okeru Seibusuhiyou no Mondaiten to Syourai*. National Institute Environmental Studies. Ibaraki. [in Japanese]
- Kobayashi, N. 1992. Ephemeroptera. pp. 18-24 In: Moriya, H. (ed.). *Sagamihara no Teiseidoubutsu II*. Sagamihara City Government. [in Japanese]
- Kobayashi, N. and M. Nishino. 1992. Ephemeroptera. pp. 4-17 In: Nishino, M. (ed.). *Biwako no Teiseidoubutsu II*. Shiga Prefectural Government. [in Japanese]
- Kondratieff, B.C. and J.R. Voshell, Jr. 1984. The North and Central American species of *Isonychia* (Ephemeroptera: Oligoneuriidae). *Trans. Amer. Entomol. Soc.* 110: 129-244.
- Kuhara, N., N. Kobayashi, Y. Nagayasu, and T. Ito. 1993. Aquatic insect fauna of the Naibetsu River, Chitose, Hokkaido, northern Japan (A preliminary report). *Biol. Inl. Wat.* 8: 15-20. [in Japanese]
- Kushiro Insect Lover's Society, 1995. Insects of Kushiro Marsh. *Sylvicola supplement* 2: 168. [in Japanese]
- Leach, W.E. 1815. Entomology. *Brewster's Edinburgh Encycl.* 9: 57-172.
- Lestage, J.A. 1917. Contribution à l'étude des larves de Ephéméroptères paléarctiques. *Ann. Biol. Lacust.* 8: 215-458.
- Lestage, J.A. 1930. La dispersion holactique de quelques Ephéméroptères. *Bull. Ann. Soc. Entomol. Beig.* 70: 201-207.
- Lestage, J.A. 1938. Contribution à l'étude des Ephéméroptères. *Bull. Ann. Soc. Entomol. Beig.* 78: 381-394.
- Linnaeus, C. 1758. *Systema Naturae*. Ed. X. Holmae. Vol. 1.
- Linnaeus, C. 1761. Fauna Suecica. ed.ii. no. 1474.
- Linnaeus, C. 1767. *Ephemera*. pp. 906-907 In: C. Linnaeus. *Systema Naturae*. 12th Ed., Pl. 2, Nos. 1-11.
- Makibayashi, I. 1998. Saitamaken no Kageroumoku [Ephemeroptera]. pp. 9-20 In: Makibayashi, I. (ed.). *Saitamaken Kontyishi I [Insects of Saitama Japan I]*. Saitama Kontyû Danwakai. [in Japanese]
- Malzacher, P. 1996. *Caenis nishinoae*, a new species of the family Caenidae from Japan (Insecta: Ephemeroptera). *Stuttgarter Beitr. Natur. Ser. A. Nr. 547*: 1-5.
- Matsumura, S. 1904. Ephemeraida, 1000 Illustrated Insect of Japan Empire: 19-20, 159-161. fig. 6, pl. 10. Keiseisha Shoten. Tokyo. [in Japanese]
- Matsumura, S. 1911. Erster Beitrag zur Insekten-Fauna von Sachalin. *J. Coll. Agri. Tohoku Univ.* 4: 1-145.
- Matsumura, S. 1931a. Ephemeraida. pp. 1466-1480 In: *6000 Illustrated Insect of Japan Empire*. Toko-shoin. Tokyo. [in Japanese]
- Matsumura, S. 1931b. New genera and new species described in "6000 Illustrated Insect of Japan Empire". *Insecta Matsumurana* 6: 199-200. [in Japanese]
- Matsumura, S. 1933. *Illustrated Common Insect of Japan*. V. Neuroptera / Orthoptera. Odonata: pp. 60-67, pl. 20-21. Shunyodo, Tokyo. [in Japanese with English description]
- McCafferty, W.P. 1996. The Ephemeroptera species of North America and index to their complete nomenclature. *Trans. Amer. Entomol. Soc.* 122: 1-54.
- McCafferty, W.P. and R.D. Waltz. 1995. *Labiobaetis* (Ephemeroptera: Baetidae) new status, new North American species, and rerated new genus. *Entomol. News* 106: 19-28.
- McDunough, J. 1924. New North American Ephemeraidae. *Canad. Entomol.* 56: 221-226.
- McDunough, J. 1933. The nymph of *Cinygma interrum* and description of a new heptagenine

- genus. *Canad. Entomol.* 65: 73-76.
- McLachlan, R. 1875. A sketch of our present knowledge of the Neuropterous fauna of Japan (excluding Odonata and Trichoptera). *Trans. Entomol. Soc.* 2: 167-190.
- Morishita, I. 1961. Yakushima no Suiseikontyu [Aquatic insects from Yakushima Is.]. *Kansai Shizenkagakuishi* 14: 4-10. [in Japanese]
- Morgan, A. H. 1911. Mayflies of Fall Creek. *Ann. Entomol. Soc. Amer.* 4: 93-119.
- Müller-Liebenau, I. 1985. Baetidae from Taiwan with remarks on *Baetiella* Ueno, 1931 (Insecta, Ephemeroptera). *Arch. Hydrobiol.* 104: 93-110.
- Nakamura, M. 1941. A list of insects collected with trap-apparatus at Utsunomiya V. *Kontyū* 15(1): 37-43. [in Japanese]
- Nakasone, K., H. Mitsumoto, M. Yonamine, T. Kishimoto, Y. Higa, and T. Omija, 1998. Studies on suspended material content in sediment and benthic macro-invertibrate fauna of river in U. S. Military Base. *Ann. Rep. Okinawa Pref. Inst. Heal. Env.* 32: 161-167.
- Navás, L. 1912. Quelques Nevroptères de la Sibérie méridionale-orientale. *Entomol. Obozr.* 12: 414-422.
- Navás, L. 1915. Neue Neuropteren. Este Serie. *Ent. Mitt.* 4: 146-153.
- Navás, L. 1919. Neuropteros (Ins.) del Japon. *Racmad.* 18: 157-164.
- Navás, L. 1933. Insecta orientalia. XII series. *Mem. Soc. Accad. Sci., Nuovi Lincei* (2)17: 75-18.
- Navás, L. 1934. Insectti Neurotteri e affini del Piemonte *Mem. Soc. Entomol. Ital.* 1933 12: 150-162.
- Needham, J.G. 1909. A peculiar new mayfly from Scandage Park. *New York State Mus. Bull.* 134: 71-75.
- Novikova, E.A. and N.Ju. Kluge. 1987. Systematics of the genus *Baetis* (Ephemeroptera, Baetidae) with description of a new species from Middle Asia. *Vestn. Zool.* 4: 8-19.
- Okazaki, H. 1984. Observation of the eggs of Japanese mayflies by scanning electron microscope. *Biol. Inl. Wat.* 3: 19-27. [in Japanese]
- Pactl, J. 1994. *Ephacerella*, a replacement name for *Acerella* Allen, 1971 (Ephemeroptera), nec Berlesia, 1909 (Protura). *Ent. News* 105(5): 283-284.
- Pictet, F.J. 1843-1845. Histoire naturelle générale et particulière des insectes Névroptère. Famille des Ephémérines. Kessmann & Cherbuliez, Geneve: 1-300, 49 pl.
- Poda, N. 1761. Insecta Musaei Graecensis: 97-98.
- Retzius, A.J. 1783. Caroli De Geer genera et species insectorum et generalissimi autoris scriptis extraxit, digessit, latine quand. Partem reddidit, et treminologiam insectorum Linneanam addidit. Druse, Lipsiae, vi+220p.
- Rostock, M. 1877. Die Ephemeren und Psociden Sachsens mit Berücksichting der meisten ubrigen deutshen arten. *Jber. Ver. Naturk. Zwickau* 15: 76-100.
- Sato, T. 1969. Succession of aquatic insects with the clarification of water in Sandankyo Gorge. pp.124-188 In: Education Board of Hiroshima Pref. (ed.). *Limnological researches of the Sandankyo District*. Hiroshima. [in Japanese]
- Sinitshenkova, N.D. 1978. A redescription of larval and imaginal stages of *Isonychia aesculus* (Imanishi) 1934 and the evolution of the gill apparatus in the larvae of the g. *Isonychia* Eaton, 1881 (Ephemeroptera, Heptageniidae). *Bull. Mosk. Obschch. Ispyt. Prir.* 83(5): 49-56. [in Russian]
- Stephens, J.F. 1835. Mandibulata. Illustrations of British Entomology. London 6: 53-70. pl. 29.
- Takahashi, Y. 1924. Some Japanese mayflies. *Annat. Zool. Japon* 36: 377-380. [in Japanese]
- Takahashi, Y. 1929. Two new mayflies from Kagawa-ken, Japan. *Lansania* 1(5): 75-80.
- Takahashi, Y. 1930. A new species of mayfly and *Cloeon dipterum* L. *Insect World* 34: 114-118.
- Takara, T. and S. Azuma. 1972. Fauna of the Iriomote Island, Ryukyus. II List of the insects (1) Order Protura, Collembola, Thysanura, Ephemeroptera, Odonata and Plecoptera 19: 87-97. [in Japanese]

- Takemon, Y. 1990. Mayfly fauna of Kyoto Prefecture - Its characteristics and taxonomical problems. *Sci. Eng. Rev. Doshisha Univ.* 31(1): 49-63. [in Japanese]
- Tanaka, H. 1966. Ecological studies an aquatic insects in upper reaches of the Kinu-gawa River, Tochigi Prefecture. *Bull. Fresh. Fish. Res. Lab.* 15(2): 123-147. [in Japanese]
- Tanida, K. 1974. Preliminary reports on the aquatic insects of the Ryukyus with some ecological remarks 1. In: Ikehara, S. (ed.). *Ecol. Stud. Nat. Ryukyus* 1: 161-174. [in Japanese]
- Traver, J.R. 1934. New North American species of mayflies (Ephemerida). *Elisha Mitchel Sci. Soc. J.* 50: 189-254.
- Tshernova, O.A. 1949. Mayfly nymphs from lake Teletskoye and River Bija. *Trudy Zool. Inst. Akad. Nauk. SSSR* 7: 139-158. [in Russian]
- Tshernova, O.A. 1952. Mayflies from the River Amur and neighboring waters, and their part in the nutrition of Amur-fishes. *Trudi Amur. Ichtiol. Eksped.* 3: 229-360. [in Russian]
- Tshernova, O.A. 1972. Some new species of mayflies from Asia (Ephemeroptera, Heptageniidae, Ephemerellidae). *Entomol. Obozr.* 51: 604-614. [in Russian]
- Tshernova, O.A. and V. V. Belov. 1982. On the systematics of the adults of the Palearctic mayflies of the genus *Cinygmulia* McDunnough, 1933 (Ephemeroptera, Heptageniidae). *Entomol. Obozr.* 61: 278-296. [in Russian]
- Tshernova, O.A., N.Ju. Kluge, N.D. Sinichenkova, and V.V. Belov. 1986. Order Ephemeroptera. pp. 99-142 In: Lehr, P.A. (ed.). *Opredelitel' Nasekomykh Dal'nego Vostoka SSSR*. Nauk, Leningrad. [in Russian]
- Uchida, S. 1986. Aquatic insects of Boso Hills. *Bull. Biol. Soc. Chiba* 36: 1-15. [in Japanese]
- Ueno, T. 1995. Nishiharimachihou niokeru Togeerakagerou-zoku 2syu ni tsuite [On the larvae of genus *Thraulus* from Nishi-harima District, central Japan]. *Hyogo Fresh-water Biology* 46: 25-27. [in Japanese]
- Uéno, M. 1928. Some Japanese mayfly nymphs. *Mem. Coll. Sci. Kyoto Imp. Univ.*, ser. B, 4, 1, Art.2: 19-63, pls.3-17, Figs 1-18.
- Uéno, M. 1931a. Einige neue Ephemeropteren und Plecopteren aus Mittel-Japan. *Annat. Zool. Japon.* 13: 91-99.
- Uéno, M. 1931b. A new Japanese mayflies. *Lanzania* 3(24): 57-59.
- Uéno, M. 1931c. Contributions to the knowledge of Japanese Ephemeroptera. *Annat. Zool. Japon.* 13: 189-231.
- Uéno, M. 1938. Stoneflies and mayflies collected in Okinawa, Ryukyu Islands. *Biogeographica-Trans.* 3(1): 92-99. [in Japanese]
- Uéno, M. 1950. Ephemeroptera. pp. 120-130 In: Ishii, T. et al. (eds). *Nippon Kontyu Zukan*. Hokuryukan. Tokyo. [in Japanese]
- Uéno, M. 1969. Mayflies (Ephemeroptera) from various regions of Southeast Asia. *Oriental Insects* 3(3): 221-238.
- Uéno, M. 1973. Ephemeroptera. pp. 515-525 In: Uéno, M. (ed.). *Freshwater Biology of Japan*. Hokuryukan, Tokyo. [in Japanese]
- Uéno, M. and H. Okamoto. 1932. Ephemeroptera. pp. 1950-1962 In: Uchida, S. et al., (eds). *Nippon Kontyu Zukan*. Hokuryukan. Tokyo. [in Japanese]
- Ulmer, G. 1912. H. Sauter's Formosana-Ausbeute. Ephemeren. *Ent. Mitt.* 1(12): 369-375.
- Ulmer, G. 1919 (1920). Neue Ephemeropteren. *Arkiv. Naturgesch.* 85(11): 1-80.
- Ulmer, G. 1924. Einige alte und neue Ephemeropteren. *Konowia* 3: 23-37.
- Ulmer, G. 1927. Entomologische Ergebnisse der schwedischen Kamtchatka-expedition 1920-1922. II. Trichopteren und Ephemeropteren. *Arkiv. Zoologi* 19(8): 1-17.
- Walsh, B.D. 1863. List of the Pseudoneuroptera of Illinois contained in the cabinet of the writer, with descriptions of over forty new species, and notes on their structural affinities. *Proc. Acad. Nat. Sci. Philadelphia* 1862: 361-402.
- Waltz, R.D. and W.P. McCafferty. 1987. Systematics of *Pseudocloeon*, *Acentrella*, *Baetiella*, and

- Liebebiella*, new genus (Ephemeropreta: Baetidae). *J. New York Entomol. Soc.* 95(4): 557-568.
- Waltz, R.D., W.P. McCafferty, and A. Thomas. 1994. Systematics of *Alainites* n. gen., *Diphotor*, *Indobaetis*, *Nigrobaetis* n. stat., and *Takobia* n. stat. (Ephemeroptera, Baetidae). *Bull. Soc. Hist. Nat. Toulouse* 130: 33-36.
- Williamson, H. 1802. On the *Ephoron leukon* usually called the white fly of the Passaic River. *Trans. Amer. Philo. Soc.* 5: 71-73.
- Wu, X.-Y. 1987. A new species of *Potamanthus* (Ephemeroptera: Potamanthidae). *Acta Zootaxonomica Sinica* 12(4): 421-423. [in Chinese]
- Yamasaki, T. 1987. Ephemeroptera of the Tamagawa River System. In: Analytic studied on the distribution of some insect-groups in the Tamagawa River System and its upper reaches. Tokyu Foundation for the Better Environment: 81-120+3pls. [in Japanese]
- Yokkaichi City Government, 1990. 15 Suiseikonchu Rui. pp. 348-354. Yokkaichi-shi Shishi '1. Yokkaichi City Government. [in Japanese]
- Zurwerra, A. and I. Tomka. 1985. *Electrogenera* gen. nov., eine neue Gattung der Heptageniidae (Ephemeroptera). *Entomol. Ber. Luzern* 13: 99-104.