

THE CRATO FOSSIL BEDS OF BRAZIL

Window into an Ancient World

Edited by

DAVID M. MARTILL

University of Portsmouth

GÜNTER BECHLY

Staatliches Museum für Naturkunde, Stuttgart

ROBERT F. LOVERIDGE

University of Portsmouth



CAMBRIDGE
UNIVERSITY PRESS

CAMBRIDGE UNIVERSITY PRESS
Cambridge, New York, Melbourne, Madrid, Cape Town, Singapore, São Paulo

Cambridge University Press
The Edinburgh Building, Cambridge CB2 8RU, UK

Published in the United States of America by Cambridge University Press, New York

www.cambridge.org
Information on this title: www.cambridge.org/9780521858670

© Cambridge University Press 2007

This publication is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without
the written permission of Cambridge University Press.

First published 2007

Printed in the United Kingdom at the University Press, Cambridge

A catalogue record for this publication is available from the British Library

ISBN 978-0-521-85867-0 hardback

Cambridge University Press has no responsibility for the persistence or
accuracy of URLs for external or third-party internet websites referred to
in this publication, and does not guarantee that any content on such
websites is, or will remain, accurate or appropriate.

Crato fossils

The first records of Zygentoma from the Crato Formation are those of Bechly (1998a). Two specimens of Lepismatidae were described by Sturm (1998). Since then, a further undescribed specimen (Plate 7a) was found in the Stuttgart (SMNS) collection. Due to their relatively poor state of preservation, neither a determination of the sex, nor a formal taxonomic description as a new species, has been possible. Consequently, these specimens have only been determined as “Lepismatidae gen. spec. ‘Araripe’” by Sturm (1998).

Lepismatidae

Lepismatidae gen. spec. ‘Araripe’ Sturm, 1998

Material: no. B 99 at SMF; no 1998 III/4 at BSPGM; and no. SMNS 66535 (Plate a). A fourth specimen without number from AMNH was figured by Grimaldi and Engel (2005: 152, figure 5.6).

Diagnosis: body 10.5–14 mm long and 3–4 mm wide; head orthognathous; filiform antennae 10.5–13.5 mm long; filiform cerci (12–18 mm long, the only 8.8-mm-long cerci in specimen B 99 are obviously broken off) and terminal filum (13–21 mm long), all provided with setae; wingless; body robust; legs stout with flattened, broad and oval-shaped coxae.

11.3 Persisting-type stem group Ephemeroptera

Rainer Willmann

Organisms do not evolve at equal rates. While many Recent taxa are of entirely modern appearance, others are plesiomorphic in many respects, and in some insects plesiomorphic structures determine their body plan. This is not necessarily related to the age of the respective taxa, but of course some taxa, be it species or large species groups, have become separated from their sister group only recently, while others are very old and have changed little, even over several tens of millions of years. In both cases the latter types of organism have been called living fossils, a term introduced by Charles Darwin. However, such types have lived at any time in the history of life. As the term living fossil cannot well be applied to relatively plesiomorphic fossil species or species groups, the term persisting type, first introduced by Huxley, may be applied. Such types belonging to ephemeropteroid insects in its widest sense were in existence in the Lower Cretaceous of Brazil (Figures 11.2 and 11.3; Plates 7c–h).

Numerous mayflies of the crown group Ephemeroptera (about 3,100 described Recent species) have been described from Triassic, Jurassic, Cretaceous and Cenozoic strata including the Crato Formation (see review by McCafferty, 1990). No




Fig. 11.2. Stem group Ephemeroptera; *Cretereisma* spp. wing venation of holotypes: (a) *Cretereisma antiqua* sp. nov., SMNS 66546, left fore- and hind wings (right wings of the living animal); (b) same specimen as (a), showing basis of left forewing (right wing of the living animal) with additional structures as observed in the right wing; (c) *Cretereisma schwickerstorium*, SMNS 66598. Right forewing (left forewing of the living animal). Note the costal brace which does not reach the anterior wing margin. Both original specimens are seen in ventral view and convex veins are preserved as concave and vice versa. Drawings by R. Willmann.

such form is known from the Palaeozoic. On the other hand, numerous stem group representatives of Ephemeroptera are known from the Carboniferous and Permian, and it has been assumed that most of them, and all of those with more than seven pairs of nymphal abdominal gills, were extinct by the end of the Permian.

During a meeting of the fossil-insects network of the European Scientific Foundation in Portsmouth in 1998, David Martill exhibited specimens of a nymph from




Fig. 11.3. Stem group Ephemeroptera; *Cretereisma* sp. nov., anatomical details of nymphs. (a) Forewing pad and proximal part of hind wing pad, SMNS 66673. (b) First abdominal left gill plate overlying femur-tibia-joint of hind leg. Its basal area is overlain by the tip of the hind wing pad. The dark area near the left margin is the posterior part of the coxa, SMNS 66673. (c) Bases of terminal appendages, SMNS 66601. (d) Left circus (on the right) and paracercus, showing setation, SMNS 66604. (e) *Cretereisma* sp., nymph, no. 512 MURJ. Photographs (a)–(d) by R. Willmann, photograph (e) by G. Bechly.

the Crato Beds that appeared to belong in mayflies but clearly not into any taxon previously known from the late Mesozoic. Comparison with stem group representatives of mayflies showed that the new finds belong here. Later, Günter Bechly obtained specimens of large adult mayflies that showed characters that were as plesiomorphic as those of the nymphs. I completely agree with Bechly *et al.* (2001a), who discussed and figured beautiful larval and adult specimens as ancestral mayflies similar to *Prottereismatidae*, that both nymphs and adults belong to the same archaic taxon. These specimens are described below.

Systematics

Including ephemeroid-like insects, the mayflies have been classified into the crown group and the stem group representatives. An early example is Handlirsch (1908: 1292 and plate LX), who distinguished between a subclass Ephemeroidea including the Plectoptera (= Agnatha), and the Protephemeroidea. Crampton (1938: 170) wrote that the Ephemerida were probably derived from the Protephemeroidea, and that the Ephemerida and Protephemeroidea might be grouped into a superorder called the Panephemeroptera or Ephemeroptera. The names Protephemeroptera and Protephemeroidea refer to the Carboniferous *Triplosoba pulchella* (Brongniart, 1893), which is, however, possibly not a stem group representative of the mayflies (Willmann, 1999). Therefore, one might abandon the names Panephemeroptera or Ephemeria. Lauterbach (1973), however, proposed to call a crown group and the entirety of its stem group representatives with the prefix Pan-, and accordingly, under a phylogenetic view, the name Panephemerida would still be both useful and acceptable. Numerous stem group representatives of the mayflies have been described since the discovery of *Triplosoba*.

Categorical ranks have varied from subclass (Ephemeroidea *sensu* Handlirsch, 1908) through superorder (Ephemeroidea *sensu* Weber, 1933), cohorts (Ephemeriformes; Rohdendorf, 1977) to order (Ephemeroptera), suborder and infraorder. The latter two ranks have been applied to the crown group mayflies, or Euplectoptera, which were, however, consistently called Ephemeroptera, Ephemerida or Plectoptera when only Recent species were considered. Carpenter (1992) called the mayflies *sensu latissimo* (including *Triplosoba*) Ephemeroptera and assigned them ordinal rank. Categorical ranks will not be used here because of the inapplicability in a phylogenetic context.

Several taxon names for stem group representatives do not refer to monophyla and should be abandoned (for Permoplectoptera, see remark of Illies, 1968: 6). Willmann (2006) reviewed the phylogenetic relationships among the early mayflies *sensu lato* and systematized the stem group representatives, including *Cretereisma*,

as follows, attempting to preserve established names (synonyms are given only as examples).

1. Panephemeroptera Crampton (= *Ephemerata* Boudreaux; = *Ephemeridea* Rasnitsyn, 2002 (in Rasnitsyn and Quicke, 2002), = *Ephemeroptera* *sensu* Tshernova, 1962, Carpenter, 1992 and others, = *Ephemerida* *sensu* Kukalová-Peck, 1985)
 - ? 1.1 *Triplosoba* ('Protephemeroptera')
 - 1.2.1 *Bojophlebia*
 - 1.2.2 *Ephemerontoida* (= *Ephemerida* *sensu* Boudreaux, 1979: 271)
 - 1.2.2.1 *Syntonoptera*
 - 1.2.2.2 *Ephemeronta*
 - 1.2.2.2.1 *Lithoneura*
 - 1.2.2.2.2 *Reticulata* (= *Plectoptera* *sensu* Tshernova, 1962, 1970, = *Ephemerida* *sensu* Kluge and Sinitshenkova, 2002 (in Rasnitsyn and Quicke, 2002) and others, = *Ephemeroptera* *sensu* Grimaldi and Engel, 2005)
 - 1.2.2.2.2.1 *Prottereisma* (+*Misthodotes*?)
 - 1.2.2.2.2.2 *Heptabranchia*
 - 1.2.2.2.2.2.1 *Cretereisma*
 - 1.2.2.2.2.2.2 n.n.
 - 1.2.2.2.2.2.2.1 *Litophlebia*
 - 1.2.2.2.2.2 *Triangulifera* (= *Euplectoptera* *sensu* Tshernova, 1962, 1970,? = *Euplectoptera* *sensu* Tillyard, 1932) (= *Paedephemera multinervosa*+remaining ephemeroids, which include, among others, the *Ephemerida*) (*Ephemerida* = *Ephemeroptera* *vel* *Plectoptera* auct., = *Euplectoptera* *sensu* Grimaldi and Engel, 2005,? = *Euplectoptera* *sensu* Tillyard, 1932; = crown group of the entire ephemeropteran lineage).

In the following, the vernacular term ephemeroid is used for any taxon which is considered to be more closely related to modern mayflies (*Ephemerida*) than to any other Recent taxon.

Cretereisma gen. nov.

Type species: *C. antiqua* sp. nov., by present designation.

Etymology: named after the Cretaceous age and the similar fossil genus *Prottereisma*.

Diagnosis: adult fore- and hind wings of almost equal size, narrow, distal portions only slightly broader than their basal parts. Costal brace long and only slightly arched. Anal veins long, meeting the hind margin of the wings at low angle (plesiomorphies shared with *Prottereisma* and, in part, earlier branches of ephemeroids). Several longitudinal veins run pairwise and almost parallel to each other towards the wing margin, each pair consisting of a concave and a convex

vein (IRS_{2a} and R_{2b} ; RS_{3+4} and MA_1 ; IMA and MA_2 ; MP_1 and IMP ; MP_2 and CuA_1 ; $ICuA$ and CuA_2 respectively). Nomenclature of veins as developed by Till-yard (1932) for *Prottereisma*. The main CuA vein appears to be CuA_2 , while this is CuA_1 in *Prottereisma*.

Nymph caterpillar-like or gammarid-like in appearance, with undivided tarsus bearing one claw. Pro-, meso- and metathorax subequal in size, hind wing pads only slightly shorter than forewing pads. Abdomen with seven pairs of elongated plate-like gill appendages.

Comment: these generic characters – that is, characters found in both species of *Cretereisma* – are not repeated in the following descriptions of the two species.

Cretereisma antiqua sp. nov. (Figures 11.2a and b; Plate 7c)

Material: holotype SMNS 66546 (old no. 28) (figured by Bechly *et al.*, 2001a: 48, Abb. 37).

Type horizon: Lower Cretaceous, Upper Aptian, Nova Olinda Member of the Crato Formation.

Etymology: *antiquus*, Latin, old. The name refers to the very archaic structure of the species, which resembles the Permian *Prottereisma* in many aspects.

Description (Figures 11.2a and b): specimen with all four wings spread out, visible from its ventral side. Body length 18 mm. Forewing length 24 mm. Hind wing measuring about four-fifths of forewing length. Sc running towards wing apex, curving posteriorly towards R. CuA and CuP simple. A_1 short (not extending beyond the first quarter of the wing), probably simple; remaining analis difficult to trace (the veinal description applies for both the fore- and the hind wings). Abdominal segments as broad as those of the thorax, head as broad as thorax, with large bulging eyes. Antennae and abdominal appendages not preserved.

Cretereisma schwickertorum sp. nov. (Figure 11.2c; Plate 7d)

Material: holotype SMNS 66598 (old no. H51).

Type horizon: Lower Cretaceous, Upper Aptian, Nova Olinda Member of the Crato Formation.

Etymology: the species is named in honour of Mr and Mrs Schwickert, Sulzbachtal, Germany, for generously supporting scientific research on Crato Formation fossils for nearly 15 years.

Description (Figure 11.2c): left wings missing except the very bases, right hind wing very fragmentary. Body length 32 mm. Forewing length 31 mm. Wing venation as in *C. antiqua* but A_1 almost extending to the middle of the wing and with a well-developed triad. With more anal veins behind A_1 than in the previous species.

Head, thorax and abdomen of almost the same width. Antennae, legs and abdominal appendages not preserved.

Comment: the circumstance that almost no crossveins have been noticed in the costal areas of both *C. antiqua* and *C. schwickerstorium* is certainly due to preservation.

Comparison with *C. antiqua*: the differences in venation between the two species are a result of size differences, *C. schwickerstorium* being considerably larger. In the latter, more anal branches are present to strengthen the posterior basal wing area. I do not believe that the differences, especially the absence/presence of a well-pronounced triad on A₁, is due to intraspecific variation or sexual dimorphism.

Two further well-preserved adult specimens of *Cretereisma* spp. are deposited without number in the collection of SMF.

Nymphs (Figures 11.3a–d; Plates 7e–h)

According to the size and morphology of the flying stages, the nymphs of the two species described above must have been of considerable size, have had four wing pads of almost equal size and, as a necessity of the latter property, a well-developed meso- and metathorax. Many specimens fulfilling these requirements have been found in the Crato Formation and are considered to be members of *Cretereisma*. These nymphs are not so rare in the Crato Formation, and are well-known to the local quarry workers under the vernacular term *Ananas* or pineapple.

Material: there are 10 nymphs with the nos SMNS 66547 (old no. 33; figured by Bechly *et al.*, 2001a: 47, Abb. 36), SMNS 66548 (old no. L74), SMNS 66549, SMNS 66599 (old no. 30), SMNS 66600 (old no. 31), SMNS 66601 (old no. 32), SMNS 66602 (old no. H54), SMNS 66603 (old no. L73), SMNS 66604 (old no. H52), and SMNS 66605 (old no. 29); three further specimens have been recently donated to SMNS (SMNS 66673, SMNS 66674, SMNS 66675). A very nice nymph is specimen MB.I.2028 at MNB, and additional specimens are held by SMF. One of the best-preserved nymphs is specimen no. 512 in coll. MURJ (Figure 11.3e). Another nymph from AMNH was figured by Grimaldi and Engel (2005: 166, figure 6.13).

Specimen SMNS 66673 (Figures 11.3a and b; Plate 7e) was chosen as the basis of nymphal description mainly because of its good general state and the complete preservation of its left wing pads: the animal is visible from its left side. Body length 33 mm. Thoracomes and abdominal segments are very high (body not flattened), which give the animal a somewhat caterpillar-like appearance. Sclerites with numerous fine pores are present which may have been the insertion points of bristles (see description of additional material).

Description (Figures 11.3a and b): head short, about as long as prothorax, and probably with burrowing devices. Mouthparts only fragmentarily preserved.

Antenna long, with big basal segments (only a very faint impression of one antenna directed forwards is left). Pro-, meso- and metathorax of almost equal length, mesothorax slightly longer than the other two thoracomes. Thoracic terga with pronounced hind margins. Meso- and metathorax with wing pads of subequal shape, the hind wing pad being slightly shorter than that of the forewing. The forewing base is attached to the thorax as follows: laterally, the tergite has two knob-like elevations opposite to the anterior basal wing area; these elevations being set apart from the wing pad by a deep longitudinal incision. The posterior lateral area of the tergite is slightly elevated and separated by a longitudinal groove from a still more lateral elevation near the middle of the wing base. The hind margin of the wing pad runs continuously into the hind margin of the mesothoracic tergite. The situation in the metathorax is the same, but the structures are less pronounced due to the slightly smaller size of the hind wing pad. As already proposed by Bechly *et al.* (2001a: 48), it appears that the wing pads were moveable along the deep incision mentioned above. Such moveable wing pads represent a very archaic character state, which is otherwise only known from Palaeozoic pterygote larvae. Legs stout; coxae very large and broad. Fore coxa slightly shorter than the fore femur, middle and hind coxae almost as long as the respective femora. The fore legs are the longest.

The first eight abdominal segments are of subequal length, the remaining ones being considerably shorter. Segments 1–7 each with a pair of strongly scleritized elongated lateral gill appendices which are directed downwards in the fossil. Their distal parts appear to have been strongly scleritized into plate-like structures and are slightly curved forwards. Gill plates with a pronounced margin around their distal portions. The gill appendices insert laterally at the abdomen with a broad base. As the bases are irregularly crumbled it appears that this portion was not developed as a plate in the living animal. Abdominal tergites 8–10 extend more ventrally than the preceding ones. Cerci are lacking in this specimen due to preservation.

Additional material as listed above shows well-preserved long antennae and fairly long cerci and paracercus. Antennae about as long as head and thorax combined (e.g. SMNS 66599, SMNS 66600, SMNS 66547). Paracercus with row of long setae on each side, cerci with one row of setae only. Cercal joints and joints of paracercus short. Femora with strong bristles or hairs (SMNS 66599 and SMNS 66604), coxa similarly equipped (SMNS 66547; other leg segments do not show such structures possibly due to preservation of the specimens available for study). Tarsi one-segmented, claws pointed. Specimen SMNS 66674, which is partly preserved as a negative imprint, shows impressions of longitudinally directed long body hairs on the abdomen. The small pores on the sclerites of other specimens may correspond to these hairs and might have been insertion places. Gill appendages fringed with hairs.

Body lengths in the additional material range from 10.5 mm (SMNS 66605) to 26 mm (SMNS 66547 and SMNS 66604) and even 30 mm in one specimen (SMNS 66549), without antennae and abdominal appendages.

It was not possible to distinguish, in the nymphs, between the two species described above from adult specimens.

With exception of specimen no. MSF Z78, all nymphs of *Cretereisma* are embedded in a lateral position. Furthermore, these nymphs clearly have a laterally compressed body, which seems to be a unique autapomorphy, because such a habitus is unknown from other fossil and Recent mayfly nymphs. Bechly *et al.* (2001a: 49) therefore suggested that these peculiar larvae had a very different lifestyle than all other known mayflies.

Comparisons

The general wing structure of *Cretereisma* resembles closely that of *Prottereisma*. A main difference is the pairwise alignment of some of the longitudinal veins which is an autapomorphy of *Cretereisma*. Another difference lies in the structure of the first anal vein, which is simple in the forewing of *Prottereisma* and *Misthodotes* while it has a triad in the hind wing (Carpenter, 1933). In *Cretereisma*, however, A₁ is apparently either simple in both wings (*C. antiqua*) or has a triad at least in the forewing (*C. schwickerstorii*; structure of hind wing analis unknown). The general tendency towards a reduction of anal branches is probably correlated to the reduction of wing width. Nymphs are different from those of *Prottereisma* (and *Phthartus*) in their unique body shape and their plate-like gill appendages, and both *Prottereisma* and *Phthartus* have nine abdominal gill pairs while *Cretereisma* is more derived in having seven. Epeorusmimidae have seven pairs of plate-like gill appendages as well; however, this is the only remarkable similarity worth mentioning. In other structures, epeorusmimid nymphs differ fundamentally from *Cretereisma*, for example in body shape, a small prothorax and posteriorly directed wing pads, of which the hind pair is only small.

Phylogenetic position

Cretereisma shares with the Permian *Prottereisma* and *Misthodotes* a reduced anal area in the hind wing and the possession of a costal brace which is only slightly curved and does not reach the costal margin. These characters were not present (or have not been observed) in earlier ephemeroids. *Cretereisma* is more derived than *Prottereisma* in having only seven gill appendages in the nymph. On the other hand, the Upper Triassic *Litophlebia*, which resembles *Prottereisma* in many aspects, possibly shares at least one synapomorphy (shortened anal veins) with the

Triangulifera but not with *Cretereisma*. Thus it appears that *Cretereisma* is the sister taxon of *Litophlebia*+*Triangulifera*. *Prottereisma* and *Misthodotes* were abundant in the Lower Permian while *Litophlebia* lived about 60 myr later. Hence *Cretereisma*, with a phylogenetic position between *Litophlebia* and the two Permian taxa, is the only known representative of an ephemeroid lineage which had retained many plesiomorphic features for about 110–160 myr.

11.4 Ephemeroptera: mayflies

Arnold H. Staniczek

Mayflies represent one of the basal branches of winged insects and have a world wide distribution. Stem group representatives of mayflies date back to the Carboniferous. More than 3,000 Recent species have been described.

The larvae of mayflies are obligatorily bound to freshwater habitats. The imagines of mayflies are short-lived insects with a life span that varies from a few hours to a few days. Mayflies are the only pterygote insects to retain more than one fully winged stage. The subimago is the alate penultimate stage and undergoes another moulting to become the imago. The subimaginal moulting is generally regarded as a vestigial adult moulting as it is present in the apterygote insect groups. The adult stages of mayflies are easily recognized by the presence of usually very long cerci, often accompanied by a median long terminal filament (paracercus). At rest the wings are folded vertically over the abdomen. The hind wings are considerably smaller than the forewings and can also be entirely lost in some taxa. The imaginal wings are generally translucent and glabrous, but the subimaginal wings usually have a greyish or milky tinge and are equipped with setae. The venation is characterized by the presence of multiple crossveins, a pronounced costal brace especially in the forewing, and the presence of vein MP. The wings generally retain a pronounced pleating and corrugation of the longitudinal veins. The adult stages of mayflies do not feed, and their mouth parts are atrophied. The antennae are very short and bristle-like, convergent on Odonata. Other features are connected with the aerial mating flight that the males perform above the water: the male eyes are generally larger than the female ones, in some taxa (Baetidae, Leptophlebiidae) they are extended to so-called turbinate eyes, which are also morphologically divergent. The male forelegs are elongated to grasp the females around the base of the forewing during copulation. The male claspers, modified abdominal leglets of abdominal segment VIII, additionally clutch the female abdomen during copulation. Males have paired penes, and females have paired gonopores, respectively. The female ovipositor is completely reduced, and only some taxa have developed a secondary egg guide. After mating the female mayfly simply drops its eggs into the water. The eggs are equipped with highly variable adhesive structures to prevent drifting.

belong to the same genus. A careful revision by a specialist on fossil Trichoptera would be very useful. Furthermore, there are several putative new genera and species to be described, such as specimens nos SMNS 66282 (Figure 11.87d), SMNS 66287 (Figure 11.88a) and SMNS 66568 (Figure 11.88b), which are clearly distinct from any described species.

Crato Lepidoptera: moths and butterflies

Lepidoptera are extremely rare in the Crato Formation and only members of the microlepidopteran grade occur. Martins-Neto and Vulcano (1989b) and Martins-Neto (1999, 2001b) described the following five taxa.

- *Parasabatinca caldasae*: Micropterygidae; body length 3.5–5 mm; forewing length 3–3.5 mm; R_2 and R_3 ending near apex. Besides the holotype, a single specimen (SMNS 66279) has been identified as this species (Figure 11.87c).
- *Undopterix caririensis*: Undopterygidae; hind wing length 3.1 mm; M_1+M_2 fork slightly posterior to R_4+R_5 fork; CuA ending at anal margin near apex.
- *Gracilepterix pulchra*: in familia incertae sedis; similar to *Undopterix*; body length and forewing length about 3 mm; hind wing length 2.3 mm; R_1 forking anteriorly of R_2+R_3 and R_4+R_5 forks in forewings; Sc and R_1 unbranched in hind wings. One specimen of this species (SMNS 66277) has been identified (Figure 11.87d). This fossil butterfly was incorrectly listed as *Gracilepteryx* (*sic*) by Martins-Neto (2005b).
- *Xena nana* Martins-Neto, 1999b: Eolepidopterigidae. This taxon seems to be similar to *Eolepidopterix* and is said to be described by ‘Martins-Neto, 2000’ according to Martins-Neto (2005b), which seems to be a *lapsus* and should be ‘Martins-Neto, 1999b’.
- *Psamateia calipsa* Martins Neto, 2002: the present author could not locate the publication with this description, which is mentioned by Martins-Neto (2005b).

All the above species belong to the most primitive and most basal grade of Lepidoptera, and thus not within the Glossata, although Martins-Neto (2005b: 480) attributed all five species to the Eolepidopterigidae, but without explication. However, Grimaldi and Engel (2005: 562, figure 13.16) figured an unnamed primitive moth (AMNH SF 46441) with a piercing oviscapt similar to the Recent families Eriocraniidae and Acanthopteroctetidae that are indeed basal Glossata. This fossil could be the most derived Lower Cretaceous lepidopteran yet discovered, if it should not turn out to be just a small trichopteran like *Cratorella media*.

References

- Achtelik, M. 1967. Ueber die Anatomie des Kopfes von *Raphidia flavipes* Stein und die Verwandtschaftsbeziehungen der Raphidiidae zu den Megaloptera. *Zoologische Jahrbücher, Abteilung für Anatomie und Ontogenie der Tiere* **84**: 249–312.

- Afzelius, B. A. and Dallai, R. 1988. Spermatozoa of Megaloptera and Raphidioptera (Insecta, Neuropteroidea). *Journal of Ultrastructural and Molecular Structure Research* **101**: 185–191.
- Alexander, B. A. 1992. An exploratory analysis of cladistic relationships within the superfamily Apoidea, with special reference to sphecid wasps (Hymenoptera). *Journal of Hymenoptera Research* **1**: 25–61.
- Alexeev, A. V. 1994. Jurassic and Lower Cretaceous Buprestidae (Coleoptera) from Eurasia. *Paleontological Journal* **27**(I A): 9–34.
- 1996. New taxa of metallic wood-boring beetles (Coleoptera, Buprestidae) from the Mesozoic of Russia, Kazakhstan, and Mongolia. *Paleontological Journal* **30**: 310–117.
- 2000. On Mesozoic Buprestids (Coleoptera: Buprestidae) from Russia, Kazakhstan, and Mongolia. *Paleontological Journal* **34** (suppl. 3): 323–326.
- Andersen, N. M. 1998. Water striders from the Paleogene of Denmark with a review of the fossil record and evolution of semiaquatic bugs (Hemiptera: Gerromorpha). *Det Kongelige Danske Videnskabernes Selskab, Biologiske Skrifter* **50**: 1–152.
- and Weir, T. A. 2004. Australian water bugs. Their biology and identification (Hemiptera-Heteroptera, Gerromorpha and Nepomorpha). *Entomonograph* **14**: 1–344.
- Andersen, S. 2001. Silky lacewings (Neuroptera: Psychopsidae) from the Eocene–Palaeocene transition of Denmark with a review of the fossil record and comments on phylogeny and zoogeography. *Insect Systematics and Evolution* **32**: 419–438.
- Anderson, R. S. 2005. New Oxycoryninae from Central and South America: phylogenetic and biogeographical implications (Coleoptera: Belidae). *Systematic Entomology* **30**: 644–652.
- Ansorge, J. and Mostovski, M. B. 2000. Redescription of *Prohirmoneura jurassica* Handlirsch 1906 (Diptera: Nemestrinidae) from the Lower Tithonian lithographic limestone of Eichstätt (Bavaria). *Neues Jahrbuch für Geologie und Paläontologie Monatshefte* **2000**: 235–243.
- Anufriev, G. A. and Emeljanov, A. F. 1988. Tsikadovye (Auchenorrhyncha, Cicadinea). [Suborder Cicadinea (Auchenorrhyncha)], pp. 12–495. In Lehr, A. P. (ed), *Oprelelitel' nasekomykh Dal'nego Vostoka SSSR*. St Petersburg: Nauka.
- Arnol'di, L. V., Zherikhin, V. V., Nikritin, L. M. and Ponomarenko, A. G. 1992. *Mesozoic Coleoptera*. Washington DC: Smithsonian Institution Libraries and The National Science Foundation.
- Asche, M. 1988. Preliminary thoughts on the phylogeny of Fulgoromorpha (Homoptera, Auchenorrhyncha). *Proceedings of the 6th Auchenorrhyncha Meeting*, Turin, Italy, 7–11 September 1987, pp. 47–53.
- Aspöck, U. and Aspöck, H. 2004. Two significant new snakeflies from Baltic amber, with discussion on autapomorphies of the order and its included taxa (Raphidioptera). *Systematic Entomology* **29**: 11–19.
- Azar, D. and Nel, A. 2002. New Cretaceous psychodid flies from Lebanese amber and Santana Formation (Chapada do Araripe, Brazil) (Diptera). *Annales de la Societe Entomologique de France* (n.s.) **38**: 253–262.
- Baudoin, R. 1980. Sur les Gerris des miroirs d'eau actuels et les Chresmoda des lagunes post-récifales portlandiennes de Solnhofen. *Annales des Sciences Naturelles / Zoologie* **2**: 111–116.
- Bechly, G. 1996. Review: SANTANA 1.0 (1996): CD-ROM by digital-fossil. *Petalura* **2**: <http://www.bernstein.naturkundemuseum-bw.de/odonata/pet2p03.htm>.

- 1997a. Die Libellen (und einige andere fossile Insekten) aus der Unterkreide von Brasilien (Santana). 4. *Fachgespräch "Fossile Insekten"*, Clausthal-Zellerfeld, 28–29 June 1997, Abstracts.
 - 1997b. Dragonflies from the Lower Cretaceous of Brazil. *Meganeura* **1**: 27–28.
 - 1997c. Dragonflies from the Lower Cretaceous of Brazil. *Inclusion Wrostek* **27**: 9.
 - 1998a. Santana – Schatzkammer fossiler Insekten. *Fossilien* **2/98**: 95–99.
 - 1998b. Santana – Forschungsgeschichte und Fauna. *Fossilien* **3/98**: 148–156.
 - 1998c. New fossil dragonflies from the Lower Cretaceous Crato Formation of north-east Brazil (Insecta: Odonata). *Stuttgarter Beiträge zur Naturkunde, Serie B* **264**: 1–66.
 - 1998d. [Santana – Eldorado of Mesozoic fossils]. *Catalogue of the Taipei Natural History Show* **1998**: 148–156.
 - 1999a *Phylogeny and Systematics of Fossil Dragonflies (Insecta: Odonatoptera) with Special Reference to some Mesozoic Outcrops*. Published PhD thesis, Eberhard-Karls-Universität Tübingen.
 - 1999b. Notices. *Meganeura* **4**: 9.
 - 2000. Two new fossil dragonfly species (Insecta: Odonata: Anisoptera: Araripegomphidae and Lindeniiidae) from the Crato Limestone (Lower Cretaceous, Brazil). *Stuttgarter Beiträge zur Naturkunde, Serie B* **296**: 1–16.
 - 2002. Phylogenetic systematics of Odonata. In Schorr, M. and Lindeboom, M. (eds), *Fundamentals of Odonatological Research*. ODOLITonCD 1.0 (ISSN 1438–034x). www.dragonflyresearch.de/.
 - 2003. The phylogenetic relationships of the three extant suborders of Odonata. *Entomologische Abhandlungen* **61**: 127–128.
 - 2005. A re-description of "Stenophlebia" casta (Insecta: Odonata: Parastenophlebiidae n. fam.) from the Upper Jurassic Solnhofen Limestone in Germany. *Stuttgarter Beiträge zur Naturkunde, Serie B* **359**: 1–12.
 - and Ueda, K. 2002. The first fossil record and first New World record for the dragonfly clade Chlorogomphida (Insecta: Odonata: Anisoptera: Araripechlorogomphidae n. fam.) from the Crato Limestone (Lower Cretaceous, Brazil). *Stuttgarter Beiträge zur Naturkunde, Serie B* **328**: 1–11.
 - , Nel, A. Martínez-Delclòs, X. and Fleck, G. 1998. Four new dragonfly species from the Upper Jurassic of Germany and the Lower Cretaceous of Mongolia (Anisoptera: Hemeroscopidae, Sonidae, and Proterogomphidae fam. nov.). *Odonatologica* **27**: 149–187.
 - , Haas, F., Schwaller, W., Schmalfuss, H. and Schmid, U. 2001a. Ur-Geziefer – Die faszinierende Evolution der Insekten. *Stuttgarter Beiträge zur Naturkunde, Serie C* **49**: 1–96.
 - , Brauckmann, C., Zessin, W. and Gröning, E. 2001b. New results concerning the morphology of the most ancient dragonflies (Insecta: Odonatoptera) from the Namurian of Hagen-Vorhalle (Germany). *Journal of Zoological Systematics and Evolutionary Research* **39**: 209–226.
 - , Nel, A. Martínez-Delclòs, X. et al. 2001c. A revision and phylogenetic study of Mesozoic Aeshnoptera, with description of several new families, genera and species (Insecta: Odonata: Anisoptera). *Neue paläontologische Abhandlungen* **4**: 1–219.
 - , Dietl, G. and Schweigert, G. 2003. A new species of *Stenophlebia* (Insecta: Odonata: Stenophlebiidae) from the Nusplingen Lithographic Limestone (Upper Jurassic, SW Germany). *Stuttgarter Beiträge zur Naturkunde, Serie B* **338**: 1–10.
- Becker-Migdisova, E. E. 1946. Ocherki po sravnitel'noi morfologii sovremennyykh i permskikh Homoptera. Chast' I. *Izvestia AN SSSR, Biology Series* **6**: 741–766.

- 1947. *Cicadoprosbole sogutensis* gen. et sp. nov. – perekhodnaya forma mezhdu permeskimi Prosbolidae i sovremennymi Cicadidae. *Doklady Akademii Nauk SSSR* 5: 445–448.
- 1949. Mezozojskie Homoptera Srednei Azii. *Trudy Palaeontologicheskogo Instituta Nauk* 22: 1–68.
- 1962a. Nekotorye novye poluzhestokrylye i senoedy. *Palontologicheskiy Zhurnal* 1: 89–104.
- 1962b. Otryad Homoptera. Ravnokrylye, pp. 162–208. In Rohdendorf, B. B. (ed.), *Osnovy paleontologii. Chlenistonogie trakheinye i khelitserovye*. Moscow: Academy of Sciences USSR.
- and Popov, Yu., A. 1963. Some representatives of Heteroptera from the Jurassic of Karatau. *Palaeontological Journal of the Academy of Science USSR* 2: 74–82.
- Bedford, G. O. 1978. The biology and ecology of the Phasmatodea. *Annual Review of Entomology* 23: 125–149.
- Beier, M. 1968. Mantodea (Fangheuschrecken). *Handbuch der Zoologie*, vol. IV, Band Arthropoda, 2 Hälfte: Insecta. Zweite Auflage: 1–47.
- Belayeva, N. V. 2002. Order Termitida, pp. 270–273. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- Bennett, D. J. and Engel, M. S. 2005. A primitive sapygid wasp in Burmese amber (Hymenoptera: Sapygidae). *Acta Zoologica Cracoviensia* 48B(1–2): 1–9.
- Béthoux, O. and Nel, A. 2002. Venation pattern and revision of Orthoptera sensu nov. and sister groups. Phylogeny of Palaeozoic and Mesozoic Orthoptera sensu nov. *Zootaxa* 96: 1–88.
- Nel, A., Lapeyrie, J., Gand, G. and Galtier, J. 2002. *Raphogla rubra* gen. n., sp. n., the oldest representative of the clade of modern Ensifera (Orthoptera: Tettigoniidea, Gryllidea). *European Journal of Entomology* 99: 111–116.
- Beutel, R. G. 1995. The Adephaga (Coleoptera): phylogeny and evolutionary history, pp. 173–217. In Pakaluk, J. and Ślipiński, S. A. (eds), *Biology, Phylogeny, and Classification of Coleoptera*, vol. 1. Warszawa: Muzeum I Instytut Zoologii PAN.
- Bischoff, H. 1927. *Biologie der Hymenopteren, Eine Naturgeschichte der Hautflügler*. Biologische Studienbücher. Berlin: J. Springer.
- Bitsch, C. and Bitsch, J. 2000. The phylogenetic interrelationships of the higher taxa of apterygote insects. *Zoologica Scripta* 29: 131–156.
- Blagoderov, V., Grimaldi, D. A. and Fraser, N. C. 2007. How time flies for flies: diverse Diptera from the Triassic of Virginia and early radiation of the order. *American Museum Novitates* (in press).
- Bode, A. 1953. Die Insektenfauna der Ostniedersächsischen oberen Liás. *Palaeoentomographica A* 103: 1–406.
- Bohart, R. M. and Menke, M. S. 1976. *Sphecid wasps of the world. A generic revision*. Los Angeles: University of California Press, Berkeley.
- Bohn, H. 2003a. Ordnung Blattoptera, Schaben, pp. 197–223. In Dathe, H. H. (ed), *Lehrbuch der Speziellen Zoologie, Band 1, Teil 5 (Insecta)*. Heidelberg: Spektrum.
- 2003b. Ordnung Isoptera, Termiten, pp. 223–250. In Dathe, H. H. (ed.), *Lehrbuch der Speziellen Zoologie, Band 1, Teil 5 (Insecta)*. Heidelberg: Spektrum.
- Bordy, E. M., Bumby, A. J., Catuneanu, O. and Eriksson, P. G. 2004. Advanced early Jurassic termite (Insecta: Isoptera) nests: evidence from the Clarens Formation in the Tuli Basin, southern Africa. *Palaios* 19: 68–78.
- Börner, C. 1909. Neue Homologien zwischen Crustaceen und Hexapoden. Die Beißmandibel der Insekten und ihre phylogenetische Bedeutung. Archi- und Metapterygota. *Zoologischer Anzeiger* 34: 100–125.

- Boudreaux, H. B. 1979. *Arthropod Phylogeny with Special Reference to Insects*. New York: Wiley.
- Boukary, I. B., Tourneur, J. C. and Gingras, J. 1996. Life cycle of *Forficula senegalensis* Serv. (Dermoptera: Forficulidae) and its relationship to the development of bulrush millet in Sundanese-Sahelian zone of Niger. *Canadian Entomologist* **128**: 831–838.
- Boulard, M. and Nel, A. 1990. Sur deux cigales fossiles des terrains tertiaires de la France [Homoptera, Cicadoidea]. *Revue Française d'Entomologie (n.s.)* **12**: 37–45.
- Bourgoin, Th. 1993. Cladistic analysis of the Meenopliidae–Kinnaridae genera: the Kinnaridae, a paraphyletic family (Hemiptera, Fulgoromorpha). In Drosopoulos, S., Petrakis, P. V., Claridge, M. F. and de Vrijer, P. W. F. (eds), *Proceedings of the 8th Auchenorrhyncha Congress*, Delphi, Greece, 9–13 August 1993: 22–24.
- and Campbell, B. C. 2002. Inferring a phylogeny for Hemiptera: falling into the ‘autapomorphic trap’, pp. 67–82. In Holzinger, W. (ed.), *Zikaden – Leafhoppers, Planthoppers and Cicadas (Insecta: Hemiptera: Auchenorrhyncha)*. Denisia **04**, Zugleich Kataloge, Neue Folge Nr. 176, Linz: OÖ. Landesmuseums.
- Steffen-Campbell, J. D. and Campbell, B. C. 1997. Molecular phylogeny of Fulgoromorpha (Insecta, Hemiptera, Archaeorrhyncha). The enigmatic Tettigometridae: evolutionary affiliations and historical biogeography. *Cladistics* **17**: 207–224.
- Bradley, C. 1955. Notes on the synonymy, distribution and affinities of the subfamily Fedtschenkiinae of the Sapygidae (Hymenoptera). *Entomological News* **66**: 230–233.
- Brandao, C. R. F., Martins-Neto, R. G. and Vulcano, M. A. 1989. The earliest known fossil ant (First Southern Hemisphere mesozoic record) (Hymenoptera: Formicidae: Myrmeciinae). *Psyche* **96**: 195–208.
- Brito, I. M. 1984. Nota preliminar sobre os insetos da Formação Santana, Cretáceo inferior da Chapada do Araripe. *XXXIII Congresso Brasileiro de Geologia, Rio de Janeiro*: 530–535.
- 1987. Nota preliminar sobre uma nova efêmera do Cretáceo do Ceará (Insecta Ephemeroptera). *Anais do X Congresso Brasileiro de Paleontologia* **2**: 593–597.
- Brogniart, C. 1893. *Recherches pour servir à l'Histoire des Insectes Fossiles des Temps Primaires, Précédées d'une Étude sur la Nervation des Ailes des Insects*. Thèse présentée à la Faculté des Sciences de Paris, no. 821; Paris.
- Brothers, D. J. 1975. Phylogeny and classification of the aculeate Hymenoptera, with special reference to Mutillidae. *University of Kansas Science Bulletin* **50**: 483–648.
- Burckhardt, D. and Agosti, D. 1991. New records of South American Peloridiidae (Homoptera: Coleorrhyncha). *Revista Chilena Entomología* **19**: 71–75.
- and Cekalovic, T. K. 2002. An anomalous moss-bug from Southern Chile and notes on *Pantinia darwini* (Hemiptera, Coleorrhyncha, Peloridiidae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft / Bulletin de la Société entomologique Suisse* **75**: 57–59.
- Burnham, L. 1978. Survey of Social Insects in the Fossil Record. *Psyche* **85**: 85–133.
- Carle, F. L. 1982. The wing vein homologies and phylogeny of the Odonata: a continuing debate. *S.I.O. Rapid Communications* **4**: 1–66.
- 1995. Evolution, taxonomy, and biogeography of ancient Gondwanian Libelluloids, with comments on anisopterid evolution and phylogenetic systematics (Anisoptera: Libelluloidea). *Odonatologica* **24**: 383–424.
- and Wighton, D. C. 1990. Odonata, pp. 51–68. In Grimaldi, D. A. (ed.), *Insects from the Santana Formation, Lower Cretaceous, of Brazil*. *Bulletin of the American Museum of Natural History*, vol. 195.

- Carpenter, F. M. 1933. The Lower Permian Insects of Kansas. Part 6. *Proceedings of the American Academy of Arts and Sciences* 68: 411–503.
- 1979. The Lower Permian insects from Oklahoma. Part 2. Orders Ephemeroptera and Palaeodictyoptera. *Psyche* 86: 261–290.
- 1992. Superclass Hexapoda. In Kaesler, R. L. (ed), *Treatise on Invertebrate Paleontology, R, Arthropoda 4, 3/4*. Boulder and Lawrence: Geological Society of America and University of Kansas.
- Carpenter, J. M. 1982. The phylogenetic relationships and natural classification of the Vespoidea (Hymenoptera). *Systematic Entomology* 7: 11–38.
- 1987. Phylogenetic relationships and classification of the Vespinae (Hymenoptera: Vespidae). *Systematic Entomology* 12: 423–431.
- and Rasnitsyn, A. P. 1990. Mesozoic Vespidae. *Psyche* 97(1–2): 1–20.
- Carver, M., Gross, G. F. and Woodward, T. E. 1992. Hemiptera, pp. 429–509. In Naumann, I. D. (ed), *The Insects of Australia*. Melbourne: CSIRO and Melbourne University Press.
- Cassola, F. and Werner, K. 2004. A fossil tiger beetle specimen from the Brazilian Mesozoic: *Oxycheilopsis cretacicus* n. gen., n. sp. *Mitteilungen der Münchener entomologischen Gesellschaft* 94: 75–81.
- Chadwick, M. A., Hunter, H., Feminella, J. W. and Henry, R. P. 2002. Salt and water balance in *Hexagenia limbata* (Ephemeroptera: Ephemeridae) when exposed to brackish water. *Florida Entomologist* 85: 650–651.
- Chen, S. and Tian, C.-c. 1973. A new family of Coleoptera from the Lower Cretaceous of Kansu. *Acta Entomologica Sinica* 16: 169–179.
- Chin, K. and Gill, B. D. 1996. Dinosaurs, dung beetles, and conifers: participants in a Cretaceous food web. *Palaios* 11: 280–285.
- Cockerell, T. D. A. 1913. The first fossil Mydaid fly. *The Entomologist* 46: 207–208.
- 1917. Insects in Burmese amber. *Annals of the Entomological Society of America* 10: 323–329.
- Colgan, D. J., Cassis, G. and Beacham, E. 2003. Setting the molecular phylogenetic framework for the Dermaptera. *Insect Systematics and Evolution* 34: 60–80.
- Compton, S. G. and Ware, A. B. 1991. Ants disperse the elaiosome-bearing eggs of an African stick insect. *Psyche* 98: 207–213.
- Crampton, G. C. 1938. The interrelationships and lines of descent of living insects. *Psyche* 45: 165–181.
- Crowson, R. A. 1981. *The Biology of Coleoptera*. New York: Academic Press.
- Cryan, J. R. 2005. Molecular phylogeny of Cicadomorpha (Insecta: Hemiptera: Cicadoidea, Cercopoidea and Membracoidea): adding evidence to the controversy. *Systematic Entomology* 30: 563–574.
- Currie, D. C. and Grimaldi, D. 2000. A new blackfly (Diptera: Simuliidae) genus from mid-Cretaceous (Turonian) amber of New Jersey, pp. 473–485. In Grimaldi, D. (ed), *Studies on Fossils in Amber, with Particular Reference to the Cretaceous of New Jersey*. Leiden: Backhuys.
- da Costa-Lima, A. 1950. Ninfas de Efemerídeo fossil do Ceará. *Anais Academia, Brasileira, Ciências* 22: 419–420.
- Darling, D. C. and Sharkey, M. J. 1990. Order Hymenoptera, pp. 123–153. In Grimaldi, D. A. (ed.), *Insects from the Santana Formation, Lower Cretaceous, of Brazil. Bulletin of the American Museum of Natural History* 195.
- Day, M. C., Else, G. R. and Morgan, D. 1981. The most primitive Scoliidae (Hymenoptera). *Journal of Natural History* 15: 671–684.

- Deitz, L. L. and Dietrich, C. H. 1993. Superfamily Membracoidea (Homoptera: Auchenorrhyncha). I. Introduction and revised classification with new family-group taxa. *Systematic Entomology* **18**: 287–296.
- Demoulin, G. 1955. Sur une larve siphlonuridienne d'Ephémère fossile du Brésil. *Bulletin et Annales de la Société Royale d'Entomologie de Belgique* **91**: 270–271.
- 1971. Contribution à l'étude morphologique, systématique et phylogénique des Ephemeroptères Jurassiques (1). VI. L'aile postérieure des Hexagenites Scudder et les rapports Hexagenitidae-Chromarcyidae-Oligoneuriidae. *Bulletin* **47(29)**: 1–10.
- Desutter-Grandcolas, L. 2003. Phylogeny and the evolution of acoustic communication in extant Ensifera (Insecta, Orthoptera). *Zoologica Scripta* **32**: 252–261.
- Dietrich, C. H. 1999. The role of grasslands in the diversification of leafhoppers (Homoptera: Cicadellidae): a phylogenetic perspective, pp. 44–48. In Warwick, C. (ed.), *Proceedings of the Fifteenth North American Prairie Conference*. Bend, OR: Natural Areas Association.
- 2002. Evolution of Cicadomorpha (Insecta, Hemiptera), pp. 155–170. In Holzinger, W. (ed.), *Zikaden – Leafhoppers, Planthoppers and Cicadas (Insecta: Hemiptera: Auchenorrhyncha)*, Denisia, 4, Neue Folge Nr. 176, zugleich Kataloge des OÖ. Linz: Landesmuseums.
- 2005. Keys to the families of Cicadomorpha and subfamilies and tribes of Cicadellidae (Hemiptera: Auchenorrhyncha). *Florida Entomologist* **88**: 502–517.
- and Rakitov, R. A. 2002. Some remarkable new deltocephaline leafhoppers (Hemiptera: Cicadellidae: Deltocephalinae) from the Amazonian rainforest canopy. *Journal of the New York Entomological Society* **110**: 1–48.
- McKamey, S. H. and Dietz, L. L. 2001. Morphology-based phylogeny of the treehopper family Membracidae (Hemiptera: Cicadomorpha: Membracoidea). *Systematic Entomology* **26**: 213–239.
- Deitz, L. L. and Dietrich, C. H. 1993. Superfamily Membracoidea (Homoptera: Auchenorrhyncha). I. Introduction and revised classification with new family-group taxa. *Systematic Entomology* **18**: 287–296.
- Distant, W. L. 1883. Contribution to a proposed monograph of the homopterous family Cicadidae. *Proceedings of the Zoological Society of London* **1883**: 187–194.
- 1905. Rhynchotal notes XXXIII. *Annals and Magazine of Natural History* **7**: 1622–1635.
- Dolin, V. G. and Nel, A. 2002. Trois nouveaux Elateridae fossiles du Mésozoïque supérieur de Chine (Coleoptera). *Bulletin de la Société entomologique de France* **107**: 341–346.
- Ehrmann, R. 2002. *Mantodea. Gottesanbeterinnen der Welt*. Munich: Natur und Tier Verlag GmbH.
- Eichwald, E. 'd. 1864. Sur un terrain jurassique a poissons et Insectes d'eau douce de la Sibérie orientale. *Bulletin de Societe Geologique de France* **21**: 19–25.
- Emeljanov, A. F. 1991. K voprosu ob obeme i podrazdeleniyakh sem. Achlidiae (Homoptera, Cicadina). *Entomologicheskoe Obozrenie* **70**: 373–393. [Published in English as: Emel'yanov, A. F. 1992. Toward the problem of the limits and subdivisions of Achlidiae (Homoptera, Cicadina). *Entomological Review* **71**: 53–73].
- 1992. Opisanie trib podsem. Achilinae (Homoptera, Achilidae) i utochnenie ikh sostava. *Entomologicheskoe Obozrenie* **71**: 574–594. [Published in English as: Emel'yanov, A. F. 1993. Description of tribes of the subfamily Achilinae (Homoptera, Achilidae) and revision of their composition. *Entomological Review* **72**: 7–27].

- 1994 Pervaya iskopayemaya nakhodka semeistva Derbidae i pereopisanie paleogenovogo roda *Hooleya* Cockerell (Achilidae) (Insecta: Homoptera, Fulgoroidea). *Paleontologicheskii Zhurnal* **3**: 76–82. [Published in English as: Emeljanov, A. F. 1995. The first find of fossil Derbidae, and a redescription of Paleogene achilid *Hooleya* Cockerell (Insecta: Homoptera, Fulgoroidea). *Paleontological Journal* **28**: 92–101].
- 1999. Notes on the delimitation of families of the Issidae group with description of a new species of Caliscellidae belonging to a new genus and tribe (Homoptera, Fulgoroidea). *Zoosystematica Rossica* **8**: 61–72.
- 2002. Contribution to classification and phylogeny of the family Cixiidae (Hemiptera, Fulgoromorpha), pp. 103–112. In Holzinger, W. (ed), *Zikaden – Leafhoppers, Planthoppers and Cicadas (Insecta: Hemiptera: Auchenorrhyncha)* Denisia, 04, Zugleich Kataloge, Neue Folge Nr. 176, Linz: OÖ. Landesmuseums.
- Engel, M. S. 2001. A monograph of the Baltic amber bees and evolution of the Apoidea (Hymenoptera). *Bulletin of the American Museum of Natural History* **259**: 1–192.
- 2002. The smallest snakefly (Raphidioptera: Mesoraphidiidae): a new species in Cretaceous amber from Myanmar, with a catalogue of fossil snakelies. *American Museum Novitates* **3363**: 1–22.
- and Grimaldi, D. A. 2004. New light shed on the oldest insect. *Nature* **427**: 627–630.
- and Chatzimanolis, S. 2005. Early Cretaceous earwigs (Dermaptera) from the Santana Formation, Brazil. *Polskie Pismo Entomologiczne* **74**: 219–226.
- Esaki, T. 1949. The occurrence of the Mesozoic insect *Chresmoda* in the Far East. *Insecta Matsumarana* **17**: 4–5.
- Eskov, K. Yu. 1984. Dreif materikov i problema isoricheskoi biogeografi, pp. 24–92. In: *Faunogenet i filogenet*. Moscow: Nauka.
- and Golovatch, S. I. 1986. On the origin of Trans-Pacific disjunctions. *Zoologisches Jahrbuch für Systematik* **113**: 265–285.
- Evans, J. W. 1956. Palaeozoic and Mesozoic Hemiptera. *Australian Journal of Zoology* **4**: 165–258.
- 1981. A review of the present knowledge of the family Peloridiidae and new genera and new species from New Zealand and New Caledonia (Hemiptera: Insecta). *Records of the Australian Museum* **34**: 381–406.
- Evenhuis, N. L. 1994. *Catalogue of the Fossil Flies of the World (Insecta: Diptera)*. Leiden: Backhuys.
- Fanenbruck, M., Harzsch, S. and Wägele, J.-W. 2004. The brain of the Remipedia (Crustacea) and an alternative hypothesis on their phylogenetic relationships. *Proceedings of the National Academy of Sciences USA* **101**: 3868–3873.
- Farrell, B. D. 1998. “Inordinate Fondness” explained: why are there so many beetles? *Science* **281**: 555–557.
- Fennah, R. G. 1954. The higher classification of the family Issidae (Hemiptera, Fulgoroidea), with description of new species. *Transactions of the Royal Entomological Society* **105**: 455–474.
- 1961. The occurrence of a Cixiinae Fulgoroid (Homoptera) in a Weald Clay. *Annals and Magazine of Natural History* **13**: 161–163.
- 1987. A new genus and species of Cixiidae (Homoptera, Fulgoroidea) from Lower Cretaceous amber. *Journal of Natural History* **21**: 1237–1240.
- Fleck, G., Bechly, G., Martínez-Delclòs, X., Jarzembski, E. A., Coram, R. and Nel, A. 2003. Phylogeny and classification of the Stenophlebioptera. (Insecta, Odonata, Epiproctophora). *Annales de la Société entomologique de France (n.s.)* **39**: 55–93.

- Nel, A., Bechly, G. and Escuillié, F. 2002. The larvae of the Mesozoic family Aeschnidiidae and their phylogenetic implications (Insecta, Odonata, Anisoptera). *Palaeontology* **45**: 165–184.
- Flook, P. K. and Rowell, C. H. F. 1997. The effectiveness of mitochondrial rRNA gene sequences for the reconstruction of the phylogeny of an insect order (Orthoptera). *Molecular Phylogenetics and Evolution* **8**: 177–192.
- and Rowell, C. H. F. 1998. Inferences about orthopteroid phylogeny and molecular evolution from small subunit nuclear ribosomal DNA sequences. *Insect Molecular Biology* **7**: 163–178.
- , Klee, S. and Rowell, C. H. F. 1999. Combined molecular phylogenetic analysis of the Orthoptera (Arthropoda, Insecta) and implications for their higher systematics. *Systematic Biology* **48**: 233–253.
- , Klee, S. and Rowell, C. H. F. 2000. Molecular phylogenetic analysis of the Pneumoroidea (Orthoptera, Caelifera): molecular data resolve morphological character conflicts in the basal Acridomorpha. *Molecular Phylogenetics and Evolution* **15**: 345–354.
- Fontes, L. R. and Vulcano, M. A. 1998. Cupins fósseis do Novo Mundo, pp. 243–295. In Fontes, L. R. and Filho, E. B. (eds), *Cupins – O desafio do conhecimento*. Piracicaba: Fundação de Estudos Agrários Luiz de Queiroz FEALQ.
- and Vulcano, M. A. 2004. Catalog of the Fossil Isoptera of the New World. *Sociobiology* **44**: 345–364.
- Frank, J. H. and Thomas, M. C. 2002. http://creatures.ifas.ufl.edu/misc/beetles/fl_rove_beetles.htm.
- Fraser, F. C. 1957. A reclassification of the order Odonata, based on some new interpretations of the venation of the dragonfly wing. *Handbook Royal Zoological Society of New South Wales* **12**: 1–133.
- Furth, D. G. 1994. Frank Morton Carpenter (1902–1994): academic biography and list of publications. *Psyche* **101**: 127–144.
- Gauld, I. and Bolton, B. 1988. *The Hymenoptera*. British Museum (Natural History). Oxford: Oxford University Press.
- Gebicki, C. and Wegierek, P. 1993. *Oligocixia electrina* gen. et sp. nov. (Homoptera, Auchenorrhyncha, Cixiidae) from Dominican amber. *Annales Naturhistorisches Museum, Wien* **95A**: 121–125.
- Germar, E. F. 1839. Die versteinerten Insekten Solenhofens. *Nova Acta Leopoldina Carolina Akademia* **19**: 187–222.
- Godoy, C. and Webb, M. D. 1994. Recognition of a new subfamily of Cicadellidae from Costa Rica based on a phenetic analysis with similar taxa (Hemiptera Homoptera Auchenorrhyncha). *Tropical Zoology* **7**: 131–144.
- Godunko, R. J. 2004. A new fossil subgenus and species of the genus *Ecdyonurus* Eaton, 1868 from Eocene Baltic amber (Ephemeroptera: Heptageniidae). *Genus* **15**: 323–328.
- and Neumann, C. 2006. Fossil mayfly collections of the Museum für Naturkunde, Humboldt University, Berlin. I. *Electroletus soldani* gen. et sp. nov. (Ephemeroptera: Amelitidae) from the Eocene Baltic Amber.- *Annales Zoologici* **56**: 437–441.
- Gomez-Pallerola, J. E. 1984. Nuevos Paleontíidos del yacimiento Infracretácico de la ‘Pedrera de Meiá’ (Lérida). *Boletín Geológico y Minero España* **95**: 301–309.
- 1986. Nuevos insectos fósiles de las calizas litográficas del Cretacico Inferior del Montsec (Lérida). *Boletín Geológico y Minero España* **97**: 27–46.

- Goodwyn, P. J. P. 2002. A new genus of water measurer from the Lower Cretaceous Crato Formation in Brazil (Insecta: Heteroptera: Gerromorpha: Hydrometridae). *Stuttgarter Beiträge Naturkunde, Serie B* **316**: 1–9.
- Gorochov, A. V. 1992. Fossil stick insects (Phasmatoptera) from the Jurassic and Cretaceous. *Trudy Paleontologicheskii Instituta Rossiosky Akademii, Nauk* 252: 112–116.
- 1994. Permian and Triassic walking sticks (Phasmatoptera) from Eurasia. *Paleontological Journal* **28**: 83–97.
- 1995. The system and evolution of the suborder Ensifera (Orthoptera). *Proceedings of the Zoological Institute, Russian Academy of Sciences* **260**(1): 1–223, **260**(2): 1–212.
- 2000. Phasmomimidae: are they Orthoptera or Phasmatoptera? *Paleontological Journal* **34**: 295–300.
- 2001. The most interesting finds of orthopteroid insects at the end of the 20th century and a new recent genus and species. *Journal of Orthoptera Research* **10**: 353–367.
- and Rasnitsyn, A. P. 2002. Superorder Gryllidea Laicharting, 1781 (= Orthopteroidea Handlirsch, 1903), pp. 293–303. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- Goulet, H. and Huber, J. T. 1993. *Hymenoptera of the World: an Identification Guide to Families*. Research Branch Agriculture Canada Publication 1894/EL.
- Gratshev, V. G. and Zherikhin, V. V. 1993. New fossil mantids (Insecta, Mantida). *Paleontological Journal* **27**: 148–165.
- and Zherikhin, V. V. 2000. New Early Cretaceous weevil taxa from Spain (Coleoptera, Curculionidae). *Acta Geologica Hispanica* **35**: 37–46.
- Grimaldi, D. 1990. Insects from the Santana Formation, Lower Cretaceous, of Brazil. *Bulletin of the American Museum of Natural History* **195**: 5–191.
- 1991. The Santana Formation insects, pp. 379–406. In Maisey, J. G. (ed), *Santana Fossils: an Illustrated Atlas*. Neptune City, NJ: T.F.H. Publications.
- 1996. *Amber – A window to the past*. AMNH; New York; 216 pp.
- 1999. The co-radiations of pollinating insects and angiosperms in the Cretaceous. *Annals of the Missouri Botanical Garden* **86**: 373–406.
- 2001. Insect evolutionary history from Handlirsch to Hennig, and beyond. *Journal of Paleontology* **75**: 1152–1160.
- 2003. A revision of Cretaceous mantises and their relationships, including new taxa (Insecta: Dictyoptera: Mantodea). *American Museum Novitates* **3412**: 1–47.
- and Maisey, J. G. 1990. Introduction, pp. 5–14. In Grimaldi, D. A. (ed), *Insects from the Santana Formation, Lower Cretaceous, of Brazil*. *Bulletin of the American Museum of Natural History*, **195**.
- and Cumming, J. 1999. Brachyceran Diptera in Cretaceous ambers and Mesozoic diversification of the Eremoneura. *Bulletin of the American Museum of Natural History* **239**: 1–124.
- and Engel, M. S. 2005. *Evolution of Insects*. Cambridge: Cambridge University Press.
- Engel, M. S. and Nascimbene, P. C. 2002. Fossiliferous Cretaceous amber from Myanmar (Burma): its rediscovery, biotic diversity, and paleontological significance. *American Museum Novitates* **3361**: 1–71.
- Guiglia, D. 1955. Appunti intorno al Gen. *Fedtschenkia* Saussure (Hymenoptera: Sapygidae). *Doriania Suppl. Agli* **II** (55): 1–4.
- 1969. The history of the peculiar Genus *Fedtschenkia* Saussure (1880) (Hymenoptera: Fedtschenkiidae). *Israel Journal of Entomology* **4**: 339–342.

- 1972. De la distribution géographique du genre *Fedtschenkia* (Saussure, 1880) (*Hymenoptera: Fedtschenkiidae*). *Proceedings of the XIII Congress of Entomology, Moscow* (1968) **1**: 138–140.
- Gullan, P. J. and Cranston, P. S. 2005. *The Insects – an Outline of Entomology*. Oxford: Blackwell Publishing.
- Gusenleitner, J. 1994. Das Vorkommen der Familie Sapygidae in Österreich. (Insecta: Hymenoptera: Sapygidae). *Annalen des Naturhistorischen Museums in Wien* **96B**: 173–188.
- 1996. Beitrag zur Kenntnis paläarktischer Sapyginae (Hymenoptera, Sapygidae). *Linzer biologische Beiträge* **28**: 23–38.
- 1997. Über Funde von Sapygidae aus dem östlichen Mittelmeergebiet (Sapygidae, Hymenoptera). *Linzer biologische Beiträge* **29**: 105–108.
- Haas, F. and Gorb, S. 2004. Evolution of locomotory attachment pads in the Dermaptera (Insecta). *Arthropod Structure and Development* **33**: 45–66.
- , Gorb, S. and Wootton, R. J. 2000. Elastic joints in dermapteran hind wings: Materials and wing folding. *Arthropod Structure and Development* **29**: 137–146.
- and Kukalová-Peck, J. 2001. Dermaptera hind wing structure and folding: new evidence for familial, ordinal and superordinal relationships within Neoptera (Insecta). *European Journal of Entomology* **98**: 445–509.
- and Klass, K.-D. 2003. The basal phylogenetic relationships in the Dermaptera, pp. 138–142. In Klass, K.-D. (ed), *Proceedings of the 1st Dresden Meeting on Insect Phylogeny: Phylogenetic Relationships within the Insect Orders*, Dresden, 19–21, September 2003. *Entomologische Abhandlungen* **61**.
- Haase, E. 1890. Beiträge zur Kenntnis der fossilen Arachniden. *Zeitschrift der deutschen geologischen Gesellschaft* **42**: 629–657.
- Hamilton, K. G. A. 1971. A remarkable fossil homopteran from Canadian Cretaceous amber representing a new family. *Canadian Entomologist* **103**: 943–946.
- 1972. The insect wing, Part III. Venation of the orders. *Journal of the Kansas Entomological Society* **45**: 145–162.
- 1983. Classification, morphology and phylogeny of the family Cicadellidae (Rhynchota: Homoptera), pp. 15–37. In Knight, W. J. (ed), *International Workshop on Leafhoppers and Planthoppers of Economic Importance*. London: Commonwealth Institute of Entomology.
- 1990. Homoptera, pp. 82–122. In Grimaldi, D. (ed), Insects from the Santana Formation, Lower Cretaceous, of Brazil. *Bulletin of the American Museum of Natural History* **195**.
- 1992. Lower Cretaceous Homoptera from the Koonwarra Fossil Bed in Australia with a New Superfamily and Synopsis of Mesozoic Homoptera. *Annals of the Entomological Society of America* **85**: 423–430.
- 1996. Cretaceous Homoptera from Brazil: Implications for Classification, pp. 89–110. In Schaefer, C. W. (ed), *Studies on Hemipteran Phylogeny*. Langham, MD: Thomas Say Publications in Entomology. Entomological Society of America.
- 1999. The ground-dwelling leafhoppers Myerslopiidae, new family and Sagmatiini, new tribe (Homoptera: Membracoidea). *Invertebrate Taxonomy* **13**: 207–235.
- Handlirsch, A. 1906–1908. *Die fossilen Insekten und die Phylogenie der rezenten Formen. Ein Handbuch für Paläontologen und Zoologen*. Leipzig: Engelmann.
- 1925. Palaeontology, pp. 117–306. In Schröder, C. (ed), *Handbuch der Entomologie* **3**. Jena: G. Fischer.
- 1939. Neue Untersuchungen über die fossilen Insekten mit Ergänzungen und Nachträgen sowie Ausblicken auf phylogenetische, palaeogeographische und

- allgemein biologische Probleme, II Theil. *Annalen des Naturhistorischen Museums in Wien* **48**: 1–240.
- Hasegawa, E. and Kasuya, E. 2006. Phylogenetic analysis of the insect order Odonata using 28S and 16S rDNA sequences: a comparison between data sets with different evolutionary rates. *Entomological Science* **9**: 55–66.
- Hasiotis, S. T. and Dubiel, R. F. 1995. Termite (Insecta: Isoptera) nest ichnofossils from the Triassic Chinle Formation, Petrified Forest National Park, Arizona. *Ichnos* **4**: 119–130.
- , Peterson, F., Demko, T. M. and Turner, C. 1997. Giant termite (Insecta: Isoptera) nests from the base of the Upper Jurassic Morrison Formation, northwestern New Mexico. *Geological Society of America Abstracts with Program* **28**: A461.
- Heads, S. W. 2008. The first fossil Proscopiidae (Insecta, Orthoptera, Eumastacidae) with comments on the historical biogeography and evolution of the family. *Palaeontology* (in press).
- Martill, D. M. and Loveridge, R. F. 2005. An exceptionally preserved antlion (Insecta, Neuroptera) with colour pattern preservation from the Cretaceous of Brazil. *Palaeontology* **48**: 1409–1417.
- Hennig, W. 1953. Kritische Bemerkungen zum phylogenetischen System der Insekten. *Beiträge zur Entomologie* **3**: 1–85.
- 1954. Flügelgeäder und System der Dipteren. *Beiträge zur Entomologie* **4**: 245–388.
- 1969. *Die Stammesgeschichte der Insekten*. Frankfurt: W. Kramer.
- 1973. Diptera (Zweiflügler). In *Handbuch der Zoologie*, IV. Band: Arthropoda – 2. Hälfte: Insecta – 2. Teil: Spezielles, 31. Berlin: Walter de Gruyter.
- 1981. *Insect Phylogeny*. Chichester: Wiley.
- Herczek, A. and Popov, Yu. A. 2001. Redescription of the oldest plant bugs from the Upper Jurassic of the Southern Kazakhstan (Heteroptera: Cimicomorpha, Miridae). *Annals of the Upper Silesian Museum in Bytom, Entomology* **10–11**: 121–128.
- Herman, L. H. 2001. Catalog of the Staphylinidae (Insecta: Coleoptera). 1758 to the end of the second millennium. I. introduction, history, biographical Sketches, and omaliine group. *Bulletin of the American Museum of Natural History* **265**: 1–650.
- Hoch, H., Deckert, J. and Wessel, A. 2006. Vibrational signalling in a Gondwanan relict insect (Hemiptera: Coleorrhyncha: Peloridiidae). *Biology Letters* **2**: 222–224.
- Hölldobler, B. and Wilson, E. O. 1990. *The Ants*. Cambridge: Belknap Press of Harvard University Press.
- Holzinger, W. E., Kammerlander, I., Bourgoin, Th., Chan, K. L. and Campbell, B. C. 2001. Towards a phylogeny of the Cixiidae (Fulgoromorpha) and its major subgroups: preliminary results. *2nd European Hemiptera Congress*, 20–24 June 2001, Fiesa, Slovenia: 19.
- , Emeljanov, A. F. and Kammerlander, I. 2002. The family Cixiidae – a review, pp. 113–138. In Holzinger, W. (ed), *Zikaden – Leafhoppers, Planthoppers and Cicadas (Insecta: Hemiptera: Auchenorrhyncha)*. Denisia, 04, Zugleich Kataloge Neue Folge Nr. 176, Linz: OÖ. Landesmuseums.
- Hong, Y. 1982. *Mesozoic Fossil Insects of Jiuquan Basin, Gansu Province*. Beijing: Geological Publishing House.
- Hörnschemeyer, T. 1994. Ein fossiler Tenebrionidae *Ceropria? messelense* n. sp. (Coleoptera: Tenebrionidae: Diaperinae) aus dem Mitteleozän der Grube Messel bei Darmstadt. *Courier Forschungsinstitut Senckenberg* **170**: 75–83.
- Hovmöller, R. 2006. *Molecular phylogenetics and taxonomic issues in dragonfly systematics (Insecta: Odonata)*. PhD thesis, Stockholm University.

- Hubbard, M. D. and Kukalová-Peck, J. 1980. Permian mayfly nymphs: new taxa and systematic characters, pp. 19–31. In Flannagan, F. F. and Marshall, K. E. (eds), *Advances in Ephemeroptera Biology*. New York: Plenum Press.
- and Savage, H. M. 1981. The fossil Leptophlebiidae (Ephemeroptera): A systematic and phylogenetic review. *Journal of Paleontology* **55**: 810–813.
- Hughes, L. and Westoby, M. 1992. Capitula on stick insect eggs and elaiosomes on seeds: convergent adaptations for burial by ants. *Functional Ecology* **6**: 642–648.
- Illies, J. 1968. Ephemeroptera (Eintagsfliegen). *Handbuch der Zoologie IV (Arthropoda) 2 (Insecta)* **5**: 1–63.
- Ivanov, V. D. and Sukatsheva, I. D. 2002. Order Trichoptera Kirby, 1813. The Caddisflies, pp. 199–220. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- James, M. T. 1981. Stratiomyidae, pp. 497–511. In McAlpine, J. F. and Woods, D. M. (eds), *Manual of Nearctic Diptera*. Ottawa: Biosystematics Research Institute.
- Jarvis, K. J., Haas, F. and Whiting, M. 2005. A phylogeny of earwigs (Insecta: Dermaptera) based on molecular and morphological evidence: reconsidering the classification of Dermaptera. *Systematic Entomology* **30**: 442–453.
- Jarzembowski, E. A. 1991. New insects from the Weald Clay of the Weald. *Proceedings of the Geologists' Association* **9**: 71–93.
- , Martínez-Delclòs, X., Bechly, G., Nel, A., Coram, R. and Escuillié, F. 1998. The Mesozoic non-calopterygoid Zygoptera: description of new genera and species from the Lower Cretaceous of England and Brazil and their phylogenetic significance (Odonata, Zygoptera, Coenagrionoidea, Hemiphlebioidea, Lestoidea). *Cretaceous Research* **19**: 403–444.
- Jell, P. A. and Duncan, P. M. 1986. Invertebrates, mainly insects, from the freshwater Lower Cretaceous, Koonwarra Fossil Bed (Korumburra Group), South Gippsland, Victoria, pp. 111–205. In *Plants and Invertebrates from the Lower Cretaceous Koonwarra Fossil Bed, South Gippsland, Victoria. Memoirs of the Association of Australasian Palaeontologists* **3**.
- Kaszab, Z. and Schawaller, W. 1984. Eine neue Schwarzkäfer-Gattung und -Art aus Dominikanischem Bernstein (Coleoptera, Tenebrionidae). *Stuttgarter Beiträge zur Naturkunde, Serie B* **109**: 1–6.
- Key, K. H. L. 1991. Phasmatodea, pp. 394–403. In Naumann, D. (ed), *The Insects of Australia: a Textbook for Students and Research Workers*, vol. 1, 2nd edn. Carlton: Melbourne University Press.
- Kimsey, L. S. 1991. Relationships among the tephid wasp subfamilies (Hymenoptera). *Systematic Entomology* **16**: 427–438.
- Kirejtshuk, A. G. and Ponomarenko, A. G. 1990. Fossil beetles of the Peltidae and Nitidulidae families (Coleoptera). *Paleontological Journal* **24**: 79–90.
- Klass, K. D. 1997a. The external male genitalia and the phylogeny of Blattaria and Mantodea. *Bonner Zoologische Monographien* **42**: 1–341.
- 1997b. The ovipositor of Dictyoptera (Insecta): Homology and ground-plan of the main elements. *Zoologischer Anzeiger* **236**: 69–101.
- 2001. Morphological evidence on Blattaria phylogeny: “Phylogenetic histories and stories” (Insecta, Dictyoptera). *Deutsche Entomologische Zeitschrift* **48**: 223–265.
- Kluge, N. J. 1986. Recent mayfly species (Ephemeroptera, Heptageniidae) in Baltic amber. *Paleontological Journal* **2**: 111–112.
- 1996. A new suborder of Thysanura for the Carboniferous insect originally described as larva of *Bojophlebia*, with comments on characters of the orders Thysanura and Ephemeroptera. *Zoosystematica Rossica* **4**: 71–75.

- 1998. Phylogeny and higher classification of Ephemeroptera. *Zoosystematica Rossica* **7**: 255–269.
- 2004. *The Phylogenetic System of Ephemeroptera*, Dordrecht: Kluwer Academic Publishers.
- , Studemann, D., Landolt, P. and Gonser, T. 1995. A reclassification of Siphlonuroidea (Ephemeroptera). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* **68**: 103–32.
- , Godunko, R. J. and Krzeminski, W. 2006. A new mayfly family (Insecta: Ephemeroptera) from Eocene Baltic amber. *Annales Zoologici* **56**: 181–185.
- Koch, M. 1997. Monophyly and phylogenetic position of the Diplura (Hexapoda). *Pedobiologia* **41**: 9–12.
- 2003. Towards a phylogenetic system of the Zygentoma. *Entomologische Abhandlungen* **61**: 122–124.
- Königsmann, E. 1976. Das phylogenetische System der Hymenoptera, Sapygidae. *Deutsche Entomologische Zeitschrift N.F.* **25(IV–V)**: 406–409.
- Kozlov, M. V., Ivanov, V. D. and Rasnitsyn, A. P. 2002. Order Lepidoptera Linné, 1758. The butterflies and moths, pp. 220–227. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- Krell, F.-T. 2000. The fossil record of Mesozoic and Tertiary Scarabaeoidea (Coleoptera: Polyphaga). *Invertebrate Taxonomy* **2000**: 871–905.
- Krishna, K. 1990. Isoptera, pp. 76–81. In Grimaldi, D. A. (ed), *Insects from the Santana Formation, Lower Cretaceous, of Brazil. Bulletin of the American Museum of Natural History* **195**.
- Kristensen, N. P. 1991. Phylogeny of extant hexapods, pp. 125–140. In Naumann, I. D. (ed), *The Insects of Australia: a Textbook for Students and Research Workers*, vol. 1, 2nd edn. Melbourne: Melbourne University Press.
- 1995. Forty years' insect phylogenetic systematics: Hennig's "Kritische Bemerkungen" and subsequent developments. *Zoologische Beiträge (NF)* **36**: 83–124.
- 1998. The groundplan and basal diversification of the hexapods, pp. 281–293. In Fortey, R. A. and Thomas, R. H. (eds), *Arthropod Relationships*. London: Chapman and Hall.
- Krzeminski, W. and Evenhuis, N. L. 2000. Review of Diptera paleontological records, pp. 535–564. In Papp, L. and Darvas, B. (eds), *Contributions to a Manual of Palaearctic Diptera*, vol. 1 L. Budapest: Science Herald.
- Kukalová-Peck, J. 1978. Origin and evolution of insect wings and their relation to metamorphosis, as documented by the fossil record. *Journal of Morphology* **156**: 53–125.
- 1983. Origin of the insect wing and wing articulation from the arthropodan leg. *Canadian Journal of Zoology* **61**: 1618–1669.
- 1985. Ephemeroptery wing venation based upon new gigantic Carboniferous mayflies and basic morphology, phylogeny, and metamorphosis of pterygote insects (Insecta, Ephemeroptera). *Canadian Journal of Zoology* **63**: 933–955.
- 1987. New Carboniferous Diplura, Monura, and Thysanura, the hexapod ground plan, and the role of thoracic side lobes in the origin of wings (Insecta). *Canadian Journal of Zoology* **65**: 2327–2345.
- 1991. Fossil history and the evolution of hexapod structures, pp. 141–179. In Naumann, I. D. (ed), *The Insects of Australia: a Textbook for Students and Research Workers*, vol. 1. Melbourne: Melbourne University Press.
- Kuschel, G. 2003. Nemonychidae, Belidae, Brentidae (Insecta: Coleoptera: Curculionoidea). *Fauna of New Zealand* **45**: 1–100. Manaaki Whenua Press.

- Kwok, H. K. and Corlett, R. T. 2002. Seasonality of forest invertebrates in Hong Kong, South China. *Journal of Tropical Ecology* **18**: 637–644.
- Ladiges, P. Y., Nelson, G. and Grimes, J. 1997. Subtree analysis, *Nothofagus* and Pacific biogeography. *Cladistics* **13**: 125–129.
- Lameere, A. 1917. Étude sur l'évolution des Ephémères. *Bulletin de Societe Zoologique de France* **42**: 41–81.
- Lauterbach, K. E. 1973. Schlüsselereignisse in der Evolution der Stammgruppe der Euarthropoda. *Zoologische Beiträge, Neue Folge* **19**: 251–299.
- Lawrence, J. F. 1999. The Australian Ommatidae (Coleoptera: Archostemata): new species, larva and discussion of relationships. *Invertebrate Taxonomy* **13**: 369–390.
- and Newton, Jr. A. F. 1995. Families and subfamilies of Coleoptera (with selected genera, notes, references and data on family-group names), pp. 779–1006. In Pakaluk, J. and Slipinski, S. A. (eds), *Biology, Phylogeny, and Classification of Coleoptera*, vol. 2. Warsaw: Muzeum I Instytut Zoologii PAN.
- Lin, Q.-B., and Huang, D. Y. 2001. Description of *Caenoephemera shangyuensis*, gen. nov., sp. nov. (Ephemeroptera), from the Yixian Formation. *Canadian Entomologist* **133**: 747–754.
- Nel, A. and Huang, D.-Y. 2002. Phylogenetic analysis of the Mesozoic dragonfly family Liupanshaniidae (Insecta: Aeshnoptera: Odonata). *Cretaceous Research* **23**: 439–444.
- Lo, N., Bandi, C., Watanabe, H., Nalepa, Ch. and Beninati, T. 2003. Evidence for coeladogenesis between diverse dictyopteran lineages and their intracellular endosymbionts. *Molecular Biology and Evolution* **20**: 907–913.
- Lohmann, H. 1996. Das Phylogenetische System der Anisoptera (Odonata). *Entomologische Zeitschrift* **106(6)**: 209–252; **106(7)**: 253–296; **106(9)**: 360–367.
- Lomhold, O. 1982. On the origin of the bees (Hymenoptera: Apidae, Sphecidae). *Entomologica Scandinavica* **13**: 185–190.
- Lubkin, S. H. 2003. *Paracupes svitkoi* (Coleoptera: Cupedidae), a new species from the Cretaceous of New Jersey. *Acta Zoologica Cracoviensia* **46** (suppl. Fossil Insects): 189–194.
- and Engel, M. S. 2005. *Permocoleus*, New Genus, the First Permian Beetle (Coleoptera) from North America. *Annals of the Entomological Society of America* **98**: 73–76.
- Maddison, D. R. 1995. Coptoclavidae. <http://tolweb.org/tree/tree?group=Coptoclavidae>.
- Maisey, J. G. (ed.). 1991. *Santana Fossils: an Illustrated Atlas*. Neptune City, NJ: T.F.H. Publications.
- Makarkin, V. N. and Menon, F. 2007. First record of fossil ‘rapismatid-like’ Ithionidae (Insecta, Neuroptera) from the Lower Cretaceous Crato Formation of Brazil. *Cretaceous Research* (in press).
- Malyshev, S. I. 1968. *Genesis of the Hymenoptera and the Phases of their Evolution*. London Methuen & Co.
- Marden, J. H. and Thomas, M. A. 2005. Rowing locomotion by a stonefly that possesses ancestral pterygote condition of co-occurring wings and abdominal gills. *Biological Journal of the Linnean Society* **79**: 341–349.
- Marshall, J. A and Haes, E. C. M. 1988. *Grasshoppers and allied insects of Great Britain and Ireland*. Colchester: Harley Books.
- Martill, D. M. (ed.) 1993. *Fossils of the Santana and Crato Formations, Brazil*. The Palaeontological Association Field Guides to Fossils, 5. London: The Palaeontological Association.

- 1994. Fake fossils from Brazil. *Geology Today* May/June **1994**: 111–115.
- and Frey, E. 1995. Colour patterning preserved in Lower Cretaceous birds and insects: the Crato-Formation of N.E. Brazil. *Neues Jahrbuch für Geologie und Paläontologie Monatshefte* **2**: 118–128.
- and Nel, A. 1996. A new dragonfly from the Crato Formation (Lower Cretaceous, Aptian) of N. E. Brazil. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* **5**: 279–292.
- and Davis, P. G. 1998. Did dinosaurs come up to scratch? *Nature* **396**: 528–589.
- Martínez, S. 1982. Catalogo sistemático de los insectos fosiles de America del Sur. *Revista de la Facultad de Humanidades y Ciencias, Serie Ciencias de la Tierra* **1**: 29–83.
- Martínez-Delclós, X. 1989. *Chresmoda aquatica* n. sp. insecto Chresmodidae del Cretácico inferior de la Sierra del Montsech (Lleida, España). *Revista Española de Paleontología* **4**: 67–74.
- , Nel, A., Azar, D., Bechly, G., Dunlop, J. A., Engel, M. S. and Heads, S. 2007. The enigmatic, Mesozoic family Chresmodidae (Polyneoptera: Archaeorthoptera): New palaeobiological and phylogenetic data, with the description of a new species from Brazil – New palaeobiological and phylogenetical data. *Neues Jahrbuch für Geologie und Paläontologie – Abhandlungen* (in press).
- Martins-Neto, R. G. 1987a. Um novo gênero de Orthoptera (Insecta, Grylloidea) da Formação Santana, Bacia do Araripe (Cretáceo Inferior) nordeste do Brasil. *Anais 10º Congresso Brasileiro de Paleontologia*, Rio de Janeiro: 599–609.
- 1987b. A paleoentomofauna Brasileira: estagio actual do concimento. *Anais 10º Congresso Brasileiro Paleontologia*, Rio de Janeiro v. **II**: 567–591.
- 1987c. Descrição de três novos géneros e três novas espécies de Orthoptera (Insecta, Aridoidea) da Formação Santana, Bacia do Araripe (Cretáceo Inferior) nordeste do Brasil, representando três familiias, sendo que duas novas: Archaeopneumoridae nov. fam. e Bouretidae nov. fam. *Anais da Academia Brasileira de Ciências* **59**: 444.
- 1988a. A new fossil insect (Homoptera, Cixiidae) from the Santana Formation (Lower Cretaceous), Araripe Basin, Northeast Brasil. *Interciencia* **13**: 313–316.
- 1988b. Neuropteros (Insecta: Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, nordeste do Brasil. I. Família Chrysopidae. *Anais da Academia Brasileira de Ciências* **60**: 189–201.
- 1989a. A new fossil insect (Homoptera, Cixiidae) from the Santana Formation (Lower Cretaceous) Araripe Basin, Northeast Brazil. *Acta Geologica Leopoldensia* **26**: 7–14.
- 1989b. Primeiro registro de Phasmatodea (Insecta: Orthopteromorpha) na Formação Santana, Bacia do Araripe (Cretáceo Inferior), nordeste do Brasil. *Acta Geologica Leopoldensia (Estudos Tecnológicos)* **12**: 91–104.
- 1990a. Primeiro registro de Dermaptera (Insecta, Orthopteromorpha) na Formação Santana (Cretáceo inferior), Bacia do Araripe, nordeste do Brasil. *Revista Brasileira de Entomologia* **34**: 775–784.
- 1990b. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. VI. Ensaio filogenético das espécie do genera *Blittersdorffia* Martins-Neto and Vulcano, com descrição de nova espécie. *Acta Geologica Leopoldensia* **13**: 3–12.
- 1990c. The family Locustopsidae (Insecta, Caelifera) in the Santana Formation (Lower Cretaceous, northeast Brazil). I. – Description of two new species of the genus *Locustopsis* Handlirsch and three new species of the genus *Zessinia* n. gen. *Atlas do I Simpósio sobre a Bacia do Araripe e Bacias Interiores do n Nordeste Crato*, 14 a 16 Junho de 1990: 277–91.

- 1990d. Um novo gênero e duas nova espécies de Tridactylidae (Insecta, Orthopteroidea) na Formação Santana (Cretáceo Inferior do nordeste do Brasil). *Annales Academia Brasil Ciencias* **62**: 51–59.
- 1991a. Sistemática dos Ensifera (Insecta, Orthopteroida) da Formação Santana, Cretáceo Inferior do nordeste do Brasil. *Estudio Technológico, Acta Geologica Leopoldensia* **32**: 3–162.
- 1991b. *Cratogryllus ciguaeli*, nova espécie de Ensifera (Insecta, Grylloidea) da Formação Santana (Cretáceo Inferior) Bacia do Araripe, nordeste do Brasil. *Acta Geologica Leopoldensia* **33**: 153–156.
- 1991c. Evidências de especiação alocrônica na fauna de Ensifera (Insecta, Orthopteroida) da Formação Santana, Cretáceo do nordeste do Brasil. *Revista de Geologia* **4**: 61–80.
- 1991d. A Paleoentomofauna do Nordeste Brasileiro: Estado da Arte. *Atas XIV Simposio de Geologia do Nordeste, Recife* **12**: 59–62.
- 1992a. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. V. Aspectos filogenéticos, paleoecológicos, paleobiogeográficos e descrição de novos taxa. *Anais da Academia Brasileira de Ciências* **64**: 117–148.
- 1992b. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. VII. Palaeoleontinae, nova subfamília de Myrmeleontidae e descrição de novos táxons. *Revista Brasileira de Entomologia* **36**: 803–815.
- 1992c. Nova ocorrência, variabilidade morfológica e relações filogenéticas do gênero *Cratoelcana* Martins-Neto, 1991 (Insecta, Ensifera, Elcanidae), da Formação Santana, Bacia do Araripe, Brasil. *Revista Brasil Entomologia* **36**: 817–830.
- 1992d. A paleoentomofauna das bacias mesozoicas brasileiras e seu potencial para possíveis correlações bioestratigráficas entre depósitos não-marinhos. In I. Encontro sobre a sedimentação continental das bacias mesozoicas brasileiras. *Acta Geologica Leopoldensia* **36**: 125–126.
- 1994. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. IX. Primeiros resultados da composição da fauna e descrição de novos táxons. *Acta Geologica Leopoldensia* **17**: 269–288.
- 1995a. Araripelocustidae, fam. n. uma nova família de gafanhotos (Insecta, Caelifera) da Formação Santana, Cretáceo Inferior do nordeste do Brasil. *Revista Brasil Entomologia* **39**: 311–319.
- 1995b. Complementos ao estudo sobre os Ensifera (Insecta, Orthopteroida) da Formação Santana, Cretáceo Inferior do nordeste do Brasil. *Revista Brasil Entomologia* **39**: 321–345.
- 1996a. New mayflies (Insecta, Ephemeroptera) from the Santana Formation (Lower Cretaceous), Araripe Basin, Northeastern Brazil. *Revista Española de Paleontología* **11**: 177–192.
- 1996b. Interação inseto / flora e considerações sobre a origem das angiospermas. *Revista UnG, Série Geociências* **1**: 31–36.
- 1997a. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. X. Descrição de novos táxons (Chrysopidae, Babinskaiidae, Myrmeleontidae, Ascalaphidae e Psychopsidae). *Revista Universidade de Guarulhos, Série Ciências Exatas e Technológicas* **2**: 68–83.
- 1997b. *Cratotetraspinus* nov. nom., novo nome para *Tetraspinus* Martins-Neto, 1995 (nom. preoc.). *Revista Universidade Guarulhos, Geociências* **2**: 105.

- 1998a. A new genus of the family Locustopsidae (Insecta, Caelifera) in the Santana Formation (Lower Cretaceous, northeast Brazil). *Revista Española de Paleontología* **13**: 133–138.
- 1998b. A new subfamily of Ensifera (Insecta, Grylloidea) from the Santana Formation (Lower Cretaceous), Araripe Basin, NE Brazil. *Proceedings of the First International Palaeoentomological Conference*, Moscow 1998: 91–97.
- 1998c. *Conan barbarica* n. gen. et n. sp. (Insecta, Coleoptera, Coptoclavidae) – uma gigantesca larva da Formação Santana, Cretáceo Inferior, Bacia do Araripe, Brasil. *Geociências São Paulo* **17**: 109–114.
- 1998d. Neurópteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. XI. Descrição de novos táxons de Myrmeleontidae (Palaeoleontinae e Pseudonymphinae). *Revista Universidade de Guarulhos, Série Ciências Biológicas e da Saúde* **3**: 38–42.
- 1998e. The paleoentomo fauna from Tremembé Formation (Taubaté Basin) Oligocene of São Paulo State: new Hemiptera, Auchenorrhyncha, Hymenoptera, Coleoptera, Lepidoptera and Diptera (Parte II) (Insecta). *Revista Universidade Guaralhos, Serie Geociências* **3**: 58–70.
- 1998f. Novos registro de Palaeointinideos (Insecta: Hemiptera) na formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. *Acta Geologica Leopoldensia* **21**: 69–74.
- 1999a. Estado actual del conocimiento de la paleoentomofauna brasileña. *Revista de la Sociedad Entomologica Argentina* **58**: 71–85.
- 1999b. New genus and new species of Lepidoptera (Insecta, Eolepidopterigidae) from Santana Formation (Lower Cretaceous, northeast Brazil). *5º Simpósio sobre o Cretáceo do Brasil*, Rio Claro. Boletim: 531–535.
- 2000. Remarks on the neuropterofauna (Insecta, Neuroptera) from the Brazilian Cretaceous, with keys for the identification of the known taxa. *Acta Geologica Hispanica* **35**: 97–118.
- 2001a. Review of some Insecta from Mesozoic and Cenozoic Brazilian deposits with descriptions of new taxa. *Acta Geologica Leopoldensia* **24**: 115–124.
- 2001b. Primeiro registro de Trichoptera (Insecta) na Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil, com descrição de sete novos táxons. *Simpósio sobre a Bacia do Araripe e bacias interiores do Nordeste*, 1 e 2, Crato, 1990/1997. Boletim: 212–226.
- 2002a. The Santana Formation Paleoentomofauna reviewed. Part I. Neuropteroida (Neuroptera and Raphidioptera): systematic and phylogeny, with descriptions of new taxa. *Acta Geologica Leopoldensia* **25**: 35–66.
- 2002b. *Insetos fósseis como bioindicadores em depósitos sedimentares: um estudo de caso para o mesozóico sul-americano*. São Leopoldo: CPGEO, UNISINOS.
- 2003a. Systematics of the Caelifera (Insecta, Orthopteroidea) from the Santana Formation, Araripe Basin (Lower Cretaceous, northeast Brazil), with a description of new genera and species. *Acta Zoologica Cracoviensis* **46** (suppl.): 205–228.
- 2003b. The fossil Tabanids (Diptera Tabanidae): When they began to appreciate warm blood and when they began transmit diseases? *Memorias do Instituto Oswaldo Cruz* **98** (suppl. I): 29–34.
- 2004. New Neuroptera (Nymphidae and Araripeneuridae) from Santana Formation (Lower Cretaceous, Araripe Basin, Northeast Brasil). *Acta Geologica Leopoldensia* **27**.

- 2005a. The Santana Formation palaeoentomofauna reviewed: Polyneoptera (Isoptera, Blattoptera, Dermaptera, Orthoptera) with description of six new species, one new genus and one new combination. *3rd International Congress of Palaeoentomology, Abstracts*, 7–11 February 2005, Pretoria, South Africa. **1**: 24.
- 2005b. Estágio atual da paleoartropologia brasileira: Hexápodes, miriápodes, crustáceos (Isopoda, Decapoda, Eucrustacea e Copepoda) e quelicerados. *Arquivos do Museu Nacional, Rio de Janeiro* **63**: 471–494.
- and Vulcano, M. A. 1989a. Primeiro registro de Raphidioptera (Neuropteroidea) na Formação Santana (Cretáceo Inferior), Bacia do Araripe, nordeste do Brasil. *Revista Brasileira de Entomologia* **34**: 241–249.
- and Vulcano, M. A. 1989b. Amphiesmenoptera, (Trichoptera + Lepidoptera) na Formação Santana (Cretáceo Inferior) Bacia do Araripe, Nordeste do Brasil. I – Lepidoptera (Insecta). *Anais da Academia Brasileira de Ciências* **61**: 459–465.
- and Vulcano, M. A. 1989c Neuropteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. I. Família Chrysopidae. *Anais Academia Brasileira de Ciências* **60**: 189–201.
- and Caldas, E. B. 1990. Efêmeras escavadoras (Insecta, Ephemeroptera, Ephemeroidea) na Formação Santana (Cretáceo Inferior), Bacia do Araripe Nordeste do Brasil: Descrição de três novos gêneros e três novas espécies (ninfas). *Atas do I Simpósio sobre a Bacia do Araripe e Bacias Inteiores do Nordeste Crato*, 1990: 265–275.
- and Vulcano, M. A. 1990a. Neuropteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, nordeste do Brasil. III – Superfamília Mantispoidea. *Revista Brasileira de Entomologia* **34**: 619–625.
- and Vulcano, M. A. 1990b. Neuropteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, nordeste do Brasil. VI – Ensaio filogenético das espécies do gênero *Blittersdorfia*, com descrição de nova espécie. *Acta Geologica Leopoldensia* **31**: 3–12.
- and Valarelli, J. V. 1991. Primeiro registro de Eumastacoidea (Insecta, Caelifera) da Formação Santana, Cretáceo Inferior do nordeste do Brasil. *Anais da Academia Brasileira de Ciências* **63**: 91–92.
- and Nel, A. 1992. Un nouveau fossile de Raphidoptère de la formation Santana, Crétacé inférieur du Brésil (Neuropteroidea, Raphidoptera). *Bulletin de la Société Entomologique de France* **97**: 425–428.
- and Santos, J. C. K. 1994. Um novo gênero e uma nova espécie de Mutuca (Insecta, Diptera, Tabanidae) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. *Acta Geologica Leopoldensia* **39**: 289–297.
- and Vulcano, M. A. 1997. Neuropteros (Insecta, Planipennia) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, nordeste do Brasil. VIII – Descrição de novos taxa de Myrmeleontidae, Ascalaphidae e Nemopteridae. *Revista Universidade Guarulhos, Série Ciências Biológicas* **2**: 64–81.
- and Petrulevičius, J. F. 1998. Mesozoic and Cenozoic Orthopteroida from South America. *1st Paleoentomological Conference, Abstracts*, 30 August–4 September 1998, Moscow: 24.
- and Szwedo, J. 2007. Taxonomic changes in fossil Cixiidae (Insecta: Hemiptera: Fulgoromorpha) 1 – *Cretofennahia* nom. nov. pro *Fennahia* Martins-Neto. *Alevisia* **1**: 117.
- , Popov, Yu. A., and Zamboni, J. A. 1999. First South Hemisphere Cretaceous record of Coreoidea (Insecta, Heteroptera) from Santana Formation (Lower Cretaceous, Northeast Brazil), representing a new genus et species. *Bulletim do 5^o Simposio Sobre o Cretáceo do Brasil*: 525–530.

- , Ribeiro-Júnior, C. and Prezoto, F. 2006. New fossils (Isoptera: Hodotermitidae), from the Santana Formation (Lower Cretaceous, Araripe Basin, Northeast Brazil), with descriptions of new taxa including a new subfamily. *Sociobiology* **47**: 125–134.
- , Melo, A. C. and Prezoto, F. 2007. A new species of wasp (Symphyta, Sepulcidae) from the Santana Formation (Lower Cretaceous, Northeast Brazil). *Journal of the Entomological Research Society* **9(1)**: 1–6.
- Martynov, A. V. 1925. Über zwei Grundtypen der Flügel bei den Insecten und ihre Evolution. *Zeitschrift zur Morphologie und Ökologie der Tiere* **4**: 465–501.
- Martynova, O. M. 1962 (reprinted 1991). Order Phasmatodea, pp. 213–215. In Rohdendorf, B. B. (ed), *Fundamentals of Paleontology*, vol. 9, *Arthropoda, Tracheata, Chelicerata*. Washington DC: Smithsonian Institution Libraries and The National Science Foundation.
- Matzke, D. and Klass, K.-D. 2005. Reproductive biology and nymphal development in the basal earwig *Tagalina papua* (Insecta: Dermaptera: Pygidicranidae), with a comparison of brood care in Dermaptera and Embioptera. *Entomologische Abhandlungen* **62**: 99–116.
- Mazzarolo, L. A. and Amorim, D. S. 2000. *Cratomyia macrorrhyncha*, a Lower Cretaceous brachyceran fossil from the Santana Formation, Brazil, representing a new species, genus and family of the Stratiomyomorpha (Diptera). *Insect Systematics & Evolution* **31**: 91–102.
- Mazzoni, A. F. and Hünicken, M. A. 1984. Ontogenia de los notonectidos (Insecta, Heteroptera) del Cretácico inferior de San Luis, Argentina. *Memoria III Congreso Latinoamericano de Paleontología*: 388–395.
- and Hünicken, M. A. 1987. Corixidae (Insecta, Heteroptera) del Cretácico Inferior de la Sierra del Gigante, San Luis, Argentina. *Actas IV Congresso Latinoamericano Palaeontologia, Bolivia* **2**: 731–738.
- McCafferty, W. P. 1990. Ephemeroptera, pp. 20–50. In Grimaldi, D. A. (ed), Insects from the Santana Formation, Lower Cretaceous, of Brazil. *Bulletin of the American Museum of Natural History* **195**.
- 1991. Toward a phylogenetic classification of the Ephemeroptera (Insecta): a commentary on systematics. *Annals of the Entomological Society of America* **84**: 343–360.
- 1997. Ephemeroptera, pp. 89–117. In Poole, R. W. and Gentilli, P. (eds), *Nomina Insecta Nearctica, a Checklist of the Insects of North America*, vol. 4. *Non-holometabolous Orders*. Rockville, MD: Entomological Informational Service.
- McCafferty, W. P. 1998. Ephemeroptera and the great American interchange. *Journal of the North American Benthological Society* **17**: 1–20.
- 2004. Higher classification of the burrowing mayflies (Ephemeroptera: Shappodonta). *Entomological News* **115**: 84–92.
- and Edmunds, Jr. G. F. 1979. The higher classification of the Ephemeroptera and its evolutionary basis. *Annals of the Entomological Society of America* **72**: 5–12.
- McAlpine, J. F. 1981. Morphology and terminology – adults, pp. 9–63. In McAlpine, J. F. and Woods, D. M. (eds), *Manual of Nearctic Diptera*, vol. 1. Ottawa: Biosystematics Research Institute.
- Melo, G. A. R. 1999. Phylogenetic relationships and classification of the major lineages of Apoidea (Hymenoptera), with emphasis on the crabronid wasps. *Scientific Papers of the Natural History Museum, University of Kansas* **14**: 1–55.
- Mendes, L. F. 1997a. A first contribution to the study of the Dominican amber Zygentoma (Insecta), family Ateluridae. *Pedobiologia* **41**: 40–43.
- 1998a. Second contribution to the study of the Dominican amber Zygentoma (Insecta): Family Lepismatidae. *Canadian Entomologist* **130**: 899–904.

- Mendes, M. 1991a. A utilização de caracteres morfológicos alternativos na Sistemática dos Blattoptera (Insecta) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. *XII Congresso Brasileiro de Paleontologia, 1991, São Paulo, Boletim de Resumos 1*: 54.
- 1991b. Raphidiomimidae (Insecta, Blattoptera) da Formação Santana (Cretáceo Inferior), Bacia do Araripe, Nordeste do Brasil. *XII Congresso Brasileiro de Paleontologia, 1991, São Paulo, Boletim de Resumos 1*: 55.
- 1993. Um novo gênero da Família Blattidae (Insecta, Blattoptera) na Formação Santana, Cretáceo do Nordeste do Brasil. *XIII Congresso Brasileiro de Paleontologia, 1993, São Leopoldo, Boletim de Resumos 1*: 115.
- 1995. Três Novas Espécies do Gênero Mesoblattina (Insecta, Blattoptera) na Formação Santana, Cretáceo do Nordeste do Brasil. *XIV Congresso Brasileiro de Paleontologia, 1995, Uberaba, Boletim de Resumos 1*: 98.
- 1997b. Duas Novas Espécies do Gênero Mesoblattinopsis (Insecta, Blattoptera, Blattidae) na Formação Santana, Cretáceo do Nordeste do Brasil. *XV Congresso Brasileiro de Paleontologia, 1997, São Pedro, Boletim de Resumos 1*: 58.
- 1998a. An attempt at taphonomic simulation in blattopteras from Araripe (Lower Cretaceous), from the North East of Brazil. *1st Paleoentomological Conference, Abstracts, 30 August–4 September 1998, Moscow*: 25.
- 1998b. An Attempt at taphonomic simulation in Blattopteras from Araripe (Lower Cretaceous) from the NE of Brazil. *AMBA projects International 98*: 99–101.
- 2000. Novos Blattodeas (Insecta, Blattopteroidea) da Formação Santana, Cretáceo Inferior do Nordeste do Brasil. *Revista Universidade de Guarulhos - Ciências Exatas e Tecnológicas 5(6)*: 25–35.
- Menon, F. 2005. New record of Tettigarctidae (Insecta, Hemiptera, Cicadoidea) from the Lower Cretaceous of Brazil. *Zootaxa 1087*: 53–58.
- and Heads, S. W. 2005. New species of Palaeontinidae (Insecta: Cicadomorpha) from the Lower Cretaceous Crato Formation of Brazil. *Stuttgarter Beiträge zur Naturkunde, Serie B 357*: 1–11.
- , Heads, S. W. and Martill, D. M. 2005. New Palaeontinidae (Insecta: Cicadomorpha) from the Lower Cretaceous Crato Formation of Brazil. *Cretaceous Research 26*: 837–844.
- Michener, C. D. 2000. *The Bees of the World*. Baltimore, MD: The John Hopkins University Press.
- Mickoleit, G. 1973. Über den Ovipositor der Neuropteroidea und Coleoptera und seine phylogenetische Bedeutung (Insecta, Holometabola). *Zeitschrift für Morphologie der Tiere 74*: 37–64.
- Misof, B. and Rickert, A. 1999a. A molecular approach to the phylogeny of extant Odonata. *Zoology 102* (suppl. II): 9.
- and Rickert, A. 1999b. A molecular phylogeny of Anisoptera. *Zoology 102* (suppl. II): 9.
- Mostovski, M. B. and Jarzembowski, E. A. 2000. The first brachycerous flies (Diptera: Rhagionidae) from the Lower Jurassic of Gondwana. *Paleontological Journal 34* (suppl. 3): 367–369.
- and Martínez-Delclós, X. 2000. New Nemestrinoidea (Diptera: Brachycera) from the Upper Jurassic – Lower Cretaceous of Eurasia, taxonomy and paleobiology. *Entomological Problems 31*: 137–148.
- , Jarzembowski, E. A. and Coram, R. A. 2003. Horseflies and athericids (Diptera: Tabanidae, Athericidae) from the Lower Cretaceous of England and Transbaikalia. *Paleontological Journal 37*: 162–169.

- Moulds, M. S. 1990. *Australian Cicadas*. Kensington, NSW: New South Wales University Press.
- 2005. An appraisal of the higher classification of cicadas (Hemiptera: Cicadoidea) with special reference to the Australian Fauna. *Records of the Australian Museum* **57**: 375–446.
- Nagatomi, A. and Yang, D. 1998. A review of extinct Mesozoic genera and families of Brachycera (Insecta, Diptera, Orthorrhapha). *Entomologist's Monthly Magazine* **134**: 95–192.
- and Rozkosny, R. 2000. Xylomyidae, pp. 369–378. In Papp, L. and Darvas, B. (eds), *Manual of Palaearctic Diptera*, vol. 2. Budapest: Science Herald.
- Nagy, C. G. 1969. A new taxon of the family Heterogynidae Latreille (Hym. Aculeata). *Entomologische Mitteilungen aus dem Zoologischen Staatsinstitut und Zoologischen Museum Hamburg* **64**: 299–303.
- Neboiss, A. 1960. On the family Cupedidae, Coleoptera. *Proceedings of the Royal Society of Victoria* **72**: 12–20.
- Nel, A. 1996. Un Tettigarctidae fossile du Lias Européen (Cicadomorpha, Cicadoidea, Tettigarctidae). *Ecole Pratique des Hautes Etudes, Biologie et Evolution des Insectes* **9**: 83–94.
- and Paicheler, J.-Cl. 1992. Les Heteroptera aquatiques fossiles, état actuel des connaissances (Heteroptera: Nepomorpha et Gerromorpha). *Entomologica Gallica* **3**: 159–182.
- and Martínez-Delclòs, X. 1993. Essai de révision des Aeschnidioidea (Insecta, Odonata, Anisoptera). *Cahiers de Paléontologie* **1993**: 1–99.
- and Paicheler, J.-C. 1993. Les Isoptera fossiles, pp. 103–179. In Nel, A., Martínez-Delclòs, X. and Paicheler, J.-C. (eds), *Essai de révision des Aeschnidioidea [Insecta, Odonata, Anisoptera] / Les Isoptera fossiles [Insecta, Dictyoptera]*. Paris: CNRS Editions (Cahiers de Paléontologie).
- and Escuillé, F. 1994. A new dragonfly from the Lower Cretaceous of Brazil. *Palaeontology* **37**: 923–930.
- and Paicheler, J.-C. 1994a. Les Gomphidae fossiles. Un inventaire critique (Odonata: Gomphidae). *Annales de la Société Entomologie de France* (n.s.) **30**: 55–77.
- and Paicheler, J.-C. 1994b. Les Libelluloidea fossiles autres que Libellulidae. Un inventaire critique (Odonata, Corduliidae, Macromiidae, Synthemistidae, Chlorogomphidae et Mesophlebiidae). *Nouvelle Revue d'Entomologie* (N.S.) **11**: 321–334.
- and Popov, Yu. A. 2000. The oldest known fossil Hydrometridae from the Lower Cretaceous of Brazil (Heteroptera: Gerromorpha). *Journal of Natural History* **34**: 2315–2322.
- and Waller, A. 2006. A giant water bug from the Lower Cretaceous Crato Formation of Brazil (Heteroptera: Belostomatidae: Lethocerinae). *Zootaxa* **1220**: 63–68.
- Sémeria, Y. and Martins-Neto, R. G. 1990. Un Raphidioptera fossile du Crétacé inférieur du Brésil (Neuropteroidea). *Neuroptera International* **6**: 27–37.
- , Martínez-Delclòs, Paicheler, J.-C. and Henrotay, M. 1993. Les “Anisozygoptera” fossiles. Phylogénie et classification. (Odonata). *Martinia, Numéro hors-série* **3**: 1–311.
- , Bechly, G., Jarzembski, E. A. and Martínez-Delclòs, X. 1998. A revision of the fossil petalurid dragonflies (Insecta: Odonata: Anisoptera: Petalurida). *Paleontologia Lombarda* **10**: 1–68.
- , Zarbout, Barale, G. and Philippe, M. 1998. *Liassotettigarcta africana* sp. n. (Auchenorrhyncha: Cicadoidea: Tettigarctidae), the first Mesozoic insect from Tunisia. *European Journal of Entomology* **95**: 593–598.

- , Azar, D., Martínez-Delclós, X. and Makhoul, E. 2004. A new Upper Cretaceous species of *Chresmoda* from Lebanon – a latest representative of Chresmodidae (Insecta: Polyneoptera inc. sed.): first record of homeotic mutations in the fossil record of insects. *European Journal of Entomology* **101**: 145–151.
- , Perrault, G., Perrichot, V. and Néraudeau, D. 2004. The oldest ants in the Lower Cretaceous amber of Charente – Maritime (SW France) (Insecta: Hymenoptera: Formicidae). *Geologica Acta* **2**: 23–29.
- , Bechly, G., Garouste, R., Pohl, B. and Escuillié, F. 2005. A new extraordinary neuropterid family from the Lower Cretaceous Crato Formation of Brazil: a new insect order? (Insecta, Neuroptera). *Cretaceous Research* **26**: 845–852.
- , Martínez-Delclós, X. and Hutin, A.. 2005. Mesozoic chrysopid-like Planipennia: a phylogenetic approach (Insecta: Neuroptera). *Annales de la Société Entomologie de France (n.s.)* **41**: 29–69.
- Newton, Jr, A. F. and Thayer, M. K. 1995. Protopselaphinae new subfamily for *Protopselaphus* new genus from Malaysia, with a phylogenetic analysis and review of the Omaliine Group of Staphylinidae including Pselaphidae (Coleoptera), pp. 219–320. In Pakaluk, J. and Ślipiński, S. A. (eds), *Biology, Phylogeny, and Classification of Coleoptera*, vol 1. Warsaw: Muzeum I Instytut Zoologii PAN.
- Nikolajev, G. V. 1992. Taxonomic criteria and generic composition of mesozoic lamellicorn beetles (Coleoptera, Scarabaeidae). *Paleontological Journal* **26**: 96–111.
- Novokschonov, V. G. 1997. *Early Evolution of Scorpionflies (Insecta: Panorpida)*. Moscow: Nauka.
- 2002. Order Panorpida Latreille, 1802. The scorpionflies, pp. 194–199. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- O'Brien, L. B. 2002. The wild wonderful world of Fulgoromorpha, pp. 83–102. In Holzinger, W. (ed), *Zikaden – Leafhoppers, Planthoppers and Cicadas (Insecta: Hemiptera: Auchenorrhyncha)*. Denisia 04, Zugleich Kataloge, Neue Folge Nr. 176. Linz: OÖ Landesmuseums.
- Ogden, T. H. and Whiting, M. F. 2003. The problem with the “Paleoptera Problem:” sense and sensitivity. *Cladistics* **19**: 432–442.
- 2005. Phylogeny of Ephemeroptera (mayflies) based on molecular evidence. *Molecular Phylogenetics and Evolution* **37**: 625–643.
- Ohl, M. 1995. Die phylogenetischen Beziehungen von Grabwespen und Bienen: Stand der Forschung, Probleme, Perspektiven. *Mitteilungen der Deutschen Gesellschaft für allgemeine und angewandte Entomologie* **10**: 629–632.
- 1996. Die phylogenetischen Beziehungen der Sphecinae (Hymenoptera: Apoidea: “Sphecidae”) aufgrund morphologischer Merkmale des Exoskellets. *Zoologische Beiträge (n. f.)* **37**: 3–40.
- 2000. Das phylogenetische System der aculeaten Hymenopteren sensu stricto – Stand der Forschung, Probleme, Perspektiven. *Beiträge der Hymenopterologen-Tagung Stuttgart 2000*: 10–13.
- and Bleidorn, C. 2006. The phylogenetic position of the enigmatic wasp family Heterogynaidae based on molecular data, with description of a new, nocturnal species (Hymenoptera: Apoidea). *Systematic Entomology* **31**: 321–337.
- Olsen, P. E., Remington, C. L., Cornet, B. and Thompson, K. S. 1978. Cyclic change in Late Triassic lacustrine communities. *Science* **201**: 729–733.
- Oman, P. W., Knight, W. J. and Nielson, M. W. 1990. *Leafhoppers (Cicadellidae): a Bibliography, Generic Check-list and Index to the World Literature 1956–1985*. Wallingford: CAB International Institute of Entomology.
- Osten, T. 1987. Ein neuer Fundort von *Proscolia spectator* Day, 1981 (Hymenoptera, Aculeata). *Entomofauna* **8**: 361–365.

- 1988. Die Mundwerkzeuge von *Proscolia spectator* Day (Hymenoptera: Aculeata). Ein Beitrag zur Phylogenie der “Scolioidea”. *Stuttgarter Beiträge zur Naturkunde Ser. A* **414**: 1–30.
- 1993. The enigmatic Genus *Proscolia*, its distribution and phylogenetic relationships. *Biologia Gallo – Hellenica* **20**: 177–182.
- 2001. Beitrag zur Scoliiden – Fauna der Insel Phuket (Thailand) (Hymenoptera, Scoliidae). *Entomofauna* **22**: 433–444.
- 2004. Verbreitungsmuster bei Scoliiden – Ein Beitrag zu ihrer Evolution. *Beiträge der Hymenopterologen-Tagung Stuttgart 2004*: 7–9.
- 2005. Checkliste der Dolchwespen der Welt (Insecta: Hymenoptera, Scoliidae). *Bericht der Naturforschenden Gesellschaft Augsburg* **62**: 1–62.
- Oswald, J. D. 1990. Raphidioptera, pp. 154–163. In Grimaldi, D. A. (ed), *Insects from the Santana Formation, Lower Cretaceous, of Brazil. Bulletin of the American Museum of Natural History* **195**.
- O’Tool, C. and Raw, A. 1991. *Bees of the World*. London: Blanford.
- Ouvrard, D., Campbell, B. C., Bourgois, Th. and Chan, K. L. 2000. 18S rRNA secondary structure and phylogenetic position of Peloridiidae (Insecta, Hemiptera). *Molecular Phylogenetics and Evolution* **16**: 403–417.
- Pate, V. S. L. 1946. The generic names of the Sapygidae and their type species (Hymenoptera: Aculeata). *Entomological News* **57**: 219–221.
- 1947. Neotropical Sapygidae with a conspectus of the family (Hymenoptera: Aculeata). *Acta Zoologica Lilloana* **4**: 393–426.
- Peters, W. L. and Peters, J. G. 2000. Discovery of a new genus of Leptophlebiidae: Leptophlebiinae (Ephemeroptera) in Cretaceous amber from New Jersey, pp. 127–131. In Grimaldi, D. (ed), *Studies on Fossils in Amber, with Particular Reference to the Cretaceous of New Jersey*. Leiden: Backhuys.
- Petrulevičius, J. F. 1998. First hanging fly fossil from South America. *1st Paleoentomological Conference, Abstracts, 30 August–4 September 1998, Moscow*: 34.
- and Martins-Neto, R. G. 2001. A bitacid from the Santana Formation, Lower Cretaceous of Brazil. Legal, ethical and systematic aspects. *Acta Geological Leopoldensia* **24**: 125–127.
- and Martins-Neto, R. G. 2007. A new proterogomphid dragonfly from the Lower Cretaceous of Brazil. *Palaeontology* **50** (in press).
- Pierce, W. D. 1950. Fossil arthropods from onyx marble. *Bulletin of the Southern Californian Academy of Sciences* **49**: 101–104.
- 1951. Fossil arthropods from onyx marble. *Bulletin of the Southern Californian Academy of Sciences* **50**: 44–49.
- Pinto, J. D. 1989. A second new blattoid from the Cretaceous of Brazil. *Resumo das Comunicações do Congresso Brasileiro de Paleontologia* **11**: 295–300.
- and Ornellas, L. P. 1974. New Cretaceous Hemiptera (Insects) from Codo Formation – Northern Brazil. *Anais 28th Congresso Brasileiro do Geologia*: 289–304.
- Pinto, I. R. and Purper, I. 1986. A new blattoid from the Cretaceous of Brazil. *Pesquisas Instituto de Geociências Universidad Federal do Rio Grande do Sul* **18**: 5–10.
- Poinar, J. G. O. and Danforth, B. N. 2006. A fossil bee from Early Cretaceous Burmese amber. *Science* **314**: 614.
- Polegatto, C. M. and Zamboni, J. C. 2001. Inferences regarding the feeding behavior and morphoecological patterns of fossil mayfly nymphs (Insecta Ephemeroptera) from the Lower Cretaceous Santana Formation of northeastern Brazil. *Acta Geologica Leopoldensia* **24**: 145–160.

- Ponomarenko, A. G. 1977. Palaeozoic members of Megaloptera (Insecta). *Paleontologicheskiy Zhurnal* **1977**: 78–86.
- 1986. Scarabaeiformes incertae sedis, pp. 110–112. In Rasnitsyn, A. P. (ed), *Nasekomye v rannemelovyx ekosistemakh Zapadnoy Mongoli. Trudy Sovmestnaya Sovetsko – Mongol'skaya Paleontologicheskaya Ehkspeditsiya* **28**: 1–213.
 - 1987. New Mesozoic water beetles (Insecta, Coleoptera) from Asia. *Paleontological Journal* **21**: 79–92.
 - 1997. New beetles of the Family Cupedidae from the Mesozoic of Mongolia. Ommatini, Mesocupedini, Priacmini. *Paleontological Journal* **31**: 389–399.
 - 2000. New alderflies (Megaloptera: Parasailidae) and glosselytrodeans (Glosselytrodea: Glosselytridae) from the Permian of Mongolia. *Paleontological Journal, suppl. 3* **34**: 309–311.
 - 2003. Ecological evolution of beetles (Insecta: Coleoptera). *Acta Zoologica Cracoviensis* **46** (suppl. Fossil Insects): 319–328.
 - and Martínez-Declòs, X. 2000. New beetles (Insecta: Coleoptera) from the Lower Cretaceous of Spain. *Acta Geologica Hispanica* **35**: 47–52.
- Popham, E. J. 1990. Dermaptera, pp. 69–75. In Grimaldi, D. A. (ed), *Insects from the Santana Formation, Lower Cretaceous, of Brazil. Bulletin of the American Museum of Natural History* **195**.
- Popov, Yu. A. 1971. The historical development of Hemiptera infraorder Nepomorpha (Heteroptera). *Proceedings of the Paleontological Institute, Academy of Sciences, USSR* **129**.
- 1986. Peloridiina (=Coleorrhyncha) et Cimicina (=Heteroptera). Insects in Early Cretaceous ecosystems of western Mongolia. *Proceedings of the Joint Soviet-Mongolian Paleontological Expeditions, Nauka, Moscow* **28**: 50–83.
 - 1990. Description of fossil insects. True bugs. Cimicina. In *Late Mesozoic insects of Eastern Transbaikalia. Proceedings of the Paleontological Institute, Academy of Sciences, USSR, Nauka* **239**: 20–39.
 - and Shcherbakov, D. E. 1991. Mesozoic Peloridioidea and their ancestors (Insecta: Hemiptera, Coleorrhyncha). *Geologica et Palaeontologica* **25**: 215–235.
 - and Shcherbakov, D. E. 1996. Origin and evolution of the Coleorrhyncha as shown by the fossil record, pp. 9–30. In Schaefer, C. W. (ed), *Studies on Hemipteran Phylogeny*. Proceedings. Entomological Society of America. Lanham, MD: Thomas Say Publications in Entomology.
 - and Pinto, I. D. 2000. On some mesozoic burrower bugs (Heteroptera: Cydnidae). *Paleontological Journal* **34** (suppl. 3): 298–302.
 - , Dolling, W. R. and Whalley, P. E. S. 1994. British Upper Triassic and Lower Jurassic Heteroptera and Coeleorrhyncha (Insecta: Hemiptera). *Genus* **5**: 307–347.
- Prentice, M. A. 1998. *The Comparative Morphology and Phylogeny of Apoid Wasps (Hymenoptera: Apoidea)*. Dissertation thesis, University of California, Berkley.
- Pritchard, G., McKee, M. H., Pike, E. M., Scrimgeour, G. and Zloty, J. 1993. Did the first insects live in water or in air? *Biological Journal of the Linnean Society* **49**: 31–44.
- Prokop, J., Nel, A. and Hoch, I. 2005. Discovery of the oldest known Pterygota in the Lower Carboniferous of the Upper Silesian Basin in the Czech Republic (Insecta: Archaeorthoptera). *Geobios* **38**, 383–387.
- Pulawski, W. J. and Rasnitsyn, A. P. 2000. *Cretobestiola*, a replacement name for *Bestiola* Pulawski and Rasnitsyn, 1999 (Hymenoptera: Sphecidae). *Acta Geologica Hispanica* **35**: 53.
- Rasnitsyn, A. P. 1977. A new subfamily of scoliid wasps (Hymenoptera, Scoliidae, Proscoliinae). *Zoologicheskii Zhurnal* **56**: 522–529.

- 1986a. Review of the fossil Tiphidae, with description of a new species (Hymenoptera). *Psyche* **93**: 91–101.
- 1988. An outline of evolution of the hymenopterous insects (Order Vespida). *Oriental Insects* **22**: 115–145.
- 1993. Archaeoscoliinae, an extinct subfamily of scoliid wasps (Insecta: Vespa = Hymenoptera: Scoliidae). *Journal of Hymenoptera Research* **2**: 85–96.
- 2002. Infraclass Gryllones, pp. 254–256. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- and Martínez-Delclòs, X. 1999. New Cretaceous Scoliidae (Vespa = Hymenoptera) from the Lower Cretaceous of Spain and Brazil. *Cretaceous Research* **20**: 767–772.
- and Martínez-Delclòs, X. 2000. Wasps (Insecta: Vespa = Hymenoptera) from the Early Cretaceous of Spain. *Acta Geologica Hispanica* **35**: 65–95.
- and Pritykina, L. N. 2002. Superorder Libellulida Laicharting, 1781. Order Odonata Fabricius, 1792. The dragonflies, pp. 97–106. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- and Quicke, D. L. J. (eds), 2002. *History of Insects*. Dordrecht: Kluwer.
- and Zherikhin, V. V. 2002. 4. Appendix: alphabetic list of selected insect fossil sites 4.1. impression fossils, pp. 437–444. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- , Jarzembowski, E. A. and Ross, A. J. 1998. Wasps (Insecta: Vespa = Hymenoptera) from the Purbeck and Wealden (Lower Cretaceous) of Southern England and their biostratigraphical and paleoenvironmental significance. *Cretaceous Research* **19**: 329–391.
- , Pulawski, W. J. and Martínez-Delclòs, X. 1999. Cretaceous digger wasps of the new genus *Bestiola* Pulawski and Rasnitsyn (Hymenoptera: Sphecidae: Angarosphecinae). *Journal of Hymenoptera Research* **8**: 23–34.
- Rehn, A. C. 2003. Phylogenetic analysis of higher-level relationships of Odonata. *Systematic Entomology* **28**: 181–239.
- Ren, D. 1997. First record of fossil stick-insects from China with analyses of some palaeobiological features (Phasmatodea: Hagiphasmatidae fam. nov.). *Acta Zootaxonomica Sinica* **22**: 268–282.
- 1998. Late Jurassic [sic: Early Cretaceous] Brachycera from northeastern China (Insecta: Diptera). *Acta Zootaxonomica Sinica* **23**: 65–83.
- , Lu, L., Guo, Z., Ji, S. and Han, Y. 1995. [*Fauna and Stratigraphy of Jurassic-Cretaceous of Beijing and adjacent areas*]. Beijing: Seismic Publishing House.
- , Yin, J.-C. and Dou, W.-X. 1998a. Late Jurassic palaeontinids (Homoptera: Auchenorrhyncha) from the Hebei and Liaoning provinces in China. *Entomologica Sinica* **5**: 222–232.
- , Yin, J.-C. and Dou, W.-X. 1998b. New planthoppers and froghoppers from the Late Jurassic of Northeast China (Homoptera: Auchenorrhyncha). *Acta Zootaxonomica Sinica* **23**: 281–288.
- Rentz, D. C. F. 1996. *Grasshopper country: the abundant orthopteroid insects of Australia*. Sydney: New South Wales University Press.
- and Su, Y. N. 2003. Orthoptera (grasshoppers, locusts, katydids, crickets), pp. 827–839. In Resh, V. H. and Cardé, R. T. (eds), *Encyclopedia of Insects*. San Diego: Academic Press.
- Ribeiro, G. C. and Martins-Neto, R. G. 1999. A new Tipulidae (Insecta, Diptera) from the Santana Formation (Araripe Basin, Lower Cretaceous, Northeastern Brazil).

- Boletim do 5th Simpósio sobre o Cretáceo do Brasil (UNESP, São Paulo) 1999:* 207–212.
- and Krzemiski, W. 2000. New information on Limoniidae (Diptera: Tipulomorpha) from the Lower Cretaceous Santana Formation (northeastern Brazil). *Polskie Pismo Entomologiczne* **69**: 451–457.
- Richter, S. 2002. The Tetraconata concept: hexapod-crustacean relationships and the phylogeny of Crustacea. *Organisms Diversity & Evolution* **2**: 217–237.
- Riek, E. F. 1974. Upper Triassic insects from the Molteno Formation, South Africa. *Palaeontologica Africana* **17**: 19–31.
- Rohdendorf, B. B. 1977. The rationalization of names of higher taxa in zoology. *Paleontological Journal* **11**: 149–155.
- (ed.) 1991. *Fundamentals of Paleontology 9: Arthropoda, Tracheata, Chelicerata*. New Dehli: Amerind Publishers.
- Rohr, D. M., Boucot, A. J., Miller, J., and Abbott, M. 1986. Oldest termite nest from the Upper Cretaceous of west Texas. *Geology* **14**: 87–88.
- Romero, E. J. 1986. Fossil evidence regarding the evolution of *Nothofagus* Blume. *Annals of the Missouri Botanical Garden* **73**: 276–283.
- Roth, L. M. 1994. The beetle-mimicking cockroach genera *Prosoplecta* and *Areolaria*, with a description of *Tomeisneria furthi* gen. n., sp. n. (Blattellidae: Pseudophyllodromiinae). *Entomologica Scandinavica* **24**: 419–426.
- Rowell, C. H. F. and Flook, P. K. 1998. Phylogeny of the Caelifera and the Orthoptera as derived from ribosomal gene sequences. *Journal of Orthoptera Research* **7**: 147–156.
- Rozkosny, R. 2000. Stratiomyidae, pp. 387–411. In Papp, L. and Darvas, B. (eds), *Manual of Palaearctic Diptera*, vol. 2. Budapest: Science Herald.
- Ruf, M. L., Goodwyn, P. P. and Martins-Neto, R. G. 2005. New Heteroptera (Insecta) from the Santana Formation, Lower Cretaceous (Northeastern Brazil), with description of a new family and new taxa of Naucoridae and Gelastocoridae. *Gaea* **1**: 68–74.
- Rumbucher, K. 1995. Hemerobiidae (Insecta, Planipennia), eine bisher noch nicht entdeckte Familie der Santana Formation aus der brasilianischen Unterkreide. *Bericht der Naturforschenden Gesellschaft Augsburg* **55**: 46–61.
- Sakakibara, A. M. 1979. Sobre algumas espécies brasileiras de *Deois* Fennah, 1948 (Homoptera, Cercopidae). *Revista Brasileira de Biologia* **39**: 9–30.
- Samuelson, G. A. 1994. Pollen consumption and digestion by leaf beetles, pp. 179–183. In Jolivet, P. H., Cox, M. L. and Petitpierre, E. (eds), *Novel Aspects of the Biology of Chrysomelidae*. Dordrecht: Kluwer Academic Publishers.
- Santiago-Blay, J. A. 1994. Paleontology of leaf beetles, pp. 1–68. In Jolivet, H., Cox, M. L. and Petitpierre, E. (eds.), *Novel Aspects of the Biology of Chrysomelidae*. Dordrecht: Kluwer Academic Publishers.
- Santos, M. E. C. 1971. Um novo artrópodo da Formação Areão, Estado de Minas Gerais. *Annales Academiae Brasiliensis Ciencias* **43**: 415–420.
- Schaefer, C. W. (ed.). 1996. *Studies on Hemiptera Phylogeny*. Langham, MD: Thomas Say Publications in Entomology; Proceedings Entomological Society of America.
- Schlee, D. 1969. Morphologie und Symbiose; ihre Beweiskraft für die Verwandtschaftsbeziehungen der Coleorrhyncha (Insecta, Hemiptera). *Stuttgarter Beiträge zur Naturkunde, Serie C* **210**: 1–27.
- Schlüter, T. 1990. Fossil insect localities in Gondwanaland. *Entomologia Generalis* **15**: 61–76.
- Scholtz, C. H. and Chown, S. L. 1995. The evolution of habitat use and diet in the Scarabaeoidea: a phylogenetic approach, pp. 355–374. In Pakaluk, J. and Ślipiński,

- S. A. (eds), *Biology, Phylogeny, and Classification of Coleoptera*, vol. 1. Warsaw: Muzeum I Instytut Zoologii PAN.
- Schuh, R. T. and Slater, J. A. 1995. *True Bugs of the World (Hemiptera: Heteroptera) – Classification and Natural History*. Ithaca: Cornell University Press.
- Scudder, S. H. 1880. *The Devonian Insects of New Brunswick*. Anniversary Memoires of the Boston Society of Natural History.
- Sellick, J. T. C. 1994. Phasmida (stick insect) eggs from the Eocene of Oregon. *Palaeontology* **37**: 913–921.
- 1997. Descriptive terminology of the phasmid egg capsule, with an extended key to the phasmid genera based on egg structure. *Systematic Entomology* **22**: 97–122.
 - 1998. The micropylar plate of the eggs of Phasmida, with a survey of the range of plate form within the order. *Systematic Entomology* **23**: 203–228.
- Sharov, A. G. 1968. Phylogeny of the Orthopteroidea. *Trudy Palaeontologicheskogo Instituta, Nauk* **118**: 1–251.
- Shear, W. A., Bonamo, P. M., Grierson, J. D., et al. 1984. Early land animals in North America. *Science* **224**: 492–494.
- Shcherbakov, D. E. 1986 (1985). Nasekomye v rannemelovykh ekosistemach zapadnoi Mongoli. *Transactions of the joint Soviet–Mongolian Paleontological Expedition* **28**: 47–50.
- 1988. New cicadas (Cicadina) from the later Mesozoic of Transbaikalia. *Paleontological Journal* **4**: 52–63.
 - 1992. The earliest leafhoppers (Hemiptera: Karajassidae n. fam.) from the Jurassic of Karatau. *Neues Jahrbuche für Geologie und Paläontologie, Monatshefte* **1**: 39–51.
 - 1996. Origin and evolution of the Auchenorrhyncha as shown by the fossil record, pp. 31–45. In Schaefer, C. W. (ed.), *Studies on Hemipteran Phylogeny*. Lanham, MD: Thomas Say Entomological Publications. Entomological Society of America.
 - 2000. The most primitive white-flies (Hemiptera, Aleyrodidae, Bernaeinae subfam. nov.) from the Mesozoic of Asia and Burmese amber with an overview of Burmese amber hemipterans. *Bulletin of the Natural History Museum, Geology* **56**: 29–37.
 - 2002. Order Forficulida. The earwigs and protelytropterans, pp. 288–291, 298–301. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
 - 2004. Fossil adult and nymphal Tettigarctidae, pp. 71–73. In Kerzhner, I. M. (ed), *Third European Hemiptera Congress. St. Petersburg, June 8–11, 2004. Abstracts*. St. Petersburg: Russian Academy of Sciences, Department of Biological Sciences, St. Petersburg Scientific Centre, Zoological Institute.
 - and Popov, Yu. A. 2002. 2.2.1.2.5. Superorder Cimicidea Laicharting, 1781 Order Hemiptera Linné, 1758. The Bugs, Cicadas, Plantlice, Scale Insects, etc. (= Cimicida Laicharting, 1781, = Homoptera Leach, 1815 + Heteroptera Latreille, 1810), pp. 143–157. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- Silvestri, F. 1912. Die Thysanuren des Baltischen Bernsteins. *Schriftreihe der physikalisch-ökonomischen Gesellschaft Königsberg* **53**: 42–66.
- Sinitshenkova, N. D. 1975. Mayfly nymphs of the family Hexagenitidae (Insecta, Ephemeroptera). *Paleontologicheskii Zhurnal* **1975**: 82–86.
- 1986. Mayflies. Ephemera (=Ephemeroptera), In Rasnitsyn, A. P. (ed.), *Insects in the Early Cretaceous Ecosystems of the Western Mongolia. Transactions of the Joint Soviet–Mongolian Paleontological Expedition. Moscow. Nauka* **28**: 45–46.
 - 2000. The first fossil prosopistomatid mayfly from Burmese amber (Ephemeroptera: Prosopistomatidae). *Bulletin of the Natural History Museum of London (Geology)* **56**: 23–26.

- and Coram, R. A. 2002. The first mayfly from the Lower Cretaceous of southern England (Insecta: Ephemeroidea =Ephemeroptera). *Cretaceous Research* **23**: 461–463.
- Marchal-Papier, F., Grauvogel-Stamm, L. and Gall, J.-C. 2005. The Ephemeroidea (Insecta) from the Grès à Voltzia (early Middle Triassic) of the Vosges (NE France). *Paläontologische Zeitschrift* **79**: 377–397.
- Smith, P. E., Evensen, N. M., York, D. et al. 1995. Dates and rates in ancient lakes: ^{40}Ar - ^{39}Ar evidence for an Early Cretaceous age for the Jehol Group, Northeast China. *Canadian Journal of Earth Sciences* **32**: 1426–1431.
- Snodgrass, R. E. 1935. *Principles of Insect Morphology*. New York: McGraw-Hill.
- Sorensen, J. T., Campbell, B. C., Gill, R. J. and Steffen-Campbell, J. D. 1995. Non-monophyly of Auchenorrhyncha ("Homoptera"), based upon 18S rDNA phylogeny: eco-evolutionary and cladistic implications within pre-Heteropterodea Hemiptera (s.l.) and a proposal for new monophyletic sub-orders. *Pan-Pacific Entomologist* **71**: 31–60.
- Spahr, U. 1987. Ergänzungen und Berichtigungen zu R. Keilbachs Bibliographie und Liste der Bernsteinfossiliien-Ordnung Hymenoptera. *Stuttgarter Beiträge zur Naturkunde, Serie B* **127**: 1–121.
- Staniczek, A. H. 1997. The morphology of *Siphlaenigma janae* Penniket (Ephemeroptera, Siphlaenigmatidae), and its significance for the ground plan of the Baetoidea, pp. 536–549. In Landolt, P. and Sartori, M. (eds), *Ephemeroptera and Plecoptera: Biology-Ecology-Systematics*. Freibourg: Mauron + Tinguely & Lachat SA.
- 2000. The mandible of Silverfish (Insecta: Zygentoma) and Mayflies (Ephemeroptera): its morphology and phylogenetic significance. *Zoologischer Anzeiger* **239**: 147–178.
- 2003. New fossil mayflies from Dominican amber (Insecta: Ephemeroptera: Leptophlebiidae: Atalophlebiinae). *Stuttgarter Beiträge zur Naturkunde (B)* **341**: 1–22.
- and Bechly, G. 2002. First fossil record of the mayfly family Baetiscidae from Baltic amber (Insecta: Ephemeroptera). *Stuttgarter Beiträge zur Naturkunde (B)* **322**: 1–11.
- Stuckenberg, B. R. 2001. Pruning the tree: a critical review of classifications of the Homeodactyla (Diptera, Brachycera), with new perspectives and an alternative classification. *Studia Dipterologica* **8**: 3–41.
- Sturm, H. 1998. Erstnachweis fischchenartiger Insekten (Zygentoma, Insecta) für das Mesozoikum (Untere Kreide, Brasilien). *Senckenbergiana Lethaea* **78**: 135–140.
- and Mendes, L. F. 1998. Two new species of Nicoletiidae (Zygentoma, 'Apterygota', Insecta) in Dominican amber. *American Museum Novitates* **3226**: 1–11.
- and Machida, R. 2001. *Archaeognatha*. Handbuch der Biologie Band 37/ IV. Berlin: Akademische Verlagsgesellschaft Athenaion.
- Svensson, G. J. and Whiting, M. F. 2004. Phylogeny of Mantodea based on molecular data: evolution of a charismatic predator. *Systematic Entomology* **29**: 359–370.
- Sweet, M. H. 1996. Comparative external morphology of the pregenital abdomen of the Hemiptera, pp. 119–158. In Schaefer, C. W. (ed), *Studies on Hemiptera Phylogeny*. Langham, MD: Thomas Say Publications in Entomology; Proceedings Entomological Society of America.
- Swisher III, C. C., Wang, Y.-Q., Wang, X.-L., Wang, Xu, X. and Wang, Y. 1999. Cretaceous age for the feathered dinosaurs from Liaoning, China. *Nature* **400**: 58–61.
- Szwedo, J. 2002. Amber and amber inclusions of planthoppers, leafhoppers and their relatives (Hemiptera, Archaeorrhyncha et Clypaeorrhyncha), pp. 37–56. In Holzinger, W. (ed.), *Zikaden – Leafhoppers, Planthoppers and Cicadas (Insecta:*

- Hemiptera: Auchenorrhyncha). Denisia*, 04, Zugleich Kataloge, Neue Folge Nr. 176, Linz: OÖ. Landesmuseums.
- 2004a. A new genus and six new species of ground-dwelling leafhoppers from Chile and New Zealand (Hemiptera: Cicadomorpha: Myerslopiidae). *Zootaxa* **424**: 1–20.
 - 2004b. An annotated checklist of Myerslopiidae with notes on the distribution and origin of the group (Hemiptera: Cicadomorpha). *Zootaxa* **425**: 1–15.
 - 2004c. *Niryasaburnia* gen. nov. for ‘*Liburnia*’ burmitina Cockerell, 1917 from Burmese amber (Hemiptera, Fulgoromorpha: Achilidae). *Journal of Systematic Palaeontology* **2**: 105–107.
 - 2006. A new genus *Waghilde* gen. nov. representing a new tribe of the planthopper family Achilidae from the Eocene Baltic amber (Hemiptera: Fulgoromorpha). *Annales Zoologici* **56**: 167–174.
 - and Webb, M. D. 1999. First fossil representatives of Aetalionidae (Homoptera, Membracoidea) from Oligocene / Miocene ambers of the New World. *10th International Auchenorrhyncha Congress*, Abstracts of Talks and Posters, 6–10 September 1999, Cardiff, Wales: 3–4.
 - and Stroiński A. 2001. *Ptychogroehnia reducta* gen. and sp. n. of the fossil tribe Ptychoptilini from the Eocene Baltic amber (Hemiptera: Fulgoroidea: Achilidae). *Annales Zoologici* **51**: 95–101.
 - , Bourgoin, Th. and Lefebvre, F. 2004. *Fossil Planthoppers (Hemiptera: Fulgoromorpha) of the World. An Annotated Catalogue with Notes on Hemiptera Classification*. Warsaw: Studio 1.
 - Tan, J.-J., Ren, D. and Liu, M. 2005. New ommatids from the Late Jurassic of western Liaoning, China (Coleoptera: Archostemata). *Insect Science* **12**: 207–216.
 - Tanai, T. 1986. Phytogeographic and phylogenetic history of the genus *Nothofagus* Bl. (Fagaceae) in the Southern Hemisphere. *Journal of the Faculty of Science Hokkaido University* **21**: 505–582.
 - Thorne, B. L., Grimaldi, D. A. and Krishna, K. 2000. Early fossil history of the termites, pp. 77–94. In Abe, T., Bignell, D. E. and Higashi, M. (eds), *Termites – Evolution, Sociality, Symbiosis, Ecology*. Dordrecht: Kluwer.
 - and Traniello, J. F. A. 2002. Comparative social biology of basal taxa of ants and termites. *Annual Review of Entomology* **48**: 283–306.
 - Tilgner, E. H. 2000. The fossil record of Phasmida (Insecta: Neoptera). *Insect Systematics and Evolution* **31**: 473–480.
 - Tillyard, R. J. 1922. Mesozoic insects of Queensland. Nr. 9. Orthoptera and additioins to Protorthroptera, Odonata, Hemiptera and Plannipennia. *Proceedings of the Linnaean Society of New South Wales* **47**: 447–470.
 - 1932. Kansas Permian Insects. Part 15. The Order Plectoptera. *American Journal of Science* **223**: 97–134, 237–272.
 - Tobias, V. I. 1965. Contribution to the knowledge of the family Fedtschenkiidae (Hymenoptera, Sapygoidea). *Zoologicheskii Zhurnal* **XLIV**: 706–715.
 - Trueman, J. W. H. 1996. A preliminary cladistic analysis of odonate wing venation. *Odonatologica* **25**: 59–72.
 - Tshernova, O. A. 1962. Otrjad Ephemeroptera. Podenki, pp. 55–64. In Rodendorf, B. B. (ed), *Osnovy paleontologii*. Moscow: Akademija Nauk SSSR.
 - 1965. Some fossil mayflies (Ephemeroptera Misthodotidae) from Permian-beds of the Urals. *Entomologicheskoe Obozrenie* **44**: 353–361.
 - 1967. Mayfly of the Recent family in Jurassic deposits of Transbaikalia (Ephemeroptera, Siphlonuridae). *Entomologicheskoe Obozrenie* **46**: 322–326.

- 1970. On the classification of Fossil and Recent Ephemeroptera. *Entomological Review* **49**: 71–81.
- and Sinitshenkova, N. D. 1974. A new fossil genus and species of the mayfly family Hexagenitidae (Ephemeroptera) from the South of the European part of the USSR and its connection with Recent mayflies. *Entomologicheskoe Obozrenie* **53**: 130–136.
- Ueda, K. 1997. A new palaeontinid species from the Lower Cretaceous of Brazil (Homoptera: Palaeontinidae). *Bulletin of the Kitakyushu Museum of Natural History* **16**: 99–104.
- Von Brischke, D. 1886. Die Hymenopteren des Bernsteins. *Schriften der Naturforschenden Gesellschaft in Danzig (N.F.)* **6**: 278–279.
- Vršanský, P. 1999a. Lower Cretaceous Blattaria, pp. 167–176. In Vršanský, P. (ed), *Proceedings of the First Palaeoentomological Conference Moscow 1998*. Bratislava: AMBA Projects International.
- 1999b. Two new species of Blattaria (Insecta) from the Lower Cretaceous of Asia, with comments on the origin and phylogenetic position of the families Polyphagidae and Blattulidae. *Entomological Problems* **30**: 85–91.
- 2002. Origin and evolution of mantises. *AMBA Projekty* **6(1)**: 1–16.
- 2003a. Umenocoleoidea – an amazing lineage of aberrant insects (Insecta, Blattaria). *AMBA Projekty* **7**: 1–32.
- 2003b. Unique assemblage of Dictyoptera (Insecta – Blattaria, Mantodea, Isoptera) from the Lower Cretaceous of Bon Tsagaan Nuur in Mongolia. *Entomological Problems* **33(1–2)**, 119–151.
- 2004. Cretaceous Gondwanian Cockroaches (Insecta: Blattaria). *Entomological Problems* **34(1–2)**: 49–54.
- 2005. Evolúcia druhohorných švábov. Ph.D.-thesis. Geological Institute, Slovak Academy of Sciences. 40 pp + 14 supplements.
- , Vishniakova, V. N. and Rasnitsyn, A. P. 2002a. Order Blattida Latreille, 1810. The cockroaches, pp. 263–269. In Rasnitsyn, A. P. and Quicke, D. L. J. (eds), *History of Insects*. Dordrecht: Kluwer.
- , Mostovski, M. B., Bazylev, B. A. and Bugdaeva, E. 2002b. Early Cretaceous climate changes suggested on the basis of cockroach wing variations. *Proceedings of the XVII Congress of Carpathian–Balkan Geological Association*, 1–4 September 2002 (CD), Bratislava. *Geologica Carpathica* **33** (special issue).
- Vulcano, M. A. 1985. *Cretaceosimulium araripensis* gen. et sp. nov. da Chapada do Araripe, Ceará, Brasil (Diptera: Simuliidae). *Resumos do Congresso Brasileiro de Zoologia* **12**: 107.
- and Pereira, F. S. 1975. Cupesidae (Coleoptera). *Arquivos do Instituto Biológico São Paulo* **42**: 31–68.
- and Pereira, F. S. 1987. Entomofauna fóssil da Chapada do Araripe, Ceará, Brasil – *Cretaceimelittomoides cearensis* gen. nov., sp. nov. (Coleoptera: Pyrochroidae). *X Congreso Brasileiro de Paleontología, 1987, Rio de Janeiro. Resumo das Comunicações*: 27.
- Waller, A., Caussanel, C. and Jamet, C. 1996. Variation morphologique des pieces buccales chez quelques Dermapteres. *Bulletin de la Societe entomologique de France* **101**: 523–533.
- Wang, B., Zhang, H.-C. and Fang, Y. 2006a. Some Jurassic Palaeontinidae (Insecta, Hemiptera) from Daohugou, Inner Mongolia, China. *Palaeoworld* **15**: 115–125.
- , Zhang, H. C. and Fang, Y. 2006b. *Gansucossus*, a replacement name for *Yumenia* Hong, 1982 (Insecta, Hemiptera, Palaeontinidae), with description of a new genus. *Zootaxa* **1268**: 59–68.

- , Zhang, H. C., Fang, Y. and Zhang, Z. 2006c. A new genus and species of Palaeontinidae (Insecta: Hemiptera) from the Middle Jurassic of Daohugou, China. *Annales Zoologici* **56**: 757–762.
- , Zhang, H. C. and Fang, Y. 2007. Middle Jurassic Palaeontinidae (Insecta, Hemiptera) from Daohugou of China. *Alavesia* **1**: 89–104.
- Wang, W., Zhang, L., Zheng, S. et al. 2004. A new study on the stratotype and biostratigraphy of the Yixian stage in Yixian-Beipiao region, Liaoning – establishment and study of stratotypes of the Yixian stage. *Acta Geologica Sinica* **78**: 433–447.
- , Zhang, L., Zheng, S. et al. 2005. The age of the Yixian stage and the boundary of Jurassic-Cretaceous – the establishment and study of stratotypes of the Yixian stages. *Geological Review* **51**: 234–242.
- Wang, Y. and Ren, D. 2006. Middle Jurassic *Pseudococcus* fossils from Daohugou, Inner Mongolia in China (Homoptera, Palaeontinidae). *Acta Zootaxonomica Sinica* **31**: 289–293.
- , Ren, D. and Shih, C.-K. 2007a. Discovery of Middle Jurassic Palaeontinids from Inner Mongolia, China (Homoptera: Palaeontinidae). *Progress in Natural Science* **17**: 112–116.
- , Ren, D. and Shih, C.-K. 2007b. New discovery of palaeontinid fossils from the Middle Jurassic in Daohugou, Inner Mongolia (Homoptera, Palaeontinidae). *Science in China, Series D: Earth Sciences* **50**: 481–486.
- Wappler, T., Hinsken, S., Brocks, J. J., Wetzel, A. and Meyer, C. A. 2005. A fossil sawfly of the genus *Athalia* (Hymenoptera: Tenthredinidae) from the Eocene-Oligocene boundary of Altkirch, France. *Palevol* **4**: 7–16.
- Weber, H. 1933. *Lehrbuch der Entomologie*. Jena: Gustav Fischer.
- Weitschat, W. and Wichard, W. 1998. *Atlas der Pflanzen und Tiere im Baltischen Bernstein*. Munich: Pfeil.
- Westfall, M. J. 1980. Cretaceous dragonfly discovery in Brazil. *Selsysia* **9**: 22.
- Weyenbergh, H. 1874. Varia zoologica et paleontologica. *Periodico Zoologico, Sociedad Entomologica Argentina* **1**: 77–111.
- Whalley, P. E. S. 1983. A survey of recent and fossil cicadas (Insecta, Hemiptera–Homoptera) in Britain. *Bulletin of the British Museum of Natural History (Geology)* **37**: 139–47.
- and Jarzembowksi, E. A. 1985. Fossil insects from the Lithographic Limestone of Montsech (Late Jurassic–Early Cretaceous), Lerida Province, Spain. *Bulletin of the British Museum of Natural History (Geology)* **38**: 381–412.
- Wheeler, W. C., Whiting, M., Wheeler, Q. D., and Carpenter, J. M. 2001. The phylogeny of the extant hexapod orders. *Cladistics* **17**: 113–169, 403–404.
- Wickham, H. F. 1914. Twenty new Coleoptera from the Florissant shales. *Transactions of the Entomological Society of America* **40**: 257–270.
- Wieland, F. 2006. The cervical sclerites of Mantodea discussed in the context of dictyopteran phylogeny (Insecta: Dictyoptera). *Entomologische Abhandlungen* **63**: 51–76.
- Wighton, D. C. 1987. *Gomphaeschna obliqua* spec. nov., a new species of Gomphaeschninae from the Lower Cretaceous of northeastern Brazil (Anisoptera, Aeshnidae). *Odonatologica* **16**: 311–314.
- 1988. Odonate fossils from the Lower Cretaceous of Northeastern Brazil. *Proceeding of the 18th International Congress on Entomology*, Vancouver, Abstracts: 64.
- Wilf, P., Labandeira, C. C., Kress, W. J., et al. 2000. Timing the radiations of leaf beetles: hispines on gingers from Latest Cretaceous to Recent. *Science* **289**: 291–294.

- Willmann, R. 1978. *Mecoptera* (Insecta, Holometabola). In Westphal, F. (ed), *Fossilium Catalogus Animalia, Pars 124*. The Hague: Junk.
- 1987. The phylogenetic system of the Mecoptera. *Systematic Entomology* **12**: 519–524.
- 1989. Evolution und Phylogenetisches System der Mecoptera (Insecta: Holometabola). *Abhandlungen Senckenberg Naturforschungsinstitut, Gesellschaft* **544**: 1–153.
- 1994. Raphidiodea aus dem Lias und die Phylogenie der Kamelhalsfliegen (Insecta: Holometabola). *Paläontologische Zeitschrift* **68**: 167–197.
- 1999. The Upper Carboniferous *Lithoneura lameerei* (Insecta, Ephemeroptera?). *Paläontologische Zeitschrift* **73**: 289–302.
- 2003. Die phylogenetischen Beziehungen der Insecta: offene Fragen und Probleme. *Verhandlungen Westdeutscher Entomologentag* **2003**: 1–64.
- 2006. Die Stammgruppenvertreter der Ephemeroptera und ihre Systematik (Insecta: Ephemeroidea, = Ephemeroptera, = Plectoptera; Eintagsfliegen). *Species, Phylogeny & Evolution* **1**: 23–31.
- Wilson, E. O. 1971. *The Insect Societies*, pp. 31–37. Cambridge, MA: Harvard University Press.
- 1985. Ants from the Cretaceous and Eocene Amber of North America. *Psyche* **92**: 205–216.
- , Carpenter, F. M. and Brown, W. L. 1967. The first Mesozoic Ants, with the description of a new Subfamily. *Psyche* **74**: 1–19.
- Wilson, H. M., Dunlop, J. A., Martill, D. M. and Selden, P. A. 2000. Filling the Mesozoic gap: Terrestrial arthropods from the Crato Formation of Brazil. *Palaeontological Association 44th Annual Meeting, Abstracts and Programme*, University of Edinburgh: **38**.
- and Martill, D. M. 2001. A new Japygid Dipluran from the Lower Cretaceous of Brazil. *Palaeontology* **44**: 1025–1031.
- Woodley, N. E. 1986. Parhadrestiinae, a new subfamily for *Parhadrestia* James and *Cretaceogaster* Teskey (Diptera: Stratiomyidae). *Systematic Entomology* **11**: 377–387.
- Wygodzinsky, P. 1961. On a surviving representative of the Lepidotrichidae (Thysanura). *Annals of the Entomological Society of America* **54**: 621–627.
- Yang, C.-T. 2002. Preliminary thoughts on the phylogeny of Coleorrhyncha-Heteroptera (Hemiptera). *Formosan Entomologist* **22**: 297–305.
- Yang, C.-T. and Chang, T.-Y. 2000. *The External Genitalia of Hemiptera (Homoptera – Heteroptera)*. Taichung, Taiwan: Shih Way Publishers.
- Yeates, D. K. 1999. Congruence and controversy: toward a higher-level phylogeny of diptera. *Annual Review of Entomology* **44**: 397–428.
- and Irwin, M. E. 1996. Apioceridae (Insecta: Diptera): cladistic reappraisal and biogeography. *Zoological Journal of the Linnean Society* **116**: 247–301.
- Yeh, W.-B., Yang, C.-T. and Hui, C.-F. 2005. A molecular phylogeny of planthoppers (Hemiptera: Fulgoroidea) inferred from mitochondrial 16S rDNA sequences. *Zoological Studies* **44**: 519–535.
- Yildirim, E. and Gusenleitner, J. 2001. Sapygidae (Hymenoptera, Aculeata) of Turkey. *Linzer biologische Beiträge* **33**: 671–675.
- Zamboni, J. C. 2001. Contribution to the knowledge of the aquatic paleoentomofauna from Santana Formation (Araripe Basin, Lower Cretaceous, Northeast Brazil) with description of new taxa. *Acta Geologica Leopoldensia* **24**: 129–135.
- Zeuner, F. E. 1944. Notes on Eocene Homoptera from the Isle of Mull, Scotland. *Annals and Magazine of Natural History* **11**: 110–117.

- Zhang, H. C. 1997. Jurassic palaeotinids from Karamai, Xinjiang, with a discussion of Palaeotinidae (Homoptera: Palaeotinidae) in China. *Entomologia Sinica* **4**: 312–323.
- , Rasnitsyn, A. P. and Zhang, J. 2002. The oldest known scoliid wasps (Insecta, Hymenoptera, Scoliidae) from Jehol biota of western Liaoning, China. *Cretaceous Research* **23**: 77–86.
- Zhang, J.-F. 1994. Discovery of primitive earwigs (Insecta) from Late Jurassic of Laiyang, Shandong and its significance. *Acta Palaeontologica Sinica* **33**: 229–245.
- Zhang, Z.-J. 2002. New Early Cretaceous lalacid from Jingxi Basin of Beijing, China. *Acta Zootaxonomica Sinica* **27**: 20–23.
- Zherikhin, V. V. and Gratshev, V. G. 2004. Fossil Curculionoid Beetles (Coleoptera, Curculionoidea) from the Lower Cretaceous of Northeastern Brazil. *Paleontological Journal* **38**: 528–537.
- Zhou, C.-F. and Peters, J. G. 2003. The nymph of *Siphluriscus chinensis* and additional imaginal description: a living mayfly with Jurassic origins (Siphluriscidae new family: Ephemeroptera). *Florida Entomologist* **86**: 345–352.